

# Emerging Era of Biomolecular Membrane Simulations: Automated Physically-Justified Force Field Development and Quality-Evaluated Databanks

Hanne S. Antila,<sup>†</sup> Batuhan Kav,<sup>‡</sup> Markus S. Miettinen,<sup>¶,§</sup> Hector  
Martinez-Seara,<sup>||</sup> Pavel Jungwirth,<sup>\*,||</sup> and O. H. Samuli Ollila<sup>\*,⊥</sup>

<sup>†</sup>*Department of Biomaterials, Max Planck Institute of Colloids and Interfaces, 14424  
Potsdam, Germany*

<sup>‡</sup>*Institute of Biological Information Processing: Structural Biochemistry (IBI-7),  
Forschungszentrum Jülich, Wilhelm-Johnen-Str., 52425, Jülich, Germany*

<sup>¶</sup>*Computational Biology Unit, Department of Informatics, University of Bergen, 5008  
Bergen, Norway*

<sup>§</sup>*Department of Chemistry, University of Bergen, 5020 Bergen, Norway*

<sup>||</sup>*Institute of Organic Chemistry and Biochemistry, Czech Academy of Sciences, Flemingovo  
nam. 2, 160 00 Prague 6, Czech Republic*


<sup>⊥</sup>*Institute of Biotechnology, University of Helsinki, 00014 University of Helsinki*

E-mail: pavel.jungwirth@uochb.cas.cz; samuli.ollila@helsinki.fi

## Abstract

Molecular simulations of biological membranes and proxies thereof are entering a new era characterized by several key aspects. Progress starts with the realization that the outcome of the simulations can be only as good as the underlying force field and we actually need to know precisely how good or bad the results are. Therefore, standardized procedures for data quality evaluation are being established and will be applied to biomembrane simulations available in the literature. This provides the necessary basis and impetus for new force field development. Here, we propose to systematically build up physically well-justified models which effectively account for electronic polarization effects for all components of the biomembrane systems in aqueous environments. Such a massive task can only be achieved within a reasonable timescale by applying automated parameterization tools.

## Graphical TOC Entry



# Introduction

Classical molecular dynamics (MD) simulations of biomembranes are nowadays increasingly used in various applications, ranging from molecular cell biology to lipid nanoparticle formulations for vaccines.<sup>1-6</sup> Such applications are expected to increase further because MD simulations are becoming more accessible to a wider audience thanks to tools streamlining the simulation workflow, such as CHARMM-GUI<sup>7</sup> and Schrödinger-Maestro combined with Desmond,<sup>8</sup> and impressive performance of widely available GPU-computing hardware. At the same time, software and hardware development enable experts in the field to push the limits of MD simulations towards increasingly complex and large systems, such as cell membrane mimics<sup>9</sup> and virus particles.<sup>10</sup>

In both of these lines of development, it is increasingly important to understand the consequences arising from the approximations in the classical potential functions, i.e., force field, and simulation algorithms used in MD simulations. The new wave of users, consisting of practitioners with varying experience on methodological details, are often unaware of these limitations. At the same time, increasing the size, time scale, and complexity of the simulated systems may lead to the multiplication or cancellation of errors in the physical description of the systems. Furthermore, with the growing impact of MD simulations in biomedical and other applications, the potentially incorrect or misleading results will have increasingly severe consequences. Therefore, improving the MD methodology and the protocols to evaluate the quality of MD simulations is becoming more and more important.

Contrary to many other fields, there are no currently generally accepted standards when it comes to MD. In the field of structural biology, the Protein Data Bank (PDB) defines the quality measures for protein structures determined with different experimental techniques and enables access to vast amount of data for experts to evaluate the quality of each structure by themselves.<sup>11-14</sup> For intrinsically disordered proteins, belonging to the same class of unstructured molecules as lipids, such a databank is emerging,<sup>15</sup> but universal quality measures for conformational ensembles determined with different methods, including MD

simulations,<sup>16,17</sup> are yet to be defined. While importance of such databanks for MD simulations is widely recognized<sup>18–25</sup> and different kinds of approaches are emerging,<sup>25–32</sup> generally accepted protocols and best practices are still under active development.

At the same time, accumulating evidence on simulation artefacts arising from force fields (the sets of functions that approximate the physical interactions between atoms in MD simulations) raises the need for more accurate parameters and improved description of physics, particularly for lipids and disordered proteins.<sup>16,33–38</sup> Due to the semi-empirical nature of MD simulation force fields, a consistent parameterization represents a daunting task, often requiring simultaneous refinement of almost all interactions within a force field. To this end, several fully or semi-automatic approaches for force field parameterization are emerging.<sup>39–43</sup>

Here, we discuss current issues and ongoing activities in evaluating and improving biomolecular MD simulations with quality-evaluated databanks and automated force field development approaches. We focus on systems where atomic resolution description of lipid membranes, and lipid interactions with other molecules play a major role. Coarse grained simulations and other biomolecular systems are left outside the scope of this perspective.

## **Current issues in lipid membrane MD simulation quality**

### **Controversies in lipid membrane simulations**

Dozens of force fields and combinations thereof are available for simulating lipid membranes and their interactions with other molecules, such as proteins, drugs, and ions.<sup>9,44</sup> Even though the quality of force fields has been widely evaluated against experimental data during parameterization and in separate comparison studies,<sup>33–37,44</sup> defining unambiguously their overall quality still remains unclear, and several controversial results are reported for each in the literature, as exemplified in Table 1. Because universal quality measures for membrane simulations are not available, it is difficult to estimate, even for an expert, the reliability of a certain MD simulation result or to select the best force field for a specific application.

Therefore, lipid membrane MD simulations in their current state have to be used with great care to avoid misinterpreting experimental data or even launching questionable research directions inspired by artificial MD results.

As evident from Table 1, many controversies can be traced to interactions between lipids and other molecules including water. These disagreements are not surprising from the historical perspective, because early development of lipid bilayer simulations focused on correctly capturing membrane phase and area per lipid, which depend mainly on acyl chain properties,<sup>56,75</sup> while the glycerol backbone and headgroup and their interactions with ions and other molecules have gained attention only more recently.<sup>33,34,38,45–48,76</sup> Consequently, the hydrophobic acyl chain regions are commonly described quite well by MD simulations, while force fields struggle to correctly capture behaviour of the hydrophilic regions of membranes.<sup>35</sup>

In particular, the interactions between charged water-soluble species and membranes have been challenging for MD simulations. Canonical force fields tend to predict stronger binding of sodium and calcium to membranes than experiments.<sup>34,48</sup> One way to deal with this overbinding is to introduce an empirical correction denoted as NBFIX,<sup>48–50</sup> the gist of which is to increase the short-range repulsion between specific atoms and thus effectively increase the separation between them. Another approach, which is discussed in detail below, aims at including in a mean-field way the actual physical term responsible for the overbinding, i.e., the electronic polarization<sup>34,45–47</sup> Also, lipid–protein interactions depend on force field parameters,<sup>69–71,74,77</sup> which is particularly evident for charged residues<sup>37,78</sup> and their interactions with lipid headgroups.<sup>72,74</sup> While sodium and calcium ion binding to membranes with various compositions have been quantitatively evaluated using lipid headgroup order parameter data,<sup>34,38,47</sup> such comparison is not available for proteins or amino acids interacting with membranes. The evaluation of lipid–protein interactions has focused on comparing hydrophobicity scales,<sup>37,78,79</sup> NMR data<sup>77,80</sup> or crystallized lipid–protein complexes.<sup>81</sup> While hydrophobicity scales are often qualitatively reproduced by MD simulations, and proteins are usually oriented reasonably in membranes in these studies, charged residues often appear as

**Table 1: Examples of controversial MD simulation results reported in the literature. PC = phosphatidylcholine. ECC = electronic continuum correction. NBFIX = increasing the effective separation between specific atoms.<sup>48-50</sup> PS = phosphatidylserine. JM-A = the N-terminal portion of the juxtamembrane segments. EGF = epidermal growth factor. MC3 = dilinoleylmethyl-4-dimethylaminobutyrate. GPCR = G protein-coupled receptor**

Issue	Controversy	Proposed solutions
Cation binding to zwitterionic membrane	Strong binding of Na <sup>+</sup> and Ca <sup>2+</sup> ions to zwitterionic PC lipid headgroups observed in most force fields do not agree with NMR or other experimental data. <sup>34</sup>	Ion binding in force fields with ECC is more realistic. <sup>45-47</sup> NBFIX parameters between specific atoms have been introduced in CHARMM force field. <sup>48-50</sup>
Lipid-headgroup-cholesterol interactions	Direct interactions between cholesterol and lipid headgroups are observed in simulations with the Berger/Höltje force field combination, <sup>51</sup> while CHARMM36 predicts only weak preference for the nearest lipid headgroups to locate over cholesterol. <sup>52</sup>	NMR data agrees better with CHARMM36, suggesting that headgroups are rotating more freely in the cholesterol-containing membranes, <sup>53</sup> probably due to a less dense headgroup region.
Calcium binding position in PS lipid headgroup	CHARMM36 with NBFIX parameters <sup>48,49</sup> suggests that Ca <sup>2+</sup> ions interact only with the carboxylate group of PS lipids. <sup>54</sup> Without NBFIX the same force field indicates a significant binding affinity also to the phosphate region. <sup>55</sup> Berger lipid force field <sup>56,57</sup> suggests substantial binding also to the carbonyls in the acyl chains. <sup>58</sup>	In Amber-based force field with ECC, which has the most realistic ion binding behaviour when compared with NMR experiments, Ca <sup>2+</sup> binds approximately twice as likely to the carboxylate than to the phosphate moiety of PS, and binding only to acyl chain carbonyls is almost negligible. <sup>46</sup>
Langmuir monolayer simulations	Lipid monolayer can remain stable in a non-physical negative-surface-pressure state, and porates at too low area per lipid values. <sup>59,60</sup>	Water model with the correct surface tension and a sufficiently large periodic box are required to simulate lipid monolayers and quantitatively reproduce the surface pressure-area isotherms. <sup>61,62</sup>
Melting temperature of single-component lipid bilayers	Overestimated melting temperatures for single-component lipid bilayers have been reported, <sup>36</sup> while good agreement with experiments are found in other studies. <sup>63</sup>	Unresolved
Serotonin binding in different membrane phases	Serotonin binds preferentially to disordered phase in simulations and acyl chain order parameters of ordered phase are not affected, while its signal originates mostly from ordered state in NMR experiments. <sup>64,65</sup>	Unresolved
Accumulation of the ionizable MC3-lipids	Ionizable MC3-lipids, used in lipid-nanoparticle formulations for vaccines, accumulate in the bilayer center when the CHARMM36 force field is used, <sup>6</sup> while Slipids shows almost no accumulation. <sup>66</sup>	Unresolved
Lipid-protein interactions	Controversial results arising from force-field-dependent lipid-protein interactions have been reported for several proteins, such as the EGF receptor, <sup>67,68</sup> mellittin, <sup>69</sup> cytochrome P450, <sup>70</sup> P-Glycoprotein, <sup>71</sup> amyloid peptides, <sup>72,73</sup> and a GPCR. <sup>74</sup>	Unresolved

outliers or potential sources of discrepancies with experiments.<sup>37,77–80</sup> Also, the partitioning of small neutral molecules seems to agree with experiments qualitatively,<sup>82</sup> but experimental data is more scarce for charged molecules, and a systematic comparison of MD simulations with experiments is not available.<sup>83</sup>

In conclusion, we see that many controversial results from MD simulations originate from the description of polar and charged molecules or molecular groups, including water, and particularly their interactions with lipids. Therefore, we focus in the following on the effects of water models and electronic polarizability on interactions involving lipids. Other force field inaccuracies and issues related to simulation methodology, such as insufficient sampling or algorithms, are left for other accounts.

## Water models

As water plays at least an indirect role in all biological processes, much effort has been put to describe water properties in atomistic MD simulations accurately, and available parameters to model water have improved significantly over the years.<sup>84</sup> However, using the most realistic water model in biomolecular MD simulations is often not straightforward due to the semi-empirical nature of force fields: The change of water model would, in principle, require reparameterization of the complete force field. The change of the water model without such reparameterization compromises the consistency of parameters. Therefore, early water models, such as TIP3P and its derivatives,<sup>85</sup> are still widely used in biomolecular MD simulations despite their known inaccuracies.

The most notable shortcomings of water models (leading to known artefacts in biomolecular MD simulations, demonstrated in Fig. 1) are the underestimated surface tension and viscosity, as well as the dielectric constant (see Fig. 1 and the next section for a detailed discussion on the proper treatment of the dielectric response). Underestimated surface tension leads to incorrect phase behaviour and opening of pores at too low areas per molecule in lipid monolayer simulations.<sup>60–62</sup> Too low viscosity leads to overestimated rotational dynamics of




Figure 1: A) NMR spin relaxation times calculated from simulations of HpTonB-92 C-terminal domain using Amber ff99SB-ILDN force field with different water models compared with the experimental data. The results with TIP3P differ from experiments due to too fast rotational diffusion caused by the underestimated water viscosity, while results with TIP4P are close to experiments. Figure adapted with permission from Ref. 86. Copyright 2018 American Chemical Society. B) Surface pressure-area isotherms of DPPC lipid monolayer from CHARMM36 simulations with different water models and long-range Lennard-Jones treatments compared with experimental data.<sup>62</sup> Simulations with TIP3P based water models suffer from artefacts arising from underestimated surface tension, such as negative surface pressures and pore formation at too low areas, while simulations with better water models give almost quantitative agreement with experiments.<sup>60-62</sup> Figure adapted with permission from Ref. 62. Copyright 2022 American Chemical Society.

proteins as follows from comparison between simulations and experiments. The incorrect overall rotational diffusion can be corrected during analysis for folded proteins but not for disordered proteins or other biomolecules.<sup>17,86</sup> Furthermore, incorrect bulk water properties usually imply inaccuracies in interactions between water molecules, which are likely to cause other problems in biomolecular simulations. Indeed, the commonly observed over-compact ensembles of disordered proteins have been related to inaccuracies in water models.<sup>87,88</sup>

In applications where the water viscosity or surface tension are critical, the optimal solu-



tion is to use a more advanced water model,<sup>17,61,62</sup> which without careful reparameterization, however, could lead to inconsistencies with other parameters. In practise, such inconsistencies are observed to be less severe than issues arising from incorrect water properties.<sup>61,62,86</sup> In conclusion, we are convinced that using modern water models with correct surface tension, viscosity, dielectric response, and other physical properties is essential for future MD simulations of biomolecules.

## Lack of electronic polarizability

A general problem pertinent to the vast majority of membrane simulations is the lack of electronic polarizability effects. Indeed, standard force fields employed in membrane simulations are non-polarizable, meaning that atomic charges are fixed. This does not reflect reality where charged, or polar objects (i.e., ions or polar molecules) can polarize neighbouring molecules and thus effectively change their charge distributions, which if not considered results in several artifacts listed in Table 1. Explicitly polarizable force fields<sup>89–91</sup> aim at fixing this problem. However, they have not been widely employed in biomembrane simulations yet. In general, the limited use of explicitly polarizable force fields may be traced back to several factors. They include issues concerning their accurate parameterization (potentially leading to failures in reproducing experimental data; see below for more discussion), instability of the iterative algorithm (the so-called polarization catastrophe<sup>92</sup>), the need for specialized, not always user friendly software, and significantly increased computational costs.

Recently, a simple approach for including electronic polarization in a mean-field way in classical non-polarizable force fields has been suggested.<sup>81,93</sup> Within this method, denoted as the electronic continuum correction (ECC), electronic polarization effects are accounted for by scaling the charged groups by the inverse square root of the high-frequency dielectric constant of the surrounding medium, i.e., by the inverse of its refractive index. Note that the high-frequency dielectric constants of the relevant biomembrane distinct environments—aqueous solutions, membranes, and protein interiors—have values around 2, being very sim-

ilar to each other (unlike the total dielectric constants), which renders this essentially continuous approach applicable. In principle, charge scaling should be equivalent to immersing the whole system into a dielectric continuum with a dielectric constant of about 2. However, since our charge scaling is based on previously parameterized non-polarizable force fields, it has turned out to be prudent to scale only the charges of ions and ionic groups and not, e.g., of polar molecules like water. In other words, some degree of charge scaling may have been already included implicitly in the original force field when fitting it to experimental data. Note that partial charges are not experimental observables and, therefore, their values depend to some extent on the particular procedure adopted for their derivation. Also, the dielectric properties of common water models may not be entirely consistent with the ECC approach in the sense that their dielectric constants are larger than the experimental low-frequency value. Therefore, scaling factors lying anywhere between the inverse square root of the high-frequency dielectric constant and unity have been suggested.<sup>81</sup>

In the context of phospholipid membrane modelling, one of the key experimental manifestations of the deficiency of standard non-polarizable force fields is the substantially overestimated binding of biologically important cations such as sodium and calcium to lipid headgroups. This overbinding is true for membranes formed from both zwitterionic and anionic lipids, as exemplified by the a much larger effect of these cations on the lipid headgroup order parameters in the simulations than obtained from NMR measurements.<sup>34,38</sup> We have shown that charge scaling via the ECC approach fixes this problem in a physically well-justified way and brings the simulation results in a quantitative agreement with experiments. This grossly improved agreement is demonstrated for membranes containing PC, PS and PG lipids by plotting the dependence of the lipid headgroup order parameters on the aqueous salt concentration for sodium and calcium chloride.<sup>45–47</sup> The comparison to experiment of the results from ECC vs. standard non-polarizable force fields clearly shows the dramatic improvement charge scaling brings to the description of cation–headgroup interactions. The overestimated interaction between charged species within non-polarizable force fields has far-

reaching consequences in biomembranes, since transmembrane proteins,<sup>94</sup> the glycocalyx,<sup>95</sup> and other species that interact with the membrane<sup>96</sup> are rich in charged groups. All these issues arise from incorrect electrostatics, and we showed that charge scaling can to a large extent fix that.<sup>81</sup>

In the future, we plan to revisit the ECC force field development for biomolecules, starting the work from scratch, i.e., building up *de novo* a charge-scaled force field for water, ions, amino acids (and the corresponding peptides and proteins), phospholipids, nucleic acids, sugars, and other biologically relevant molecules. In this way, we will no longer be subject to potential overscaling due to reliance on existing force fields and inconsistent dielectric responses of water models. The challenge of building, from scratch, a new force field for biomembranes that is ECC-compatible would have been unsurmountable just a few years ago. Developing a new force field is still a monumental task involving several research groups and extensive iterations, where mistakes are corrected. In fact, there is a large amount of manual work involved which also explains why all well-known force fields used to simulate biomembranes started their development more than two decades ago. Nevertheless, the constant increase of available computational power and the surge of computational techniques that potentially allow considerable automatization of such tasks (e.g., machine learning,<sup>97</sup> see below) make the present an opportune time to attempt such an endeavor. If successful in our goal, we will eventually provide to the community an accurate force field that accounts for electronic polarizability in a mean-field charge scaling way such that it is fully equivalent to immersing the system into a dielectric continuum with the high-frequency dielectric constant.

# Current status of quality-evaluated databanks and automated force field development

## Quality-evaluated databanks

The importance of sharing MD simulation data following the FAIR principles<sup>98</sup> has been widely recognized,<sup>18–25</sup> and databanks are emerging.<sup>25–32</sup> The relevance of quality evaluation of simulation trajectories in databanks regarding technical details of simulations and accuracy of the underlying physical description of the system (force field) has become evident<sup>19,23,26</sup> and such quality evaluation has in some cases also been implemented.<sup>26,28</sup> However, straightforward quality comparisons between individual simulations or force fields within these databanks remain challenging.

The quality of lipid bilayer simulations has been quite extensively evaluated under various conditions,<sup>33–38,44,47,63,99–101</sup> yet determining unambiguously the overall accuracies of different force fields remains still unclear, as exemplified in Table 1. Within the NMRlipids Project ([nmrlipids.blogspot.fi](http://nmrlipids.blogspot.fi)), we have used the open collaboration approach, inspired by the Polymath Project<sup>102</sup> and open-source software development, to find lipid force fields that would correctly capture the lipid conformational ensembles and ion binding to membranes.<sup>33–35,38,47</sup> So far, we have used the data collected through the open collaboration to evaluate against NMR data the lipid headgroup conformational ensembles and ion binding affinities to membranes in the most commonly used force fields,<sup>33,34,38,47</sup> and to propose the *inverse conformational selection* model for lipid–protein interactions.<sup>47</sup> We are currently using the NMRlipids open collaboration approach and the data collected during the project to build an openly accessible databank of lipid bilayer simulations ([github.com/NMRlipids/Databank](https://github.com/NMRlipids/Databank)). The NMRlipids Databank will conduct by default automatic evaluations of simulation trajectory qualities against NMR and X-ray scattering data, and enable arbitrary user-defined automatic analyses over the vast sets of MD simulation data.

As an example application of using the NMRlipids Databank to evaluate conformational




Figure 2: Evaluation of conformational dynamics of lipid from the preliminary NMRlipids Databank<sup>103</sup> against the effective correlation time of C–H bond rotation,  $\tau_e$ , determined using spin-relaxation data.<sup>104</sup> Figure adapted with permission from Ref. 103. Copyright 2021 American Chemical Society.

dynamics of a large set of simulation data, Fig. 2 shows the effective rotational correlation times of lipid C–H bond vectors,  $\tau_e$ , calculated from all simulations in a preliminary version of the databank:<sup>103</sup> Only Slipids and CHARMM36 force fields are in line with the experimental data, predicting conformational timescales below 5 ns for all lipid segments. Other tested force fields exhibit significantly slower dynamics in the glycerol backbone and headgroup regions, but such discrepancies are not observed in the acyl chain region. This provides further support for the results from the NMRlipids Project indicating that CHARMM36 gives the most realistic description for the lipid headgroup region, while all force fields capture acyl chain region properties with reasonable accuracy.<sup>33,35,38,47</sup>

Another application of the NMRlipids Databank is exemplified in Fig. 3, which shows the area per lipid calculated from all POPC:POPS mixture simulations currently available in the NMRlipids Databank. The area per lipid relates to the lateral density and is commonly used to characterize membrane properties. Most force fields predict a decreased area per lipid (i.e., tighter membrane packing) with an increasing amount of charged POPS lipids in the membrane, yet the slope of the decrease depends on the force field parameters (Fig. 3). Since




Figure 3: Area per lipid calculated from all simulations of POPC:POPS mixtures with sodium counterions at 298 K currently available in the NMRlipids Databank (left) ordered according to the quality of the POPS *sn*-1 acyl chain (right). The number in parenthesis describes the negative 10-base logarithm of the average probability of the acyl chain  $\text{CH}_2$ -group order parameters to reside within experimental error in a simulation, thereby decreasing with increasing simulation quality. For reference, the values extracted from the experimental X-ray scattering data using the scattering density profile (SDP) model are shown in black.<sup>105,106</sup> The data used here have been originally reported in the NMRlipids publications.<sup>33,38</sup>

the area per lipid is not a directly measurable quantity in experiments, we quality-evaluate the results using the acyl chain order parameters. These depend on the membrane order, thereby being a good proxy for the area per lipid.<sup>35,107</sup> As all force fields predict acyl chain order parameters of pure POPC bilayers with reasonable accuracy, we focused on evaluating the quality of POPS acyl chain order parameters. The qualities of POPS simulations are ranked on the right-hand side of Fig. 3 based on C–H bond order parameters of the CH<sub>2</sub> groups in the *sn*-1 acyl chain of POPS. The quality of POPS simulations systematically decreases with decreasing area per lipid—from the GROMOS-CKP simulation with the largest area per lipid to the Berger simulation with the smallest area per lipid and the worst agreement with experiment. The quality evaluation based on acyl chain order parameters agrees with the comparison of area per lipid values extracted from X-ray scattering experiments using the SDP model<sup>105,106</sup> (Fig. 3). The force field dependent counterion binding affinities can explain the differences in the simulated POPS lipid bilayer areas: The bound ions screen the electrostatic repulsion between charged PS headgroups, thus leading to decreasing area per lipid with increasing binding affinity.<sup>38</sup> In conclusion, all force fields seem to overestimate to varying extent the counter ion binding affinity to POPS lipids, except for the GROMOS-CKP simulations. However, GROMOS-CKP predicts an unexpected minimum of area per lipid for a mixed PC:PS system, which cannot be verified with the currently available experimental data.

These examples demonstrate the power of the NMRlipids Databank in navigating the complex landscape of current lipid MD simulation quality. Notably, none of the current force fields overpasses others in all aspects of quality. For example, CHARMM36 force field best captured the conformational dynamics of lipids (Fig. 2) and the main differences between PC, PE, PG, and PS lipid headgroups, although not all order parameters were within the experimental accuracy.<sup>47</sup> At the same time, CHARMM36 is not the best option for applications where the packing of membranes containing PS lipids is relevant (Fig. 3). Consequently, the most reliable force field for a specific research question must be separately

selected. Without automatic quality evaluation enabled by the NMRlipids Databank or a similar resource, performing such evaluation separately for each application would become an overwhelming task.

As exemplified here and in previous studies, already the current version of the NMRlipids Databank helps to resolve issues with contradicting simulation results related, for example, to lipid conformational ensembles, dynamics, membrane packing and ion binding. However, more robust methods and experimental data are still needed for the evaluation of drug and amino acid binding to membranes, which is related to many known controversies listed in Table 1. Some NMR data from lipid headgroup order parameters are also available for such systems,<sup>108</sup> but quantitative comparison with simulations is more complicated for molecules with higher binding affinity. Furthermore, publicly available tools and quality measures enabling quantitative evaluation of MD simulations of other biomolecular systems than lipid bilayers, such as sugars or proteins, are not yet generally available—but there is certainly ongoing work towards providing them.<sup>16,17,109,110</sup> The flexible design of the NMRlipids Databank enables also inclusion of other systems than lipid bilayers in the future.

## **Automated force field development**

As discussed in the above sections, several significant simulation artefacts arise from fundamental issues in force field parameters, such as the lack of electronic polarizability or inaccuracies in water models. Correcting such issues is highly non-trivial because it requires systematic force field reparameterization. Building a new force field or extensively re-calibrating an existing one, is a daunting task owing to its sheer size (thousands of parameters); complexity (non-linear parameter dependencies with significant cross-correlations); the lock-in effect arising from the requirement of consistency with the decades of effort already invested in a particular force field; and the fact that such parameterization work is traditionally done by hand. Consequently, advances are made slowly, with efforts geared towards small yet important parts of the force fields with fewer target parameters, such as water



models.<sup>88,111,112</sup>

Recently, however, the rise of machine learning (ML) has made the automated building of predictive models with an extensive amount of parameters a common day practise. This widely lauded development—along with the ever-increasing computational resources and the growing availability of high-fidelity target data from experiments and from increasingly realistic ab initio calculations—seems to have significantly increased interest also in automated force field development. The promise is that high-throughput approaches would revolutionize force field development, making the currently unavoidable extensive human involvement and especially the incremental hand-tuning obsolete.

Automated force field development can be roughly divided into (1) approaches where a machine learning algorithm,<sup>97,113</sup> such as a neural network<sup>114–117</sup> or a kernel-based approach<sup>118</sup> acts as the force field, predicting the forces for a specific configuration of molecules (black-box potentials), and (2) approaches where automated optimization algorithms (including ML tools<sup>119</sup>) are used to find better parameters for existing force field functions<sup>40–43,120</sup> (gray-box potentials). Quality-evaluated open access databanks of MD trajectories can aid both of these approaches, as they guide the choice of optimization targets (by pinpointing the typical failures of existing MD force fields), establish quality standards, and can even provide training data for ML methods in force field parameterization.

**(1) Black-box potentials.** Typically these methods rely on the bottom-up approach, where a ML algorithm is trained with QM data of small molecules to extract a description of immediate environments of atoms, and to predict the atom-wise energies and forces. Such an algorithm is free from the shackles of the fixed functional forms of the classical force fields and is also intrinsically polarizable. However, it is somewhat limited in describing long-range interactions that cannot be cut off, such as electrostatics. Perhaps the most prominent examples have been those using neural networks, pioneered by Behler and Parrinello,<sup>114</sup> after which several architectures, such as ANI<sup>115,116</sup> and SchNet<sup>117</sup> have emerged.

Although such ML algorithms can be well-scalable to larger molecules<sup>121</sup> so far no truly

large-scale or complex (involving many molecular species) simulation has been conducted. Therefore, it is difficult to assess whether black-box potentials can accurately produce complex biomolecular simulation data in line with experiments, even though they do predict the potential energy surfaces, forces, and vibrational spectra with precision comparable to their training data for small molecules. They might suffer from a similar drawback that the classical force fields optimized using similar QM-based bottom-up approaches seem to have, i.e., not correctly probing the intermolecular interactions. Although subject to the same limitations in the types of systems (transferability and generalizability) and the level of theory used in generating the QM data for the training,<sup>122</sup> black-box potentials do not have the same flexibility for empirical error correction (e.g., by partial re-calibration to new experimental data) as classical force fields.

The computational cost of black-box potentials currently lies between those of ab initio and classical force fields.<sup>113,122,123</sup> This does not include the computational cost of training the black box, or creating the QM training data set, the size of which depends on the ML approach of choice. The cost associated with training data can be alleviated by exploiting or creating curated, open-access databanks and by using strategies, such as active learning, to sample the training data efficiently.<sup>116</sup>

**(2) Gray-box potentials.** There are two standard strategies for parameterizing a classical force field functions for MD simulations, either by 1) computing the function parameters from highly accurate QM calculations for (as large as possible) molecules and molecular fragments and piecing the model together bottom-up; or 2) optimizing the function parameters for molecules or multi-molecule systems directly against target experimental data.

In most of the currently used force fields, these two strategies have been used in conjunction with one another, usually by obtaining the starting point from QM calculations and then re-adjusting some parameters targeting experimental data. While force fields calibrated with this approach during the last decades have reached impressive performance, they also have severe flaws, as outlined in the sections above. The ever growing need for more accu-

rate force fields and more realistic description of the underlying physics such as electronic polarization is a strong incentive to speed up the parameterization process using automated approaches.

Note that the force field function parameters are traditionally thought to have a physical meaning, even when looked at individually (say, a dihedral stiffness or an atom size). When it comes to automated optimization of many parameters simultaneously, however, it can be argued that the physical meaning of individual parameters is blurred, and thus it make only limited sense to make such individual interpretations. Only the collection of parameters as a whole, to the extent they reproduce the targeted behaviour of the simulated system, has such a physical meaning.<sup>124</sup> To emphasize this point, we call such automatically optimized force fields here gray-box potentials.

Automated force field calibration has been done since the late 1990's<sup>125</sup> but has thus far mostly remained in the realm of small molecules; that said, also force fields for complex unstructured biomolecules, such as lipids, have been developed using automated approaches.<sup>43,126,127</sup> Gradient-based minimizations,<sup>41,43,128,129</sup> gradient-free methods such as Simplex<sup>130-132</sup> and evolutionary algorithms,<sup>39,40,42,120,132</sup> surrogate or metamodel strategies,<sup>39,112,133,134</sup> and their various combinations have been used for this purpose (Table 2). In addition to the extreme complexity of the underlying optimization problem, one of the main hurdles preventing such automated approaches seems to be the computational cost arising from the fact that the optimization loop requires running MD simulations on (possibly thousands) of force field candidates. This cost can be alleviated, for example, by thermodynamic reweighing of cached MD trajectories.<sup>43,135</sup> However, it appears that in some cases the gain from reweighing is smaller than the cost of trying the reweighing with insufficient data,<sup>136</sup> and that the reweighing only works reliably with minimal perturbations.

At the same time, the swarm of (possibly thousands) of force field candidates created as a side product of the optimisation loop is not just computational waste but high-quality raw material that can be upcycled to train machine learning algorithms (such as a neural

network), to act as a surrogate for the costly MD simulation in probing the optimization landscape,<sup>39,42,134</sup> or to create an expert (machine learning) model that directly predicts force field parameters from experimental data. Additionally, the major speed up by the next generation supercomputing architectures<sup>137</sup> may accelerate further automated parameterization procedures.

The increased automatization of force field parameter development with reduced human involvement underscores the importance of the quality and proper interpretation of target data. Optimal experimental target data have high accuracy and direct connection with simulations. Instead, data requiring approximate models to connect to physical properties should be relegated to an inferior role despite being easier for humans to interpret. For example, for lipid bilayers, the intuitively intelligible area per lipid can be determined only indirectly from experiments, and with relatively low accuracy. In contrast, the C–H bond order parameters from NMR or x-ray scattering form factors have a more direct connection with experiments and higher accuracy, thereby being better quantities to evaluate MD simulation quality.<sup>35</sup> However, care must be taken also when using order parameters as target data. For example, the lack of information on their sign or forking (two C–H bonds at the same carbon having different order parameters) can misguide the fitting algorithm to incorrect values, leading to significant deviation of the conformational ensemble from experiment, see Fig. 4. A similar situation may also appear if interfaces with incorrect surface pressures are used as fitting targets.<sup>62</sup> Nevertheless, the game-changing advantage of automated parameter optimization in such cases is the ease of repeating the fitting procedure with better target parameters when they become available.

Notably, even after carefully considering the various aspects of target data and fitting algorithms, one should be aware that fitting may not lead to a unique solution and that overfitting, e.g., by ignoring the inherent data uncertainty, targeting only one type of data, or severely overweighting one measurable, may lead to an incorrect ensemble being produced by the gray-box force field. Nevertheless, a force field that reproduces the high-quality exper-

**Table 2: Automated force field calibration studies. Complex biomolecular systems highlighted with light blue. ReaXFF = Reactive force field. EA = Evolutionary Algorithm. GM = Gradient-based Minimization. PSO = Particle Swarm Optimization. LS-SVR = Least-squares support-vector regression EXP = Experiments. QM = Quantum Mechanics. BI = Bayesian inference. SS = Systematic search. NN = Neural network. RW = Reweighting.**

authors	year	system	method	target
Gray-box potentials				
Hunger et al. <sup>125</sup>	1998	Mo(CO) <sub>3</sub> compounds	EA	EXP
Norrby and Liljefors <sup>138</sup>	1998	Ethane	GM	QM + EXP
Wang and Kollman <sup>40</sup>	2001	Small organic molecules	EA + SS	QM
Mostaghim et al. <sup>139</sup>	2004	Small organic molecules	EA + PSO	QM
Tafipolsky and Schmid <sup>140</sup>	2009	Metal organic frameworks	EA	QM
Liu et al. <sup>141</sup>	2011	Protein	LS-SVR	EXP
Handley and Deeth <sup>142</sup>	2011	Iron amide	EA	QM
Li and Hartke <sup>143</sup>	2013	ReaXFF for azobenzene	EA	QM
Dittner et al. <sup>144</sup>	2015	ReaXFF for SiOH	EA	EXP
Gao et al. <sup>145</sup>	2015	(CD)Benzene	EA	QM
Betz and Walker <sup>146</sup>	2015	Small organic molecules	EA	QM
McKiernan et al. <sup>126</sup>	2016	DPPC lipid	GM	EXP
Zahariev et al. <sup>120</sup>	2017	Phosphine oxides	EA + Simplex	QM
Wang et al. <sup>147</sup>	2017	Proteins	GM	QM
Dutta et al. <sup>112</sup>	2018	Water	BI	MD
Messerly et al. <sup>135</sup>	2018	Small hydrocarbons	Surrogate + RW	MD
Cesari et al. <sup>129</sup>	2019	RNA	GM + RW	EXP
Oliveira et al. <sup>148</sup>	2020	Haloalkanes	GM + RW	EXP
Krishnamoorthy et al. <sup>149</sup>	2020	2H-MoSe <sub>2</sub>	EA	QM
Oliveira and Hünenberger <sup>150</sup>	2021	Organic molecules	GM + RW	EXP
Yu et al. <sup>43,127</sup>	2021	Lipids	GM + RW	EXP
Yang et al. <sup>151</sup>	2021	ZrS <sub>2</sub>	EA	QM
Boothroyd et al. <sup>136</sup>	2021	Organic molecules	GM + RW	EXP
Wang et al. <sup>119</sup>	2021	Small organic molecules	NN + RW	QM
Befort et al. <sup>134</sup>	2021	Hydrofluorocarbons	Surrogate	QM + MD
Black-box potentials				
Morawietz and Behler <sup>152</sup>	2013	Water	NN	QM
Smith et al. <sup>153</sup>	2017	Organic molecules	NN	QM
Thaler and Zvadlav <sup>154</sup>	2021	Diamond	NN+RW	EXP+QM

imental data with a robust connection to simulations has a higher likelihood of representing the correct, underlying physical behaviour.

In conclusion, the black-box ML potentials and the automated parameterization of classical force fields to create gray-box potentials are promising tools for improving MD models, e.g., by introducing more accurate water models and including electronic polarization effects, improve thus their consistency with experiments, thus paving the way toward more realistic MD simulations of membranes and other biomolecular systems. However, although black-box ML potentials can reach QM-like precision, and methods like gradient-based optimization have consistently shown their advantages in improving the classical force fields, no single approach has yet emerged as clearly superior to tackle this task. For this to happen, practitioners need to keep pushing the applications towards larger and more complex systems. Considering this, we expect that the full potential of these approaches and the most successful methodologies of complex biomolecules will only be revealed in the coming years.

## Future perspectives

We are confident that a new era of biological membrane simulations is just about to emerge. It will be characterized by computational data with quantitative relation to existing experiments and providing high fidelity predictions thanks to the following key features: (i) Force fields will be developed systematically based on physically well-justified models, including accurate representation of the aqueous solvent and an effective account for electronic polarization, (ii) Force field development will employ automated parameterization tools making the whole procedure much more efficient and thus feasible within a reasonable timescale, and (iii) A standard procedure for quality evaluation will be established and performed for published simulation data. All these features will benefit from the creation of databanks of MD simulations where quality control against experiments is enforced.




Figure 4: (*top*) Headgroup and glycerol backbone C–H bond order parameters from standard CHARMM36 parameters are close to experiments, while the polarizable Drude-CHARMM predicts forking (inequivalent order parameters  $S_{CH}$  for the two C–H bonds in the same carbon) for the headgroup  $\alpha$  and  $\beta$  carbons. (*bottom*) Both force fields are close to experiments without the information on the signs and forking, probably because the average of absolute values over hydrogens attached to the same carbon were used as target parameters in the fitting procedure for the polarizable Drude-CHARMM.<sup>89</sup> C–H bonds are labelled as in Fig. 3. Data is taken from the NMRLipids VI project ([github.com/NMRLipids/NMRLipidsVIpolarizableFFs](https://github.com/NMRLipids/NMRLipidsVIpolarizableFFs)).

## Acknowledgement

PJ and HMS thank the Czech Science Foundation (EXPRO grant no. 19-26854X) for support. O.H.S.O. acknowledges the Academy of Finland (Academy Research Fellow Grants 315596, 319902, and 345631) for funding and CSC–IT Center for Science (Espoo, Finland) for computational resources. HSA thanks the Instrumentarium Science Foundation and Alexander von Humboldt foundation for financial support.

## References

- (1) Hedger, G.; Sansom, M. S. Lipid interaction sites on channels, transporters and receptors: Recent insights from molecular dynamics simulations. *Biochim. Biophys. Acta* **2016**, *1858*, 2390–2400.
- (2) Enkavi, G.; Javanainen, M.; Kulig, W.; Róg, T.; Vattulainen, I. Multiscale Simulations of Biological Membranes: The Challenge To Understand Biological Phenomena in a Living Substance. *Chem. Rev.* **2019**, *119*, 5607–5774.
- (3) Corradi, V.; Sejdiu, B. I.; Mesa-Galloso, H.; Abdizadeh, H.; Noskov, S. Y.; Markink, S. J.; Tieleman, D. P. Emerging Diversity in Lipid–Protein Interactions. *Chem. Rev.* **2019**, *119*, 5775–5848.
- (4) Reid, L. M.; Verma, C. S.; Essex, J. W. The role of molecular simulations in understanding the mechanisms of cell-penetrating peptides. *Drug Discov. Today* **2019**, *24*, 1821–1835.
- (5) Róg, T.; Girysh, M.; Bunker, A. Mechanistic Understanding from Molecular Dynamics in Pharmaceutical Research 2: Lipid Membrane in Drug Design. *Pharmaceuticals* **2021**, *14*.
- (6) Park, S.; Choi, Y. K.; Kim, S.; Lee, J.; Im, W. CHARMM-GUI Membrane Builder for



- Lipid Nanoparticles with Ionizable Cationic Lipids and PEGylated Lipids. *J. Chem. Inf. Model.* **2021**, *61*, 5192–5202.
- (7) Jo, S.; Kim, T.; Iyer, V. G.; Im, W. CHARMM-GUI: A web-based graphical user interface for CHARMM. *J. Comput. Chem.* **2008**, *29*, 1859–1865.
- (8) Bowers, K. J.; Chow, D. E.; Xu, H.; Dror, R. O.; Eastwood, M. P.; Gregersen, B. A.; Klepeis, J. L.; Kolossvary, I.; Moraes, M. A.; Sacerdoti, F. D. et al. Scalable Algorithms for Molecular Dynamics Simulations on Commodity Clusters. SC '06: Proceedings of the 2006 ACM/IEEE Conference on Supercomputing. 2006; pp 43–43.
- (9) Marrink, S. J.; Corradi, V.; Souza, P. C.; Ingólfsson, H. I.; Tieleman, D. P.; Sansom, M. S. Computational Modeling of Realistic Cell Membranes. *Chem. Rev.* **2019**, *119*, 6184–6226.
- (10) Tarasova, E.; Nerukh, D. All-Atom Molecular Dynamics Simulations of Whole Viruses. *J. Phys. Chem. Lett* **2018**, *9*, 5805–5809.
- (11) Berman, H.; Henrick, K.; Nakamura, H. Announcing the worldwide Protein Data Bank. *Nat. Struct. Mol. Biol.* **2003**, *10*, 980–980.
- (12) Read, R. J.; Adams, P. D.; Arendall III, W. B.; Brunger, A. T.; Emsley, P.; Joosten, R. P.; Kleywegt, G. J.; Krissinel, E. B.; Lütke, T.; Otwinowski, Z. et al. A New Generation of Crystallographic Validation Tools for the Protein Data Bank. *Structure* **2011**, *19*, 1395–1412.
- (13) Montelione, G. T.; Nilges, M.; Bax, A.; Güntert, P.; Herrmann, T.; Richardson, J. S.; Schwieters, C. D.; Vranken, W. F.; Vuister, G. W.; Wishart, D. S. et al. Recommendations of the wwPDB NMR Validation Task Force. *Structure* **2013**, *21*, 1563–1570.
- (14) Henderson, R.; Sali, A.; Baker, M. L.; Carragher, B.; Devkota, B.; Downing, K. H.;

- Egelman, E. H.; Feng, Z.; Frank, J.; Grigorieff, N. et al. Outcome of the First Electron Microscopy Validation Task Force Meeting. *Structure* **2012**, *20*, 205–214.
- (15) Lazar, T.; Martínez-Pérez, E.; Quaglia, F.; Hatos, A.; Chemes, L.; Iserle, J. A.; Méndez, N. A.; Garrone, N. A.; Saldaño, T.; Marchetti, J. et al. PED in 2021: a major update of the protein ensemble database for intrinsically disordered proteins. *Nucleic Acids Res.* **2020**, *49*, D404–D411.
- (16) Robustelli, P.; Piana, S.; Shaw, D. E. Developing a molecular dynamics force field for both folded and disordered protein states. *Proc. Natl. Acad. Sci.* **2018**, *115*, E4758–E4766.
- (17) Virtanen, S. I.; Kiirikki, A. M.; Mikula, K. M.; Iwai, H.; Ollila, O. H. S. Heterogeneous dynamics in partially disordered proteins. *Phys. Chem. Chem. Phys.* **2020**, *22*, 21185–21196.
- (18) Feig, M.; Abdullah, M.; Johnsson, L.; Pettitt, B. Large scale distributed data repository: design of a molecular dynamics trajectory database. *Future Gener. Comput. Syst.* **1999**, *16*, 101–110.
- (19) Tai, K.; Murdock, S.; Wu, B.; Ng, M. H.; Johnston, S.; Fangohr, H.; Cox, S. J.; Jeffreys, P.; Essex, J. W.; Sansom, M. S. BioSimGrid: towards a worldwide repository for biomolecular simulations. *Org. Biomol. Chem.* **2004**, *2*, 3219–3221.
- (20) Silva, C. G.; Ostropytsky, V.; Loureiro-Ferreira, N.; Berrar, D.; Swain, M.; Dubitzky, W.; Brito, R. M. M. P-found: The Protein Folding and Unfolding Simulation Repository. 2006 IEEE Symposium on Computational Intelligence and Bioinformatics and Computational Biology. 2006; pp 1–8.
- (21) Abraham, M.; Apostolov, R.; Barnoud, J.; Bauer, P.; Blau, C.; Bonvin, A. M.; Chavent, M.; Chodera, J.; Čondić Jurčić, K.; Delemotte, L. et al. Sharing Data from Molecular Simulations. *J. Chem. Inf. Model.* **2019**, *59*, 4093–4099.

- (22) Hildebrand, P. W.; Rose, A. S.; Tiemann, J. K. Bringing Molecular Dynamics Simulation Data into View. *Trends Biochem. Sci.* **2019**, *44*, 902–913.
- (23) Hospital, A.; Battistini, F.; Soliva, R.; Gelpí, J. L.; Orozco, M. Surviving the deluge of biosimulation data. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **2020**, *10*, e1449.
- (24) Abriata, L. A.; Lepore, R.; Dal Peraro, M. About the need to make computational models of biological macromolecules available and discoverable. *Bioinformatics* **2020**, *36*, 2952–2954.
- (25) Rodríguez-Espigares, I.; Torrens-Fontanals, M.; Tiemann, J. K. S.; Aranda-García, D.; Ramírez-Anguita, J. M.; Stepniewski, T. M.; Worp, N.; Varela-Rial, A.; Morales-Pastor, A.; Medel-Lacruz, B. et al. GPCRmd uncovers the dynamics of the 3D-GPCRome. *Nat. Methods* **2020**, *17*, 777–787.
- (26) Meyer, T.; D’Abramo, M.; Hospital, A.; Rueda, M.; Ferrer-Costa, C.; Pérez, A.; Carrillo, O.; Camps, J.; Fenollosa, C.; Repchevsky, D. et al. MoDEL (Molecular Dynamics Extended Library): A Database of Atomistic Molecular Dynamics Trajectories. *Structure* **2010**, *18*, 1399–1409.
- (27) van der Kamp, M. W.; Schaeffer, R. D.; Jonsson, A. L.; Scouras, A. D.; Simms, A. M.; Toofanny, R. D.; Benson, N. C.; Anderson, P. C.; Merkle, E. D.; Rysavy, S. et al. Dynameomics: A Comprehensive Database of Protein Dynamics. *Structure* **2010**, *18*, 423–435.
- (28) Hospital, A.; Andrio, P.; Cugnasco, C.; Codo, L.; Becerra, Y.; Dans, P. D.; Battistini, F.; Torres, J.; Goñi, R.; Orozco, M. et al. BIGNASim: a NoSQL database structure and analysis portal for nucleic acids simulation data. *Nucleic Acids Res.* **2016**, *44*, D272–D278.
- (29) Mixcoha, E.; Rosende, R.; Garcia-Fandino, R.; Piñeiro, A. Cyclo-lib: a database of

- computational molecular dynamics simulations of cyclodextrins. *Bioinformatics* **2016**, *32*, 3371–3373.
- (30) Newport, T. D.; Sansom, M. S.; Stansfeld, P. J. The MemProtMD database: a resource for membrane-embedded protein structures and their lipid interactions. *Nucleic Acids Res.* **2018**, *47*, D390–D397.
- (31) Bekker, G.-J.; Kawabata, T.; Kurisu, G. The Biological Structure Model Archive (BSM-Arc): an archive for in silico models and simulations. *Biophys. Rev.* **2020**, *12*, 371–375.
- (32) Suarez-Leston, F.; Calvelo, M.; Tolufashe, G. F.; Muñoz, A.; Veleiro, U.; Porto, C.; Bastos, M.; Ángel Piñeiro,; Garcia-Fandino, R. SuPepMem: A database of innate immune system peptides and their cell membrane interactions. *Comput. Struct. Biotechnol. J.* **2022**, *20*, 874–881.
- (33) Botan, A.; Favela-Rosales, F.; Fuchs, P. F. J.; Javanainen, M.; Kanduč, M.; Kulig, W.; Lamberg, A.; Loison, C.; Lyubartsev, A.; Miettinen, M. S. et al. Toward Atomistic Resolution Structure of Phosphatidylcholine Headgroup and Glycerol Backbone at Different Ambient Conditions. *J. Phys. Chem. B* **2015**, *119*, 15075–15088.
- (34) Catte, A.; Girych, M.; Javanainen, M.; Loison, C.; Melcr, J.; Miettinen, M. S.; Monticelli, L.; Määttä, J.; Oganessian, V. S.; Ollila, O. H. S. et al. Molecular electrometer and binding of cations to phospholipid bilayers. *Phys. Chem. Chem. Phys.* **2016**, *18*, 32560–32569.
- (35) Ollila, O. S.; Pabst, G. Atomistic resolution structure and dynamics of lipid bilayers in simulations and experiments. *Biochim. Biophys. Acta* **2016**, *1858*, 2512 – 2528.
- (36) Pluhackova, K.; Kirsch, S. A.; Han, J.; Sun, L.; Jiang, Z.; Unruh, T.; Böckmann, R. A. A Critical Comparison of Biomembrane Force Fields: Structure and Dynamics of Model DMPC, POPC, and POPE Bilayers. *J. Phys. Chem. B* **2016**, *120*, 3888–3903.

- (37) Sandoval-Perez, A.; Pluhackova, K.; Böckmann, R. A. Critical Comparison of Biomembrane Force Fields: Protein–Lipid Interactions at the Membrane Interface. *J. Chem. Theory Comput* **2017**, *13*, 2310–2321.
- (38) Antila, H.; Buslaev, P.; Favela-Rosales, F.; Ferreira, T. M.; Gushchin, I.; Javanainen, M.; Kav, B.; Madsen, J. J.; Melcr, J.; Miettinen, M. S. et al. Headgroup Structure and Cation Binding in Phosphatidylserine Lipid Bilayers. *J. Phys. Chem. B* **2019**, *123*, 9066–9079.
- (39) Hunger, J.; Huttner, G. Optimization and analysis of force field parameters by combination of genetic algorithms and neural networks. *J. Comput. Chem.* **1999**, *20*, 455–471.
- (40) Wang, J.; Kollman, P. A. Automatic parameterization of force field by systematic search and genetic algorithms. *J. Comput. Chem.* **2001**, *22*, 1219–1228.
- (41) Wang, L.-P.; Martinez, T. J.; Pande, V. S. Building Force Fields: An Automatic, Systematic, and Reproducible Approach. *J. Phys. Chem. Lett.* **2014**, *5*, 1885–1891, PMID: 26273869.
- (42) Krämer, A.; Hülsmann, M.; Köddermann, T.; Reith, D. Automated parameterization of intermolecular pair potentials using global optimization techniques. *Comput. Phys. Commun* **2014**, *185*, 3228–3239.
- (43) Yu, Y.; Krämer, A.; Venable, R. M.; Simmonett, A. C.; MacKerell, A. D.; Klauda, J. B.; Pastor, R. W.; Brooks, B. R. Semi-automated Optimization of the CHARMM36 Lipid Force Field to Include Explicit Treatment of Long-Range Dispersion. *J. Chem. Theory Comput* **2021**, *17*, 1562–1580.
- (44) Leonard, A. N.; Wang, E.; Monje-Galvan, V.; Klauda, J. B. Developing and Testing of Lipid Force Fields with Applications to Modeling Cellular Membranes. *Chem. Rev.* **2019**, *119*, 6227–6269.

- (45) Melcr, J.; Martinez-Seara, H.; Nencini, R.; Kolafa, J.; Jungwirth, P.; Ollila, O. H. S. Accurate Binding of Sodium and Calcium to a POPC Bilayer by Effective Inclusion of Electronic Polarization. *J. Phys. Chem. B* **2018**, *122*, 4546–4557.
- (46) Melcr, J.; Ferreira, T. M.; Jungwirth, P.; Ollila, O. H. S. Improved Cation Binding to Lipid Bilayers with Negatively Charged POPS by Effective Inclusion of Electronic Polarization. *J. Chem. Theo. Comput.* **2020**, *16*, 738–748.
- (47) Bacle, A.; Buslaev, P.; Garcia-Fandino, R.; Favela-Rosales, F.; Mendes Ferreira, T.; Fuchs, P. F. J.; Gushchin, I.; Javanainen, M.; Kiirikki, A. M.; Madsen, J. J. et al. Inverse Conformational Selection in Lipid–Protein Binding. *J. Am. Chem. Soc.* **2021**, *143*, 13701–13709.
- (48) Venable, R. M.; Luo, Y.; Gawrisch, K.; Roux, B.; Pastor, R. W. Simulations of Anionic Lipid Membranes: Development of Interaction-Specific Ion Parameters and Validation Using NMR Data. *J. Phys. Chem. B* **2013**, *117*, 10183–10192.
- (49) Kim, S.; Patel, D.; Park, S.; Slusky, J.; Klauda, J.; Widmalm, G.; Im, W. Bilayer Properties of Lipid A from Various Gram-Negative Bacteria. *Biophys. J.* **2016**, *111*, 1750 – 1760.
- (50) Han, K.; Venable, R. M.; Bryant, A.-M.; Legacy, C. J.; Shen, R.; Li, H.; Roux, B.; Gericke, A.; Pastor, R. W. Graph–Theoretic Analysis of Monomethyl Phosphate Clustering in Ionic Solutions. *J. Phys. Chem. B* **2018**, *122*, 1484–1494.
- (51) Alwarawrah, M.; Dai, J.; Huang, J. Modification of Lipid Bilayer Structure by Diacylglycerol: A Comparative Study of Diacylglycerol and Cholesterol. *J. Chem. Theor. Comput.* **2012**, *8*, 749–758.
- (52) Leeb, F.; Maibaum, L. Spatially Resolving the Condensing Effect of Cholesterol in Lipid Bilayers. *Biophys. J.* **2018**, *115*, 2179–2188.

- (53) Antila, H. S.; Wurl, A.; Ollila, O. H. S.; Miettinen, M. S.; Ferreira, T. M. Rotational decoupling between the hydrophilic and hydrophobic regions in lipid membranes. *Biophys. J.* **2021**,
- (54) Valentine, M. L.; Cardenas, A. E.; Elber, R.; Baiz, C. R. Physiological Calcium Concentrations Slow Dynamics at the Lipid-Water Interface. *Biophys. J.* **2018**, *115*, 1541 – 1551.
- (55) Hallock, M. J.; Greenwood, A. I.; Wang, Y.; Morrissey, J. H.; Tajkhorshid, E.; Rienstra, C. M.; Pogorelov, T. V. Calcium-Induced Lipid Nanocluster Structures: Sculpturing of the Plasma Membrane. *Biochemistry* **2018**, *57*, 6897–6905.
- (56) Berger, O.; Edholm, O.; Jähnig, F. Molecular dynamics simulations of a fluid bilayer of dipalmitoylphosphatidylcholine at full hydration, constant pressure, and constant temperature. *Biophys. J.* **1997**, *72*, 2002 – 2013.
- (57) Mukhopadhyay, P.; Monticelli, L.; Tieleman, D. P. Molecular Dynamics Simulation of a Palmitoyl-Oleoyl Phosphatidylserine Bilayer with Na<sup>+</sup> Counterions and NaCl. *Biophys. J.* **2004**, *86*, 1601 – 1609.
- (58) Melcrová, A.; Pokorna, S.; Pullanchery, S.; Kohagen, M.; Jurkiewicz, P.; Hof, M.; Jungwirth, P.; Cremer, P. S.; Cwiklik, L. The complex nature of calcium cation interactions with phospholipid bilayers. *Sci. Reports* **2016**, *6*, 38035.
- (59) Baoukina, S.; Monticelli, L.; Marrink, S. J.; Tieleman, D. P. Pressure–Area Isotherm of a Lipid Monolayer from Molecular Dynamics Simulations. *Langmuir* **2007**, *23*, 12617–12623.
- (60) Lamberg, A.; Ollila, O. H. S. Comment on “Structural Properties of POPC Monolayers under Lateral Compression: Computer Simulations Analysis”. *Langmuir* **2015**, *31*, 886–887.

- (61) Javanainen, M.; Lamberg, A.; Cwiklik, L.; Vattulainen, I.; Ollila, O. H. S. Atomistic Model for Nearly Quantitative Simulations of Langmuir Monolayers. *Langmuir* **2018**, *34*, 2565–2572.
- (62) Tempra, C.; Ollila, O. H. S.; Javanainen, M. Accurate Simulations of Lipid Monolayers Require a Water Model with Correct Surface Tension. *J. Chem. Theory Comput* **2022**, *18*, 1862–1869.
- (63) Khakbaz, P.; Klauda, J. B. Investigation of phase transitions of saturated phosphocholine lipid bilayers via molecular dynamics simulations. *Biochim. Biophys. Acta* **2018**, *1860*, 1489–1501.
- (64) Engberg, O.; Bochicchio, A.; Brandner, A. F.; Gupta, A.; Dey, S.; Böckmann, R. A.; Maiti, S.; Huster, D. Serotonin Alters the Phase Equilibrium of a Ternary Mixture of Phospholipids and Cholesterol. *Front. Physiol.* **2020**, *11*, 1389.
- (65) Bochicchio, A.; Brandner, A. F.; Engberg, O.; Huster, D.; Böckmann, R. A. Spontaneous Membrane Nanodomain Formation in the Absence or Presence of the Neurotransmitter Serotonin. *Front. Cell Dev. Biol.* **2020**, *8*, 1440.
- (66) Ermilova, I.; Swenson, J. DOPC versus DOPE as a helper lipid for gene-therapies: molecular dynamics simulations with DLin-MC3-DMA. *Phys. Chem. Chem. Phys.* **2020**, *22*, 28256–28268.
- (67) Arkhipov, A.; Shan, Y.; Das, R.; Endres, N.; Eastwood, M.; Wemmer, D.; Kuriyan, J.; Shaw, D. Architecture and Membrane Interactions of the EGF Receptor. *Cell* **2013**, *152*, 557 – 569.
- (68) Kaszuba, K.; Grzybek, M.; Orlowski, A.; Danne, R.; Róg, T.; Simons, K.; Coskun, U.; Vattulainen, I. N-Glycosylation as determinant of epidermal growth factor receptor conformation in membranes. *Proc. Natl. Acad. Sci. U.S.A.* **2015**, *112*, 4334–4339.



- (69) Fox, S. J.; Lakshminarayanan, R.; Beuerman, R. W.; Li, J.; Verma, C. S. Conformational Transitions of Melittin between Aqueous and Lipid Phases: Comparison of Simulations with Experiments. *J. Phys. Chem. B* **2018**, *122*, 8698–8705.
- (70) Mustafa, G.; Nandekar, P. P.; Mukherjee, G.; Bruce, N. J.; Wade, R. C. The Effect of Force-Field Parameters on Cytochrome P450-Membrane Interactions: Structure and Dynamics. *Sci. Rep.* **2020**, *10*, 7284.
- (71) Wang, L.; O’Mara, M. L. Effect of the Force Field on Molecular Dynamics Simulations of the Multidrug Efflux Protein P-Glycoprotein. *J. Chem. Theory Comput* **2021**, *17*, 6491–6508.
- (72) Lockhart, C.; Smith, A. K.; Klimov, D. K. Three Popular Force Fields Predict Consensus Mechanism of Amyloid  $\beta$  Peptide Binding to the Dimyristoylglycerophosphocholine Bilayer. *J. Chem. Inf. Model.* **2020**, *60*, 2282–2293.
- (73) Yang, Y.; Jalali, S.; Nilsson, B. L.; Dias, C. L. Binding Mechanisms of Amyloid-like Peptides to Lipid Bilayers and Effects of Divalent Cations. *ACS Chem. Neurosci.* **2021**, *12*, 2027–2035.
- (74) Mahmood, M. I.; Yamashita, T. Influence of Lipid Bilayer on the GPCR Structure: Comparison of All-Atom Lipid Force Fields. *Bull. Chem. Soc. Jpn.* **2021**, *94*, 2569–2574.
- (75) Egberts, E.; Marrink, S.-J.; Berendsen, H. Molecular dynamics simulation of a phospholipid membrane. *Eur. Biophys. J.* **1994**, *22*, 423–436.
- (76) Klauda, J. B.; Venable, R. M.; Freites, J. A.; O’Connor, J. W.; Tobias, D. J.; Mondragon-Ramirez, C.; Vorobyov, I.; MacKerell Jr, A. D.; Pastor, R. W. Update of the CHARMM All-Atom Additive Force Field for Lipids: Validation on Six Lipid Types. *J. Phys. Chem. B* **2010**, *114*, 7830–7843.

- (77) Wang, Y.; Zhao, T.; Wei, D.; Strandberg, E.; Ulrich, A. S.; Ulmschneider, J. P. How reliable are molecular dynamics simulations of membrane active antimicrobial peptides? *Biochim. Biophys. Acta* **2014**, *1838*, 2280–2288.
- (78) MacCallum, J. L.; Bennett, W. F. D.; Tieleman, D. P. Distribution of Amino Acids in a Lipid Bilayer from Computer Simulations. *Biophys. J.* **2008**, *94*, 3393–3404.
- (79) Johansson, A. C.; Lindahl, E. Position-resolved free energy of solvation for amino acids in lipid membranes from molecular dynamics simulations. *Proteins: Struct. Funct. Bioinf.* **2008**, *70*, 1332–1344.
- (80) Reißer, S.; Strandberg, E.; Steinbrecher, T.; Elstner, M.; Ulrich, A. S. Best of Two Worlds? How MD Simulations of Amphiphilic Helical Peptides in Membranes Can Complement Data from Oriented Solid-State NMR. *J. Chem. Theory Comput* **2018**, *14*, 6002–6014.
- (81) Duboué-Dijon, E.; Javanainen, M.; Delcroix, P.; Jungwirth, P.; Martinez-Seara, H. A practical guide to biologically relevant molecular simulations with charge scaling for electronic polarization. *J. Chem. Phys.* **2020**, *153*, 050901.
- (82) Paloncýová, M.; Fabre, G.; DeVane, R. H.; Trouillas, P.; Berka, K.; Otyepka, M. Benchmarking of Force Fields for Molecule–Membrane Interactions. *J. Chem. Theory Comput* **2014**, *10*, 4143–4151.
- (83) Bittermann, K.; Spycher, S.; Endo, S.; Pohler, L.; Huniar, U.; Goss, K.-U.; Klamt, A. Prediction of Phospholipid–Water Partition Coefficients of Ionic Organic Chemicals Using the Mechanistic Model COSMOmic. *J. Phys. Chem. B* **2014**, *118*, 14833–14842.
- (84) Kadaoluwa Pathirannahalage, S. P.; Meftahi, N.; Elbourne, A.; Weiss, A. C. G.; McConville, C. F.; Padua, A.; Winkler, D. A.; Costa Gomes, M.; Greaves, T. L.; Le, T. C. et al. Systematic Comparison of the Structural and Dynamic Properties of Commonly

- Used Water Models for Molecular Dynamics Simulations. *J. Chem. Inf. Model.* **2021**, *61*, 4521–4536.
- (85) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **1983**, *79*, 926–935.
- (86) Ollila, O. H. S.; Heikkinen, H. A.; Iwai, H. Rotational Dynamics of Proteins from Spin Relaxation Times and Molecular Dynamics Simulations. *J. Phys. Chem. B* **2018**, *122*, 6559–6569.
- (87) Best, R. B.; Zheng, W.; Mittal, J. Balanced Protein–Water Interactions Improve Properties of Disordered Proteins and Non-Specific Protein Association. *J. Chem. Theory Comput* **2014**, *10*, 5113–5124.
- (88) Piana, S.; Donchev, A. G.; Robustelli, P.; Shaw, D. E. Water Dispersion Interactions Strongly Influence Simulated Structural Properties of Disordered Protein States. *J. Phys. Chem. B* **2015**, *119*, 5113–5123.
- (89) Li, H.; Chowdhary, J.; Huang, L.; He, X.; MacKerell Jr, A. D.; Roux, B. Drude polarizable force field for molecular dynamics simulations of saturated and unsaturated zwitterionic lipids. *J. Chem. Theory Comput* **2017**, *13*, 4535–4552.
- (90) Chowdhary, J.; Harder, E.; Lopes, P. E.; Huang, L.; MacKerell Jr, A. D.; Roux, B. A polarizable force field of dipalmitoylphosphatidylcholine based on the classical drude model for molecular dynamics simulations of lipids. *J. Phys. Chem. B* **2013**, *117*, 9142–9160.
- (91) Chu, H.; Peng, X.; Li, Y.; Zhang, Y.; Li, G. A polarizable atomic multipole-based force field for molecular dynamics simulations of anionic lipids. *Molecules* **2018**, *23*, 77.

- (92) Thole, B. Molecular polarizabilities calculated with a modified dipole interaction. *Chem. Phys.* **1981**, *59*, 341–350.
- (93) Leontyev, I. V.; Stuchebrukhov, A. A. Electronic continuum model for molecular dynamics simulations of biological molecules. *J. Chem. Theory Comput* **2010**, *6*, 1498–1508.
- (94) Gadsby, D. C. Ion channels versus ion pumps: the principal difference, in principle. *Nat. Rev. Mol. Cell Biol.* **2009**, *10*, 344–352.
- (95) Weinbaum, S.; Tarbell, J. M.; Damiano, E. R. The Structure and Function of the Endothelial Glycocalyx Layer. *Annu. Rev. Biomed. Eng* **2007**, *9*, 121–167.
- (96) Shi, X.; Bi, Y.; Yang, W.; Guo, X.; Jiang, Y.; Wan, C.; Li, L.; Bai, Y.; Guo, J.; Wang, Y. et al. Ca<sup>2+</sup> regulates T-cell receptor activation by modulating the charge property of lipids. *Nature* **2013**, *493*, 111–115.
- (97) Unke, O. T.; Chmiela, S.; Sauceda, H. E.; Gastegger, M.; Poltavsky, I.; Schütt, K. T.; Tkatchenko, A.; Müller, K.-R. Machine Learning Force Fields. *Chem. Rev.* **2021**, *121*, 10142–10186.
- (98) Wilkinson, M. D.; Dumontier, M.; Aalbersberg, I. J.; Appleton, G.; Axton, M.; Baak, A.; Blomberg, N.; Boiten, J.-W.; da Silva Santos, L. B.; Bourne, P. E. et al. The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* **2016**, *3*, 160018.
- (99) Zhuang, X.; Makover, J. R.; Im, W.; Klauda, J. B. A systematic molecular dynamics simulation study of temperature dependent bilayer structural properties. *Biochim. Biophys. Acta* **2014**, *1838*, 2520 – 2529.
- (100) Zhuang, X.; Dávila-Contreras, E. M.; Beaven, A. H.; Im, W.; Klauda, J. B. An exten-

- sive simulation study of lipid bilayer properties with different head groups, acyl chain lengths, and chain saturations. *Biochim. Biophys. Acta* **2016**, *1858*, 3093–3104.
- (101) Bera, I.; Klauda, J. B. Molecular Simulations of Mixed Lipid Bilayers with Sphingomyelin, Glycerophospholipids, and Cholesterol. *J. Phys. Chem. B* **2017**, *121*, 5197–5208.
- (102) Gowers, T.; Nielsen, M. Massively collaborative mathematics. *Nature* **2009**, *461*, 879–881.
- (103) Antila, H. S.; M. Ferreira, T.; Ollila, O. H. S.; Miettinen, M. S. Using Open Data to Rapidly Benchmark Biomolecular Simulations: Phospholipid Conformational Dynamics. *J. Chem. Inf. Model.* **2021**, *61*, 938–949.
- (104) Ferreira, T. M.; Ollila, O. H. S.; Pigliapochi, R.; Dabkowska, A. P.; Topgaard, D. Model-free estimation of the effective correlation time for C-H bond reorientation in amphiphilic bilayers:  $^1\text{H}$ - $^{13}\text{C}$  solid-state NMR and MD simulations. *J. Chem. Phys.* **2015**, *142*, 044905.
- (105) Kučerka, N.; Nieh, M. P.; Katsaras, J. Fluid phase lipid areas and bilayer thicknesses of commonly used phosphatidylcholines as a function of temperature. *Biochim. Biophys. Acta* **2011**, *1808*, 2761–2771.
- (106) Pan, J.; Cheng, X.; Monticelli, L.; Heberle, F. A.; Kucerka, N.; Tieleman, D. P.; Katsaras, J. The molecular structure of a phosphatidylserine bilayer determined by scattering and molecular dynamics simulations. *Soft Matter* **2014**, *10*, 3716–3725.
- (107) Leftin, A.; Molugu, T.; Job, C.; Beyer, K.; Brown, M. Area per Lipid and Cholesterol Interactions in Membranes from Separated Local-Field  $^{13}\text{C}$  NMR Spectroscopy. *Biophys. J.* **2014**, *107*, 2274 – 2286.

- (108) Beschiaschvili, G.; Seelig, J. Peptide binding to lipid membranes. Spectroscopic studies on the insertion of a cyclic somatostatin analog into phospholipid bilayers. *Biochim. Biophys. Acta*, volume=1061, number=1, pages=78–84, year=1991, publisher=Elsevier
- (109) Palivec, V.; Kopecký, V.; Jungwirth, P.; Bouř, P.; Kaminský, J.; Martinez-Seara, H. Simulation of Raman and Raman optical activity of saccharides in solution. *Phys. Chem. Chem. Phys.* **2020**, *22*, 1983–1993.
- (110) Palivec, V.; Johannessen, C.; Kaminský, J.; Martinez-Seara, H. Use of Raman and Raman optical activity to extract atomistic details of saccharides in aqueous solution. *PLOS Comput. Biol* **2022**, *18*, e1009678.
- (111) Handley, C. M.; Popelier, P. L. A. Dynamically Polarizable Water Potential Based on Multipole Moments Trained by Machine Learning. *J. Chem. Theory Comput* **2009**, *5*, 1474–1489, PMID: 26609841.
- (112) Dutta, R.; Brotzakis, Z. F.; Mira, A. Bayesian calibration of force-fields from experimental data: TIP4P water. *J. Chem. Phys.* **2018**, *149*, 154110.
- (113) Noé, F.; Tkatchenko, A.; Müller, K.-R.; Clementi, C. Machine Learning for Molecular Simulation. *Annu. Rev. Phys. Chem.* **2020**, *71*, 361–390.
- (114) Behler, J.; Parrinello, M. Generalized Neural-Network Representation of High-Dimensional Potential-Energy Surfaces. *Phys. Rev. Lett.* **2007**, *98*, 146401.
- (115) Gao, X.; Ramezanghorbani, F.; Isayev, O.; Smith, J. S.; Roitberg, A. E. TorchANI: A Free and Open Source PyTorch-Based Deep Learning Implementation of the ANI Neural Network Potentials. *J. Chem. Inf. Model.* **2020**, *60*, 3408–3415, PMID: 32568524.
- (116) Smith, J. S.; Nebgen, B.; Lubbers, N.; Isayev, O.; Roitberg, A. E. Less is more: Sampling chemical space with active learning. *J. Chem. Phys.* **2018**, *148*, 241733.

- (117) Schütt, K. T.; Kessel, P.; Gastegger, M.; Nicoli, K. A.; Tkatchenko, A.; Müller, K.-R. SchNetPack: A Deep Learning Toolbox For Atomistic Systems. *J. Chem. Theory Comput* **2019**, *15*, 448–455.
- (118) Saucedo, H. E.; Gastegger, M.; Chmiela, S.; Müller, K.-R.; Tkatchenko, A. Molecular force fields with gradient-domain machine learning (GDML): Comparison and synergies with classical force fields. *J. Chem. Phys.* **2020**, *153*, 124109.
- (119) Wang, Y.; Fass, J.; Kaminow, B.; Herr, J. E.; Rufa, D.; Zhang, I.; Pulido, I.; Henry, M.; Chodera, J. D. End-to-End Differentiable Molecular Mechanics Force Field Construction. 2020; <https://arxiv.org/abs/2010.01196>.
- (120) Zahariev, F.; De Silva, N.; Gordon, M. S.; Windus, T. L.; Pérez García, M. ParFit: A Python-Based Object-Oriented Program for Fitting Molecular Mechanics Parameters to ab Initio Data. *J. Chem. Inf. Model.* **2017**, *57*, 391–396, PMID: 28169538.
- (121) Gastegger, M.; Behler, J.; Marquetand, P. Machine learning molecular dynamics for the simulation of infrared spectra. *Chem. Sci.* **2017**, *8*, 6924–6935.
- (122) Rosenberger, D.; Smith, J. S.; Garcia, A. E. Modeling of Peptides with Classical and Novel Machine Learning Force Fields: A Comparison. *J. Phys. Chem. B* **2021**, *125*, 3598–3612, PMID: 33798336.
- (123) Smith, J. S.; Roitberg, A. E.; Isayev, O. Transforming Computational Drug Discovery with Machine Learning and AI. *ACS Med. Chem. Lett* **2018**, *9*, 1065–1069.
- (124) Hunger, J. In *Philosophy Of Chemistry: Synthesis of a New Discipline*; Baird, D., Scerri, E., McIntyre, L., Eds.; Springer Netherlands: Dordrecht, 2006; pp 129–156.
- (125) Hunger, J.; Beyreuther, S.; Huttner, G.; Allinger, K.; Radelof, U.; Zsolnai, L. How to Derive Force Field Parameters by Genetic Algorithms: Modelling tripod-Mo(CO)<sub>3</sub> Compounds as an Example. *Eur. J. Inorg. Chem.* **1998**, *1998*, 693–702.

- (126) McKiernan, K. A.; Wang, L.-P.; Pande, V. S. Training and validation of a liquid-crystalline phospholipid bilayer force field. *J. Chem. Theory Comput* **2016**, *12*, 5960–5967.
- (127) Yu, Y.; Krämer, A.; Venable, R. M.; Brooks, B. R.; Klauda, J. B.; Pastor, R. W. CHARMM36 Lipid Force Field with Explicit Treatment of Long-Range Dispersion: Parametrization and Validation for Phosphatidylethanolamine, Phosphatidylglycerol, and Ether Lipids. *J. Chem. Theory Comput* **2021**, *17*, 1581–1595, PMID: 33620194.
- (128) Huang, J.; MacKerell, A. D., Jr CHARMM36 all-atom additive protein force field: validation based on comparison to NMR data. *J. Comput. Chem.* **2013**, *34*, 2135–2145.
- (129) Cesari, A.; Bottaro, S.; Lindorff-Larsen, K.; Banaš, P.; Šponer, J.; Bussi, G. Fitting corrections to an RNA force field using experimental data. *J. Chem. Theory Comput* **2019**, *15*, 3425–3431.
- (130) Reith, D.; Meyer, H.; Müller-Plathe, F. CG-OPT: A software package for automatic force field design. *Comput. Phys. Commun* **2002**, *148*, 299–313.
- (131) Fogarty, J. C.; Chiu, S.-W.; Kirby, P.; Jakobsson, E.; Pandit, S. A. Automated Optimization of Water–Water Interaction Parameters for a Coarse-Grained Model. *J. Phys. Chem. B* **2014**, *118*, 1603–1611, PMID: 24460506.
- (132) Betz, R. M.; Walker, R. C. Paramfit: Automated optimization of force field parameters for molecular dynamics simulations. *J. Comput. Chem.* **2015**, *36*, 79–87.
- (133) Sestito, J. M.; Thatcher, M. L.; Shu, L.; Harris, T. A.; Wang, Y. Coarse-Grained Force Field Calibration Based on Multiobjective Bayesian Optimization to Simulate Water Diffusion in Poly- $\epsilon$ -caprolactone. *J. Phys. Chem. A* **2020**, *124*, 5042–5052.



- (134) Befort, B. J.; DeFever, R. S.; Tow, G. M.; Dowling, A. W.; Maginn, E. J. Machine Learning Directed Optimization of Classical Molecular Modeling Force Fields. *J. Chem. Inf. Model.* **2021**, *61*, 4400–4414.
- (135) Messerly, R. A.; Razavi, S. M.; Shirts, M. R. Configuration-sampling-based surrogate models for rapid parameterization of non-bonded interactions. *J. Chem. Theory Comput* **2018**, *14*, 3144–3162.
- (136) Boothroyd, S.; Wang, L.-P.; Mobley, D. L.; Chodera, J. D.; Shirts, M. R. Open Force Field Evaluator: An Automated, Efficient, and Scalable Framework for the Estimation of Physical Properties from Molecular Simulation. *J. Chem. Theory Comput.* **2022**, <https://doi.org/10.1021/acs.jctc.1c01111>.
- (137) Shaw, D. E.; Adams, P. J.; Azaria, A.; Bank, J. A.; Batson, B.; Bell, A.; Bergdorf, M.; Bhatt, J.; Butts, J. A.; Correia, T. et al. Anton 3: Twenty Microseconds of Molecular Dynamics Simulation before Lunch. Proceedings of the International Conference for High Performance Computing, Networking, Storage and Analysis. New York, NY, USA, 2021; pp 1–11.
- (138) Norrby, P.-O.; Liljefors, T. Automated molecular mechanics parameterization with simultaneous utilization of experimental and quantum mechanical data. *J. Comput. Chem.* **1998**, *19*, 1146–1166.
- (139) Mostaghim, S.; Hoffmann, M.; Konig, P. H.; Frauenheim, T.; Teich, J. Molecular force field parametrization using multi-objective evolutionary algorithms. Proceedings of the 2004 Congress on Evolutionary Computation (IEEE Cat. No. 04TH8753). 2004; pp 212–219.
- (140) Tafipolsky, M.; Schmid, R. Systematic first principles parameterization of force fields for metal-organic frameworks using a genetic algorithm approach. *J. Phys. Chem. B* **2009**, *113*, 1341–1352.

- (141) Liu, Y.-Y.; Slotine, J.-J.; Barabási, A.-L. Controllability of complex networks. *Nature* **2011**, *473*, 167–173.
- (142) Handley, C. M.; Deeth, R. J. A multi-objective approach to force field optimization: structures and spin state energetics of d6 Fe (II) complexes. *J. Chem. Theory Comput* **2012**, *8*, 194–202.
- (143) Li, Y.; Hartke, B. Approximate photochemical dynamics of azobenzene with reactive force fields. *J. Chem. Phys.* **2013**, *139*, 224303.
- (144) Dittner, M.; Müller, J.; Aktulga, H. M.; Hartke, B. Efficient global optimization of reactive force-field parameters. *J. Comput. Chem.* **2015**, *36*, 1550–1561.
- (145) Gao, D. Z.; Federici Canova, F.; Watkins, M. B.; Shluger, A. L. Efficient parametrization of complex molecule–surface force fields. *J. Comput. Chem.* **2015**, *36*, 1187–1195.
- (146) Betz, R. M.; Walker, R. C. Paramfit: Automated optimization of force field parameters for molecular dynamics simulations. *J. Comput. Chem.* **2015**, *36*, 79–87.
- (147) Wang, L.-P.; McKiernan, K. A.; Gomes, J.; Beauchamp, K. A.; Head-Gordon, T.; Rice, J. E.; Swope, W. C.; Martínez, T. J.; Pande, V. S. Building a more predictive protein force field: a systematic and reproducible route to AMBER-FB15. *J. Phys. Chem. B* **2017**, *121*, 4023–4039.
- (148) Oliveira, M. P.; Andrey, M.; Rieder, S. R.; Kern, L.; Hahn, D. F.; Riniker, S.; Horta, B. A.; Hünenberger, P. H. Systematic optimization of a fragment-based force field against experimental pure-liquid properties considering large compound families: application to saturated haloalkanes. *J. Chem. Theory Comput* **2020**, *16*, 7525–7555.
- (149) Krishnamoorthy, A.; Mishra, A.; Grabar, N.; Baradwaj, N.; Kalia, R. K.; Nakano, A.; Vashishta, P. Evolutionary multi-objective optimization and Pareto-frontal uncer-

- tainty quantification of interatomic forcefields for thermal conductivity simulations. *Comput. Phys. Commun* **2020**, *254*, 107337.
- (150) Oliveira, M. P.; Hünenberger, P. H. Systematic optimization of a fragment-based force field against experimental pure-liquid properties considering large compound families: application to oxygen and nitrogen compounds. *Phys. Chem. Chem. Phys.* **2021**, *23*, 17774–17793.
- (151) Yang, L.; Tiwari, S. C.; Jo, S. S.; Hong, S.; Mishra, A.; Krishnamoorthy, A.; Kalia, R. K.; Nakano, A.; Jaramillo, R.; Vashishta, P. Unveiling oxidation mechanism of bulk ZrS<sub>2</sub>. *MRS Advances* **2021**, *6*, 303–306.
- (152) Morawietz, T.; Behler, J. A density-functional theory-based neural network potential for water clusters including van der Waals corrections. *J. Phys. Chem. A* **2013**, *117*, 7356–7366.
- (153) Smith, J. S.; Isayev, O.; Roitberg, A. E. ANI-1: an extensible neural network potential with DFT accuracy at force field computational cost. *Chem. Sci* **2017**, *8*, 3192–3203.
- (154) Thaler, S.; Zavadlav, J. Learning neural network potentials from experimental data via Differentiable Trajectory Reweighting. *Nat. Commun.* **2021**, *12*, 1–10.




Hanne S. Antila obtained her doctorate in computational chemistry from Aalto University, Finland, in 2016. She did postdoctoral work at Northwestern University (USA), after which she became a postdoctoral researcher at the Max Planck Institute for Colloids and Interfaces (Germany), where she currently works as an Alexander von Humboldt fellow. Her research interests are in understanding interfacial effects and how those connect to transport, as well as in computational method and model development for soft matter.



Batuhan Kav obtained his BSc. and MSc. degrees in chemistry and physics from Bilkent University and Koç University, Turkey, respectively with a short stay at the University of Cambridge. In 2019 he completed his PhD in theoretical physics at the Max Planck Institute of Colloids and Interfaces in Potsdam, Germany where he investigated the role of lipid-anchored saccharides in membrane adhesion using molecular dynamics simulations under the supervision of Thomas Weikl and Emanuel Schneck. Since 2019, he is working in the group of Birgit Strodel at the Institute of Biological Information Processing (IBI-7), Forschungszentrum Jülich, Germany with the focus on developing novel molecular dynamics force fields to study the amyloid aggregation. Since 2018, he is a

member of the NMRlipids Project. His research focuses on benchmarking and developing both polarisable and non-polarisable all-atom molecular dynamics force fields for intrinsically disordered proteins and lipid bilayers.



Markus S. Miettinen received his Ph.D. in Theoretical and Computational Physics from Aalto University (Helsinki, Finland) in 2010. After postdoctoral work in Germany, as an EMBO Long-Term Fellow at the University of Potsdam and a Volkswagen Foundation Fellow at the Freie Universität Berlin, he moved to the Max Planck Institute of Colloids and Interfaces, where he was a Research Group Leader at the Department of Theory and Bio-Systems 2017–2022. Currently he is an Associate Professor of Computational Chemistry at the University of Bergen (Norway), where he also leads the High-Fidelity Biomolecular Modelling Group of the Computational Biology Unit. In his research he develops methods for automated data-driven calibration of MD models of disordered biomolecules, and for calculation of local materials properties from such calibrated simulations. He is a co-founder of the NMRlipids Project.



Hector Martinez-Seara received Ph.D. in chemistry from Barcelona University in 2010, focusing on studying the structure of biological membranes

using computational methods. During his postdoctoral work at Tampere University of Technology (Finland) under the supervision of Prof. Ilpo Vattulainen, he enlarged his simulation skills to include dynamic properties and coarse-grained methods applied to biomembranes. He is currently established at the Czech Academy of Sciences (Czech Republic) as a team leader and head of the high performance computing service group. Now, he focuses on the computational study of the glycocalyx that adds glycoscience to his expertise. He enjoys combining spectroscopic experimental results and simulations to access structural details that otherwise remain hidden. He also aims to improve simulation protocols used in biosystems (e.g., ECC) to promote simulations as a reference tool in his spare time.



Pavel Jungwirth (PhD. 1993) is a Distinguished Chair at the Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences and a full professor in physics (External Faculty) at the Charles University in Prague. His scientific interests involve modeling of ion–protein interactions, solvated electrons, and molecular processes in cellular membranes. He never thought he will end up developing force fields with the aim to try to be useful to the scientific community, but never say never.



O. H. Samuli Ollila received Ph.D. in biophysics from Tampere University of Technology, Finland, in 2010. After post doctoral work in Lund University (Sweden), Aalto University (Finland) and Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences (Czech Republic), he is currently an Academy of Finland Research Fellow in the Institute of Biotechnology, University of Helsinki, Finland. His research is focused to develop methods to study disordered biomolecules, such as lipids and disordered proteins, by combining MD simulations and NMR experiments. He is a co-founder and the leader of the NMRlipids open collaboration project.