



# Generalized isotopic shift construction for APN functions

Lilya Budaghyan<sup>1</sup> · Marco Calderini<sup>1</sup> · Claude Carlet<sup>1,2</sup> · Robert Coulter<sup>3</sup> · Irene Villa<sup>1</sup>

Received: 18 March 2020 / Revised: 24 June 2020 / Accepted: 8 September 2020  
© The Author(s) 2020

## Abstract

In this work we give several generalizations of the isotopic shift construction, introduced recently by Budaghyan et al. (IEEE Trans Inform Theory 66:5299–5309, 2020), when the initial function is a Gold function. In particular, we derive a general construction of APN functions which covers several unclassified APN functions for  $n = 8$  and produces fifteen new APN functions for  $n = 9$ .

**Keywords** APN functions · Isotopic shift · Vectorial Boolean functions

**Mathematics Subject Classification** 94A60 · 11T71 · 06E30

## 1 Introduction

For  $n$  a positive integer, let  $\mathbb{F}_{2^n}$  be the finite field with  $2^n$  elements. By  $\mathbb{F}_{2^n}^*$  we denote the multiplicative group of  $\mathbb{F}_{2^n}$  and, throughout the paper,  $\zeta$  denotes one of its primitive elements,

---

Communicated by A. Pott.

---

Parts of this work were presented at *WCC 2019: The Eleventh International Workshop on Coding and Cryptography*.

---

✉ Marco Calderini  
marco.calderini@uib.no

Lilya Budaghyan  
lilya.budaghyan@uib.no

Claude Carlet  
claude.carlet@gmail.com

Robert Coulter  
coulter@udel.edu

Irene Villa  
irene.villa@uib.no

<sup>1</sup> Department of Informatics, University of Bergen, PB 7803, 5020 Bergen, Norway

<sup>2</sup> LAGA, University of Paris 8, Saint-Denis, France

<sup>3</sup> Department of Mathematical Sciences, University of Delaware, Newark, DE, USA

so that  $\mathbb{F}_{2^n}^* = \langle \zeta \rangle = \{1, \zeta, \zeta^2, \zeta^3, \dots, \zeta^{2^n-2}\}$ . An  $(n, n)$ -function is a map from  $\mathbb{F}_{2^n}$  to itself. Such function admits a unique representation as a univariate polynomial of degree at most  $2^n - 1$ , that is

$$F(x) = \sum_{j=0}^{2^n-1} a_j x^j, \quad a_j \in \mathbb{F}_{2^n}.$$

The kernel of  $F$  is defined as  $\ker(F) = \{u \in \mathbb{F}_{2^n} \text{ s.t. } F(u) = 0\}$ .

The function  $F$  is

- *linear* if  $F(x) = \sum_{i=0}^{n-1} c_i x^{2^i}$ ;
- *affine* if it is the sum of a linear function and a constant;
- *DO* (Dembowski-Ostrom) *polynomial* if  $F(x) = \sum_{0 \leq i < j < n} a_{ij} x^{2^i+2^j}$ , with  $a_{ij} \in \mathbb{F}_{2^n}$ ;
- *quadratic* if it is the sum of a DO polynomial and an affine function.

A function  $F$  is called *differentially  $\delta$ -uniform*, for  $\delta$  a positive integer, if for any pair  $(a, b) \in \mathbb{F}_{2^n}^2$ , with  $a \neq 0$ , the equation  $F(x+a) - F(x) = b$  admits at most  $\delta$  solutions. When  $F$  is used as an S-box inside a cryptosystem, the differential uniformity measures its contribution to the resistance to the differential attack [3]. The smaller is  $\delta$ , the better is the resistance to this attack.

Over fields of characteristic 2, the solutions of the equation  $F(x+a) - F(x) = b$ , that is,  $F(x+a) + F(x) = b$ , go by pairs  $\{x, x+a\}$ , and  $\delta$  is even. The best resistance is then achieved by differentially 2-uniform functions. Such functions are also called *almost perfect nonlinear*; in short, APN. The simplest known example of APN function is Gold function,  $\mathcal{G}_i(x) = x^{2^i+1}$ , that is APN whenever  $i$  is coprime with  $n$ .

APN functions have connections to optimal objects in other fields such as geometry, sequence design and combinatorics.

There are several equivalence relations of functions for which differential uniformity, and thus the APN property, is preserved. Two functions  $F$  and  $F'$  from  $\mathbb{F}_{2^n}$  to itself are called:

- affine equivalent if  $F' = A_1 \circ F \circ A_2$  where  $A_1, A_2 : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$  are affine permutations;
- EA-equivalent if  $F' = F'' + A$ , where the map  $A : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$  is affine and  $F''$  is affine equivalent to  $F$ ;
- CCZ-equivalent [12] if there exists some affine permutation  $\mathcal{L}$  of  $\mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$  such that the image of the graph of  $F$  is the graph of  $F'$ , that is,  $\mathcal{L}(G_F) = G_{F'}$ , where  $G_F = \{(x, F(x)) : x \in \mathbb{F}_{2^n}\}$  and  $G_{F'} = \{(x, F'(x)) : x \in \mathbb{F}_{2^n}\}$ .

CCZ-equivalence is the most general known equivalence relation for functions which preserves differential uniformity, while affine and EA-equivalences are its particular cases. We refer the reader to [5] and [11] for a more comprehensive overview on vectorial Boolean functions.

Inspired by the notion of *isotopic equivalence*, originally defined by Albert [1] in the study of presemifields and semifields, a new construction method for APN functions, called isotopic shift, was introduced in [6].

Given  $p$  a prime number,  $F \in \mathbb{F}_{p^n}[x]$  a function, and  $L \in \mathbb{F}_{p^n}[x]$  a linear map, the *isotopic shift* of  $F$  by  $L$  is defined as the map:

$$F_L(x) = F(x + L(x)) - F(x) - F(L(x)). \quad (1)$$

As we have shown in [6], for the case  $p = 2$ , an isotopic shift of an APN function can lead to APN functions CCZ-inequivalent to the original map. In particular, all existing quadratic APN functions over  $\mathbb{F}_{2^6}$ , which are 13 up to CCZ-equivalence, can be obtained from  $x^3$  by

isotopic shift. Moreover, a new family of quadratic APN functions, which generates a new APN function for  $n = 9$ , is constructed by isotopic shift of Gold functions [6]. In [7], the isotopic shift construction has been investigated for the case of planar functions ( $p > 2$ ), i.e. differentially 1-uniform functions. Also here, given a planar function, it is possible to obtain an inequivalent planar function from its isotopic shifts.

In the present paper we further study the isotopic shift construction over fields of even characteristic. Firstly, we verify that, over  $\mathbb{F}_{2^6}$ , any quadratic APN map can be obtained as an isotopic shift of any other quadratic APN map. Then, we consider different generalizations of the isotopic shift construction when the initial function is a monomial with a Gold exponent. In [6], we studied the APN property of the isotopic shift of  $\mathcal{G}_i(x) = x^{2^i+1}$  over  $\mathbb{F}_{2^n}$ , with  $n = km$ , given by

$$\mathcal{G}_{i,L}(x) = xL(x)^{2^i} + x^{2^i}L(x), \tag{2}$$

where  $L$  is a  $2^m$ -polynomial, that is  $L(x) = \sum_{i=0}^{k-1} A_i x^{2^{im}}$  for some  $A_i \in \mathbb{F}_{2^n}$ . This construction provides a new APN function over  $\mathbb{F}_{2^9}$ .

In the present work, we study the APN property of  $xL_1(x)^{2^i} + x^{2^i}L_2(x)$  where both  $L_1$  and  $L_2$  are  $2^m$ -polynomials. From this construction, we obtain fifteen new APN functions for  $n = 9$ . Moreover, we cover some of the functions in the lists given in [16] and [19] which are not contained in any of the known infinite families.

To show the inequivalence between some of the obtained maps, we introduce in Proposition 3.2 a new EA-invariant (such invariant was also noticed independently in [17]). Note that for quadratic APN functions, CCZ-equivalence coincides with EA-equivalence [23].

Finally, we consider the case when the isotopic shift of  $\mathcal{G}_i(x)$  is obtained using a function  $L$  not necessarily linear. In this case we obtain that all the known power APN functions in odd dimension, except the Dobbertin function, can be obtained as the nonlinear shifts of Gold functions.

## 2 Further results on the isotopic linear shift over $\mathbb{F}_{2^n}$

Before considering generalizations of the isotopic shift, we extend a result obtained in [6].

We have shown that, given a quadratic APN function  $F$ , if the isotopic shift  $F_L$  by a linear map  $L$  is APN, then the map  $L$  is either a permutation or a 2-to-1 map. From the isotopic shifts of the Gold function  $x^3$ , with both choices for  $L$  being a permutation and a 2-to-1 map, we obtained (computationally) all the quadratic APN functions over  $\mathbb{F}_{2^6}$  (up to EA-equivalence). That is, for any given quadratic APN function  $F$  over  $\mathbb{F}_{2^6}$  there exist a permutation  $L$  and a 2-to-1 map  $L'$  such that the isotopic shifts  $\mathcal{G}_{1,L}(x)$  and  $\mathcal{G}_{1,L'}(x)$  are EA-equivalent to  $F$ . The same result was computationally obtained for any quadratic APN map over  $\mathbb{F}_{2^6}$  listed in [16, Table 5] (see also [4]) in place of  $\mathcal{G}_1$ . Up to EA-equivalence (and thus CCZ-equivalence) the list is complete and, since for two quadratic maps the EA-equivalence implies EA-equivalence of the isotopic shifts (see [6, Corollary 3.2]), we can state the following result.

**Proposition 2.1** *Over  $\mathbb{F}_{2^6}$  for any two quadratic APN maps  $F$  and  $G$ , there exist a linear permutation  $L$  and a linear 2-to-1 map  $L'$  such that  $F_L$  and  $F_{L'}$  are EA-equivalent to  $G$ .*

We conclude with the observation that the isotopic shift can lead to an APN function also starting from a non-APN function.

**Remark 2.1** Consider  $\mathbb{F}_{2^6}$  and the function  $F(x) = x^5$ , which is not APN. With  $L(x) = \zeta x^8$  we construct the APN map

$$F_L(x) = x^4 L(x) + x L(x)^4 = \zeta x^{12} + \zeta^4 x^{33},$$

where  $F_L(x) = M(x^3)$  for the linear permutation  $M(x) = \zeta x^4 + \zeta^4 x^{32}$ .

### 3 Generalized isotopic shift of Gold functions

In this section we generalize the isotopic shift construction for the case of Gold functions.

#### 3.1 On the generalized linear shift over $\mathbb{F}_{2^n}$

In [6], we showed that the isotopic shift can be a useful construction method for APN functions. Let  $n = km$ , where  $m$  and  $k$  are any positive integers. An  $\mathbb{F}_{2^m}$ -polynomial is a linear map given by  $L(x) = \sum_{j=0}^{k-1} A_j x^{2^{jm}}$ , for some  $A_j \in \mathbb{F}_{2^m}$ . The construction  $\mathcal{G}_{i,L}(x)$  as in (2) leads to a family of APN functions, providing, in particular, for  $n = 9$  ( $k, m = 3$ ) a new APN function and for  $n = 8$  ( $k = 4, m = 2$ ) a function equivalent to  $x^9 + \text{Tr}(x^3)$ , which is not contained in any infinite family.

In the following, we generalize the isotopic shift construction. This generalization provides further new APN functions, as it will be shown below.

Given two positive integers  $k, m$ , let us consider the finite field  $\mathbb{F}_{2^n}$  with  $n = km$ . Denoting  $d = \gcd(2^m - 1, \frac{2^{km}-1}{2^m-1})$ , let  $d'$  be the positive integer with the same prime factors as  $d$ , satisfying  $\gcd(2^m - 1, \frac{2^{km}-1}{(2^m-1)d'}) = 1$ . Now, let  $U = \langle \zeta^{d'(2^m-1)} \rangle$  be the multiplicative subgroup of  $\mathbb{F}_{2^m}^*$  of order  $(\frac{2^{km}-1}{2^m-1})/d'$ . Note that it is possible to write every element  $x \in \mathbb{F}_{2^m}^*$  as  $x = ut$  with  $u \in W$  and  $t \in \mathbb{F}_{2^m}^*$ , where  $W = \{\zeta^s y : y \in U, 0 \leq s \leq d' - 1\}$ . Indeed, let  $\mathbb{F}_{2^{mk}}^* = \langle \zeta \rangle$ , we have  $x = \zeta^{d'z+j}$ , for some integers  $z$  and  $j$  where  $0 \leq j \leq d' - 1$ . For ease of notation, set  $l = \frac{2^{mk}-1}{(2^m-1)d'}$ . Since  $\gcd(2^m - 1, l) = 1$ , for any such  $z$ , there exist integers  $r$  and  $s$  such that  $z = r(2^m - 1) + sl$ . Hence we have

$$x = \zeta^{d'z+j} = \zeta^{d'r(2^m-1)} \zeta^j \zeta^{d'sl} = ut,$$

where, denoting  $y = \zeta^{d'r(2^m-1)} \in U$ , we have  $u = y\zeta^j \in W$  and  $t = \zeta^{d'sl} = \zeta^{s(\frac{2^{mk}-1}{2^m-1})} \in \mathbb{F}_{2^m}^*$ . Since  $|\{(u, t) : u \in W, t \in \mathbb{F}_{2^m}^*\}| = |W| \cdot |\mathbb{F}_{2^m}^*| = (d'|U|) \cdot (2^m - 1) = d' \cdot \frac{2^{mk}-1}{d'(2^m-1)} \cdot (2^m - 1) = 2^{mk} - 1 = |\mathbb{F}_{2^{mk}}^*|$ , two distinct elements in  $\mathbb{F}_{2^{mk}}^*$  cannot have the same representation, so  $u$  and  $t$  are unique.

Then it is possible to obtain the following generalization of [6, Theorem 6.3].

**Theorem 3.1** *Let  $n = km$  for  $m > 1$ . Let  $L_1(x) = \sum_{j=0}^{k-1} A_j x^{2^{jm}}$  and  $L_2(x) = \sum_{j=0}^{k-1} B_j x^{2^{jm}}$  be two  $\mathbb{F}_{2^m}$ -polynomials. Then, let  $i$  be such that  $\gcd(i, m) = 1$  and  $F \in \mathbb{F}_{2^n}[x]$  the function given by:*

$$F(x) = x L_1(x)^{2^i} + x^{2^i} L_2(x). \tag{3}$$

*Then  $F$  is APN over  $\mathbb{F}_{2^n}$  if and only if each of the following statements holds for any  $v \in W$ :*

- $(\frac{L_1(v)}{v})^{2^i} \neq \frac{L_2(v)}{v}$ ;
- If  $u \in W \setminus \{1\}$  and  $(\frac{L_1(uv)}{uv})^{2^i} = \frac{L_2(v)}{v}$ , then  $(\frac{L_1(v)}{v})^{2^i} \neq \frac{L_2(uv)}{uv}$ ;

– If  $u \in W \setminus \{1\}$  and  $(\frac{L_1(uv)}{uv})^{2^i} \neq \frac{L_2(v)}{v}$ , then  $\frac{L_1(v)^{2^i}(uv)+L_2(uv)v^{2^i}}{L_1(uv)^{2^i}v+L_2(v)(uv)^{2^i}} \notin \mathbb{F}_{2^m}^*$ .

**Proof** We need that, for any  $a \in \mathbb{F}_{2^n}^*$ , the function  $\Delta_a(x) = F(x+a) + F(x) + F(a) + F(0)$  is a 2-to-1 map, or equivalently, that  $\ker(\Delta_a(ax)) = \{0, 1\}$ . As showed before, we can rewrite  $a = st$  and  $x = uv$  with  $s, u \in \mathbb{F}_{2^m}^*$  and  $t, v \in W$ . Hence, since  $L_1$  and  $L_2$  are  $\mathbb{F}_{2^m}$ -polynomials, we have:

$$\begin{aligned} \Delta_a(ax) &= L_1(a)^{2^i}ax + L_2(a)(ax)^{2^i} + L_1(ax)^{2^i}a + L_2(ax)a^{2^i} \\ &= s^{2^i}L_1(t)^{2^i}st \cdot uv + sL_2(t)s^{2^i}t^{2^i} \cdot u^{2^i}v^{2^i} + s^{2^i}u^{2^i}L_1(tv)^{2^i}st + suL_2(tv)s^{2^i}t^{2^i} \\ &= us^{2^i+1}[(L_1(t)^{2^i}tv + L_2(tv)t^{2^i}) + u^{2^i-1}(L_2(t)t^{2^i}v^{2^i} + L_1(tv)^{2^i}t)]. \end{aligned}$$

Without loss of generality we can assume that  $s = 1$ . So,  $F$  is APN over  $\mathbb{F}_{2^n}$  if and only if  $u = 0$  or  $u = v = 1$  are the only solutions to  $\Delta_t(uvt) = 0$  for any  $t \in U$ .

If  $v = 1$ , then

$$\Delta_t(tx) = u(L_1(t)^{2^i}t + L_2(t)t^{2^i})[1 + u^{2^i-1}].$$

Since  $\gcd(i, m) = 1$ ,  $x^{2^i-1}$  is a permutation over  $\mathbb{F}_{2^m}$  and thus  $\ker(\Delta_t(tx)) = \{0, 1\}$  if and only if  $\frac{L_1(t)^{2^i}}{t^{2^i}} \neq \frac{L_2(t)}{t}$ .

Assume now that  $v \neq 1$ . Then, if  $L_2(t)t^{2^i}v^{2^i} + L_1(tv)^{2^i}t = 0$ , we have:

$$\Delta_t(tx) = u[(L_1(t)^{2^i}tv + L_2(tv)t^{2^i})].$$

This implies  $\frac{L_1(t)^{2^i}}{t^{2^i}} \neq \frac{L_2(tv)}{tv}$ .

If  $L_2(t)t^{2^i}v^{2^i} + L_1(tv)^{2^i}t \neq 0$ , then

$$[(L_1(t)^{2^i}tv + L_2(tv)t^{2^i}) + u^{2^i-1}(L_2(t)t^{2^i}v^{2^i} + L_1(tv)^{2^i}t)] = 0$$

implies  $u^{2^i-1} = \frac{L_1(t)^{2^i}tv+L_2(tv)t^{2^i}}{L_2(t)t^{2^i}v^{2^i}+L_1(tv)^{2^i}t}$ . Since  $x^{2^i-1}$  is a permutation over  $\mathbb{F}_{2^m}$  this equation admits a solution different from zero if and only if  $\frac{L_1(t)^{2^i}tv+L_2(tv)t^{2^i}}{L_2(t)t^{2^i}v^{2^i}+L_1(tv)^{2^i}t}$  is contained in  $\mathbb{F}_{2^m}^*$ .  $\square$

The obtained APN function (3) is of the form

$$F(x) = (A_0^{2^i} + B_0)x^{2^i+1} + \sum_{j=1}^{k-1} [A_j^{2^i}x^{2^i+jm+1} + B_jx^{2^i+jm+2^i}].$$

Let us see now necessary conditions on the linear functions  $L_1$  and  $L_2$  for  $F$  to be APN.

**Proposition 3.1** *Let  $n, L_1, L_2$  and  $F$  be as in Theorem 3.1. If  $F$  is APN over  $\mathbb{F}_{2^n}$ , then the following statements hold:*

- (i)  $\ker(L_1(x) + rx) \cap \ker(L_2(x) + r^{2^i}x) = \{0\}$  for any  $r \in \mathbb{F}_{2^n}$ ;
- (ii)  $|\ker(L_1(x)^{2^i} + rx) \cap \ker(L_2(x) + w^{2^i}x^{2^i})| \leq 2$  for any  $r, w \in \mathbb{F}_{2^n}$ ;
- (iii) If  $\ker(L_1) \cap \ker(L_2(x) + x) \neq \{0\}$ , then  $\ker(L_1(x) + x) \cap \ker(L_2) = \{0\}$ ;
- (iv)  $\ker(L_1(x) + rx^{2^j}) \cap \ker(L_2(x) + r^{2^i}x^{(2^j-1)2^i+1}) = \{0\}$  for any  $r \in \mathbb{F}_{2^n}$  and  $j \geq 0$ .

**Proof** For any nonzero  $a$ , we define the function  $\Delta_a(x) = F(x+a) + F(x) + F(a) + F(0)$  and, with  $t \in \mathbb{F}_{2^m}$ , we have

$$\begin{aligned}\Delta_a(x) &= aL_1(x)^{2^i} + xL_1(a)^{2^i} + x^{2^i}L_2(a) + a^{2^i}L_2(x), \\ \Delta_a(at) &= (t+t^{2^i})(aL_1(a)^{2^i} + a^{2^i}L_2(a)).\end{aligned}$$

Suppose there exists a non-zero  $a \in \ker(L_1(x) + rx) \cap \ker(L_2(x) + r^{2^i}x)$ . We clearly have  $a\mathbb{F}_{2^m} \subseteq \ker(\Delta_a)$ , but since  $m > 1$ , this contradicts  $|\ker(\Delta_a)| = 2$ . This establishes (i).

For (ii), suppose  $\{0, a, b\} \subset \ker(L_1(x)^{2^i} + rx) \cap \ker(L_2(x) + w^{2^i}x^{2^i})$ . Then

$$\Delta_a(b) = a(rb) + b(ra) + a^{2^i}(w^{2^i}b^{2^i}) + b^{2^i}(w^{2^i}a^{2^i}) = 0.$$

Next suppose  $a \in \ker(L_1) \cap \ker(L_2(x) + x)$ . Then we have  $\Delta_a(x) = a(L_1(x) + x)^{2^i} + a^{2^i}L_2(x)$ . Clearly any  $b \in \ker(L_1(x) + x) \cap \ker(L_2)$  satisfies  $\Delta_a(b) = 0$ . Since  $f$  is APN,  $\ker(\Delta_a) = \{0, a\}$ , so that  $\ker(L_1(x) + x) \cap \ker(L_2) \subset \{0, a\}$ . However,  $\ker(L_1) \cap \ker(L_1(x) + x) = \{0\}$ , so that no non-zero element of  $\mathbb{F}_{2^m}$  can lie in both  $\ker(L_1) \cap \ker(L_2(x) + x)$  and  $\ker(L_1(x) + x) \cap \ker(L_2)$ . This establishes (iii).

For (iv), suppose  $a \in \ker(L_1(x) + rx^{2^j}) \cap \ker(L_2(x) + r^{2^i}x^{(2^j-1)2^i+1})$  is non-zero. Then for any  $t \in \mathbb{F}_{2^m}$  we have

$$\begin{aligned}\Delta_a(ta) &= (t+t^{2^i})(aL_1(a)^{2^i} + a^{2^i}L_2(a)) \\ &= (t+t^{2^i})(ar^{2^i}a^{2^j2^i} + a^{2^i}r^{2^i}a^{(2^j-1)2^i+1}) = 0,\end{aligned}$$

so that  $a\mathbb{F}_{2^m} \subseteq \ker(\Delta_a)$ , a contradiction.  $\square$

### 3.2 The case $n = 8$

Applying the construction of Theorem 3.1 in dimension 8 with  $k = 4$  and  $m = 2$ , restricting the coefficients of  $L_1$  and  $L_2$  to the subfield  $\mathbb{F}_{2^4}$  we obtained several APN functions given in [16, Table 9] and one in [19, Table 6] which have not been previously identified as a part of any APN family. The functions mentioned are listed in Table 1.

The following results were obtained for  $n = 8$ .

- Considering generalized isotopic shifts of  $x^3$  it is possible to obtain maps EA-equivalent to nos. 1.2, 1.5, 1.7, 1.8, 1.10, 1.11, 1.12, 1.16, 1.17, 3.1 in Table 9 [16] and to no. 9 in Table 6 of [19].
- Considering generalized isotopic shifts of  $x^9$  it is also possible to obtain maps EA-equivalent to no. 1.3 Table 9 [16].

**Remark 3.1** The function no. 9 in Table 6 [19] has the same CCZ-invariants ( $\Gamma$ -rank,  $\Delta$ -rank and  $\mathcal{M}_{G_F}$ ) as the function number 1.9 in Table 9 of [16] (we note that the value of the  $\Gamma$ -rank given in [19] is not correct, indeed this function has  $\Gamma$ -rank = 14034).

Since two quadratic APN functions are CCZ-equivalent if and only if they are EA-equivalent [23], the CCZ-inequivalence between these two functions can be obtained by checking another invariant with respect to the EA-equivalence that we shall introduce in the next subsection.

**Table 1** APN polynomials over  $\mathbb{F}_{2^8}$  derived from Theorem 3.1

Functions	Equiv. to no. in Table 9 in [16]
$\zeta^{136}x^{66} + \zeta^{85}x^{33} + \zeta^{85}x^{18} + \zeta^{102}x^9 + \zeta^{221}x^6 + x^3$	No. 9 in Table 6 in [19]
$\zeta^{102}x^{66} + \zeta^{204}x^9 + x^3$	1.2
$\zeta^{153}x^{129} + \zeta^{204}x^{66} + \zeta^{170}x^{33} + \zeta^{85}x^{18} + \zeta^{204}x^6 + x^3$	1.5
$\zeta^{102}x^{129} + \zeta^{153}x^{66} + \zeta^{170}x^{33} + \zeta^{221}x^{18} + \zeta^{221}x^9 + \zeta^{187}x^6 + x^3$	1.7
$x^{66} + \zeta^{85}x^{33} + x^{18} + x^9 + x^3$	1.8
$\zeta^{204}x^{129} + \zeta^{170}x^{66} + \zeta^{153}x^{33} + \zeta^{85}x^{18} + \zeta^{153}x^9 + \zeta^{17}x^6 + x^3$	1.10
$\zeta^{204}x^{66} + x^{33} + x^{18} + \zeta^{153}x^9 + x^3$	1.11
$\zeta^{170}x^{129} + \zeta^{204}x^{66} + \zeta^{17}x^{33} + \zeta^{68}x^{18} + \zeta^{221}x^9 + \zeta^{204}x^6 + x^3$	1.12
$\zeta^{238}x^{129} + \zeta^{204}x^{66} + \zeta^{119}x^{33} + \zeta^{68}x^{18} + \zeta^{85}x^9 + \zeta^{119}x^6 + x^3$	1.16
$\zeta^{17}x^{129} + \zeta^{85}x^{66} + \zeta^{34}x^{33} + \zeta^{34}x^{18} + \zeta^{187}x^9 + \zeta^{187}x^6 + x^3$	1.17
$\zeta^{17}x^{129} + \zeta^{238}x^{66} + \zeta^{153}x^{33} + \zeta^{85}x^{18} + \zeta^{238}x^9 + \zeta^{102}x^6 + x^3$	3.1
$\zeta^{153}x^{129} + \zeta^{221}x^{72} + \zeta^{170}x^{33} + \zeta^{102}x^{24} + x^{12} + x^9 + \zeta^{136}x^3$	1.3

All are either new or correspond to the known but unclassified cases

### 3.3 A new EA-equivalence invariant

Let  $S(F) = \{b \in \mathbb{F}_{2^n} : \exists a \in \mathbb{F}_{2^n} \text{ s.t. } \mathcal{W}_F(a, b) = 0\}$ , where  $\mathcal{W}_F(a, b) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\text{Tr}(ax + bF(x))}$  is the Walsh coefficient of  $F$  in  $a$  and  $b$ . This set was used in [8] to study some relations between the CCZ-equivalence and the EA-equivalence.

It is easy to check that:

- if  $F'(x) = F(x) + L(x)$  with  $L$  linear, then  $b \in S(F)$  if and only if  $b \in S(F')$ .
- If  $F'(x) = A_1 \circ F \circ A_2(x)$  with  $A_1, A_2$  affine permutations, then  $b \in S(F)$  if and only if  $\bar{A}_1^*(b) \in S(F')$ , where  $\bar{A}_1^*$  is the adjoint operator of the linear map  $A_1(x) + A_1(0)$ .

From this we have the following.

**Proposition 3.2** *Let  $N_i$  be the number of the  $\mathbb{F}_2$ -vector subspaces of  $\mathbb{F}_{2^n}$  contained in  $S(F)$  of dimension  $i$ . Then, the values  $N_i$  for  $i = 1, \dots, n$  are EA-invariant.*

**Proof** If  $F'$  is EA-equivalent to  $F$ , then there exist  $A_1, A_2$  affine permutations and  $L$  linear such that  $F'(x) = A_1 \circ F \circ A_2(x) + L(x)$ . From the arguments above, denoting  $\bar{A}_1(x) = A_1(x) + A_1(0)$  we have that  $S(F') = \bar{A}_1^*(S(F))$ . □

**Remark 3.2** We computed the values  $N_i$  for the two functions and we got  $N_1 = 86, N_2 = 340$  and  $N_3 = 4$  for the new function, and  $N_1 = 86, N_2 = 340$  and  $N_3 = 8$  for the function number 1.9. Thus from Proposition 3.2 we have that the two functions are not EA-equivalent.

**Remark 3.3** Note that when  $n$  is odd, a quadratic APN function  $F$  is Almost Bent (i.e. for all  $b \in \mathbb{F}_{2^n}^*$  we have  $\{\mathcal{W}_F(a, b) : a \in \mathbb{F}_{2^n}\} = \{0, \pm 2^{(n+1)/2}\}$ ), which implies  $S(F) = \mathbb{F}_{2^n}$ . Thus, such invariant cannot be used for testing the CCZ-equivalence of quadratic APN functions in the case  $n$  odd.

**Remark 3.4** In fact, this EA-invariant was tackled independently by Göloğlu and Pavlů in [17]. In their work, they focused on plateaued functions and looked at the subspaces in the set  $\{b : \mathcal{W}_F(0, b) \neq \pm 2^{n/2}\}$  ( $n$  even). For plateaued functions, this set coincides with  $S(F)$ .

### 3.4 The case $n = 9$

For the case  $k = m = 3$  we consider the generalized linear shift as in (3) with  $L_1$  and  $L_2$  having coefficients in the subfield  $\mathbb{F}_{2^3}$ . In Table 2 we list all known APN functions for  $n = 9$ , as reported in [6, Table 1]. In Table 3, we list all new APN functions obtained from Theorem 3.1. We can observe that the family of Theorem 3.1 covers the only known example of APN function for  $n = 9$ , function 8.1 of Table 11 in [16], which has not been previously identified as a part of an APN family. Hence, currently, we do not have any known example of APN functions for  $n = 9$  which would not be covered by an APN family. Note that this latter function was not obtained from the approach studied in [16] (it does not belong to a switching class of a previously known APN map). Finally, Table 3 indicates 15 new APN functions all obtained from Theorem 3.1. In both tables we include, for each function, the CCZ-invariants  $\Gamma$ -rank,  $\Delta$ -rank and  $|\mathcal{M}_{GF}|$ .

The CCZ-inequivalence of some of these functions was obtained by checking with MAGMA the equivalence of some linear code which can be associated to an APN function (see [4]).

### 3.5 Isotopic shifts with nonlinear functions

In this section we consider the case when the function used in the shift is not necessarily linear.

In [6], it has been proved that, in even dimension, an isotopic shift of the Gold function with a linear function defined over  $\mathbb{F}_2[x]$  cannot be APN. In the following, we show that for any quadratic function  $F$  in even dimension, we cannot obtain APN functions by shifting  $F$  with a polynomial whose coefficients belong to  $\mathbb{F}_2$ .

**Proposition 3.3** *For two integers  $k$  and  $m$  let  $n = km$  and  $q = 2^k$ . Consider a function  $F \in \mathbb{F}_{2^n}[x]$  of the form*

$$F(x) = \sum_{i < j} b_{ij} x^{q^i + q^j} + \sum_i b_i x^{2^i} + c,$$

*If  $n$  is even or  $k > 1$ , then any isotopic shift  $F_L$  with  $L \in \mathbb{F}_{2^k}[x]$  cannot be APN. In particular, this holds for any quadratic function  $F \in \mathbb{F}_{2^n}[x]$  with  $n$  even and  $L \in \mathbb{F}_2[x]$ .*

**Proof** For  $F$  and  $L$  as outlined, we have

$$F_L(x) = \sum_{i < j} b_{ij} [x^{q^i} L(x)^{q^j} + x^{q^j} L(x)^{q^i}] + c$$

and  $L(x^q) = L(x)^q$ . Note that for any  $x \in \mathbb{F}_{2^k}$ ,  $F_L(x) = c$ . For  $a \in \mathbb{F}_{2^n}$ , we set  $\Delta_a(x) = F_L(x+a) + F_L(x) + F_L(a) + F_L(0)$ .

If  $k > 1$ , then  $\Delta_a(x) = 0$  for all  $x, a \in \mathbb{F}_{2^k}$ , so that  $F_L$  is not APN. If  $\mathbb{F}_4 = \{0, 1, \alpha, \alpha + 1\} \subseteq \mathbb{F}_{2^n}$ , then consider  $\Delta_\alpha(x)$ . Clearly  $\Delta_\alpha(0) = 0$ , while it is easily observed that  $\Delta_\alpha(\alpha +$



**Table 2** Previously known CCZ-inequivalent APN polynomials over  $\mathbb{F}_{2^9}$  and their relation to previously known families of APN functions

Functions	Families	no. Table 11 in [16]	$\Gamma$ -rank	$\Delta$ -rank	$ \mathcal{M}_{GF} $
$x^3$	Gold	1.1	38470	872	$9 \cdot 2^9 \cdot 511$
$x^5$	Gold	2.1	41494	872	$9 \cdot 2^9 \cdot 511$
$x^{17}$	Gold	3.1	38470	872	$9 \cdot 2^9 \cdot 511$
$x^{13}$	Kasami	4.1	58676	3086	$9 \cdot 511$
$x^{241}$	Kasami	6.1	61726	3482	$9 \cdot 511$
$x^{19}$	Welch	5.1	60894	3956	$9 \cdot 511$
$x^{255}$	Inverse	7.1	130816	93024	$2 \cdot 9 \cdot 511$
$Tr_1^9(x^9) + x^3$	[9]	1.2	47890	920	$9 \cdot 2^9$
$Tr_3^9(x^{18} + x^9) + x^3$	[10]	1.3	48428	930	$9 \cdot 2^9$
$Tr_3^9(x^{36} + x^{18}) + x^3$	[10]	1.4	48460	944	$9 \cdot 2^9$
$x^3 + x^{10} + \zeta^{438} x^{136}$	-	8.1	48608	938	$3 \cdot 7 \cdot 2^9$
$\zeta^{337} x^{129} + \zeta^{424} x^{66} + \zeta^2 x^{17} + \zeta x^{10} + \zeta^{34} x^3$	[6]	-	48596	944	$3 \cdot 7 \cdot 2^9$

**Table 3** APN polynomials over  $\mathbb{F}_{2^9}$  derived from Theorem 3.1

$\mathcal{G}_i$	Function	Eq. to known ones	$\Gamma$ -rank	$\Delta$ -rank	$ \mathcal{M}_{GF} $
$i = 1$	$x^{129} + \zeta^{146}x^{66} + x^{17} + \zeta^{365}x^{10} + x^3$	eq. to APN function in [6]	48596	944	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{219}x^{129} + \zeta^{292}x^{66} + \zeta^{292}x^{17} + \zeta^{219}x^{10} + x^3$	new	48506	936	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{365}x^{129} + \zeta^{292}x^{66} + \zeta^{365}x^{17} + \zeta^{73}x^{10} + x^3$	new	48610	938	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{365}x^{129} + \zeta^{365}x^{66} + \zeta^{146}x^{17} + \zeta^{365}x^{10} + x^3$	new	48612	938	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{365}x^{129} + \zeta^{219}x^{66} + \zeta^{292}x^{17} + \zeta^{73}x^{10} + x^3$	new	48548	928	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{73}x^{129} + \zeta^{365}x^{66} + \zeta^{73}x^{17} + \zeta^{73}x^{10} + x^3$	new	48548	928	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{365}x^{129} + \zeta^{438}x^{66} + \zeta^{292}x^{10} + x^3$	new	48506	936	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{365}x^{129} + x^{66} + \zeta^{438}x^{10} + x^3$	new	48604	928	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{73}x^{129} + \zeta^{292}x^{66} + x^{10} + x^3$	new	48564	942	$2^9 \cdot 3 \cdot 7$
$i = 1$	$\zeta^{73}x^{129} + x^{66} + \zeta^{219}x^{17} + x^3$	new	48604	928	$2^9 \cdot 3 \cdot 7$
$i = 2$	$\zeta^{146}x^{257} + \zeta^{438}x^{68} + \zeta^{438}x^{12} + x^5$	new	48546	938	$2^9 \cdot 3 \cdot 7$
$i = 2$	$\zeta^{146}x^{257} + \zeta^{365}x^{33} + \zeta^{365}x^{12} + x^5$	eq. to 8.1	48608	938	$2^9 \cdot 3 \cdot 7$
$i = 2$	$\zeta^{73}x^{257} + \zeta^{146}x^{68} + x^{33} + x^5$	new	48564	942	$2^9 \cdot 3 \cdot 7$
$i = 2$	$\zeta^{365}x^{257} + \zeta^{438}x^{68} + \zeta^{365}x^{33} + \zeta^{438}x^{12} + x^5$	new	48594	944	$2^9 \cdot 3 \cdot 7$
$i = 2$	$\zeta^{146}x^{257} + \zeta^{219}x^{68} + \zeta^{73}x^{33} + x^{12} + x^5$	new	48520	932	$2^9 \cdot 3 \cdot 7$
$i = 2$	$\zeta^{73}x^{257} + \zeta^{219}x^{68} + \zeta^{365}x^{33} + x^5$	new	48602	938	$2^9 \cdot 3 \cdot 7$
$i = 4$	$\zeta^{292}x^3 + \zeta^{146}x^{80} + \zeta^{73}x^{24} + x^{17}$	new	48520	932	$2^9 \cdot 3 \cdot 7$

All, except for the first one, are either new or correspond to the one known but unclassified case

$1) = \Delta_\alpha(1)$ . We have

$$\begin{aligned} \Delta_\alpha(\alpha + 1) &= F_L(\alpha + 1) + F_L(\alpha) + F_L(1) + c \\ &= \sum_{i < j} b_{ij} [L(\alpha + 1)^{q^i} (\alpha + 1)^{q^j} + (\alpha + 1)^{q^i} L(\alpha + 1)^{q^j} \\ &\quad + L(\alpha)^{q^i} \alpha^{q^j} + \alpha^{q^i} L(\alpha)^{q^j}] \\ &= \sum_{i < j} b_{ij} [L(\alpha + 1)(\alpha + 1)^{q^{j-i}} + (\alpha + 1)L(\alpha + 1)^{q^{j-i}} \\ &\quad + L(\alpha)\alpha^{q^{j-i}} + \alpha L(\alpha)^{q^{j-i}}]^{q^i}. \end{aligned}$$

When  $j - i$  is odd and  $\mathbb{F}_4 \not\subseteq \mathbb{F}_{2^k}$ , the term in the sum is zero as  $\alpha^{q^{j-i}} = \alpha^2 = \alpha + 1$ ,  $L(\alpha)^{q^{j-i}} = L(\alpha + 1)$  and  $L(\alpha + 1)^{q^{j-i}} = L(\alpha)$ . If  $j - i$  even or  $\mathbb{F}_4 \subseteq \mathbb{F}_{2^k}$ , then the term in the sum is again zero due to the fact that  $\alpha^{q^{j-i}} = \alpha$  and  $L(\alpha)^{q^{j-i}} = L(\alpha)$ . In either case, we have  $\Delta_\alpha(x) = 0$  for  $x = 0, 1, \alpha + 1$ , so  $F_L$  is not APN. □

### 3.5.1 Nonlinear shift for the Gold functions

If we consider an isotopic shift of a Gold function without the restriction that  $L(x)$  is a linear function, then  $L(x) = \sum_{j=0}^{2^n-1} c_j x^j$  and the isotopic shift will be of the form

$$\mathcal{G}_{i,L}(x) = x^{2^i} L(x) + xL(x)^{2^i}. \tag{4}$$

We have  $\mathcal{G}_{i,L}(x^2)^{2^{-1}} = x^{2^i} M(x) + xM(x)^{2^i}$ , where  $M(x) = \sum_{j=0}^{2^n-1} c_j x^j$ , and also  $\zeta^{-2^i-1} \mathcal{G}_{i,L}(\zeta x) = x^{2^i} N(x) + xN(x)^{2^i}$ , where  $N(x) = \sum_{j=0}^{2^n-1} c_j \zeta^{j-1} x^j$ . Hence we obtain the following.

**Proposition 3.4** *Let  $\mathbb{F}_{2^n}^* = \langle \zeta \rangle$ . Assume that  $\mathcal{G}_{i,L}$  is constructed with  $L(x) = \sum_{j=0}^{2^n-1} c_j x^j$ . Then, for any integers  $k, t$ , we have that  $\mathcal{G}_{i,L}$  is linear equivalent to  $\mathcal{G}_{i,M}$ , where  $M(x) = \sum_{j=0}^{2^n-1} (c_j \zeta^{k(j-1)})^{2^t} x^j$ .*

As for the linear shifts, it is possible to restrict the search of one possible non-zero coefficient of the function.

In the following table we recall the list of known APN power maps (the list was conjectured to be complete in [14]).

In odd dimension it is possible to obtain all the power APN functions, except the Dobbertin functions, as the isotopic shifts of a Gold function by a monomial.

**Theorem 3.2** *Over  $\mathbb{F}_{2^n}$  with  $n$  an odd integer, let  $F$  be any known APN power function outside the class of Dobbertin functions. Then there exists a monomial  $L(x) = ax^d$  and a Gold function  $\mathcal{G}_i = x^{2^i+1}$  such that the shift  $\mathcal{G}_{i,L}$  is EA-equivalent to  $F$ .*

**Proof** As shown in Table 4, excluding the Dobbertin function, the known APN power functions are the Gold functions, the Kasami functions, the Welch function, the Niho functions and the inverse function. In the following we will show that it is possible for any of the mentioned functions, to construct an isotopic shift of a Gold function that is EA-equivalent to it.

**Table 4** Known APN power functions  $x^d$  over  $\mathbb{F}_{2^n}$

Functions	Exponents $d$	Conditions	Degree	Proven
Gold	$2^i + 1$	$\gcd(i, n)=1$	2	[18,22]
Kasami	$2^{2i} - 2^i + 1$	$\gcd(i, n)=1$	$i+1$	[20,21]
Welch	$2^t + 3$	$n = 2t + 1$	3	[13]
Niho	$2^t + 2^{\frac{t}{2}} - 1, t \text{ even}$	$n = 2t + 1$	$\frac{t+2}{2}$	[14]
	$2^t + 2^{\frac{3t+1}{2}} - 1, t \text{ odd}$		$t+1$	
Inverse	$2^{2t} - 1$	$n = 2t + 1$	$n - 1$	[2,22]
Dobbertin	$2^{4i} + 2^{3i} + 2^{2i} + 2^i - 1$	$n = 5i$	$i + 3$	[15]

1. Consider the Kasami function  $x^{2^{2t}-2^t+1}$ . If  $t$  is odd, then let  $i$  be an integer such that  $n = 2i + t$ . Then, considering  $L(x) = ax^{2^{n-i}+2^{n-i+1} \dots + 2^{n-i+t-1}}$  we have

$$\begin{aligned} \mathcal{G}_{i,L}(x) &= a^{2^i} x^{2^t} + ax^{2^{n-i}+2^{n-i+1} \dots + 2^{n-i+t-1} + 2^i} \\ &= a^{2^i} x^{2^t} + ax^{2^i(2^t+2^{t+1} \dots + 2^{2t-1}+1)} \\ &= a^{2^i} x^{2^t} + ax^{2^i(2^{2t}-2^t+1)}. \end{aligned}$$

If  $t$  is even, let  $i$  be an integer such that  $t = 2i$ . Then, with  $L(x) = ax^{2^{i+2^{i+1}}+\dots+2^{3i-1}}$  we have  $\mathcal{G}_{i,L}(x) = a^{2^i} x^{2^{2t}-2^t+1} + ax^{2^{3i}}$ .

2. For the inverse function,  $x^{2^n-2}$ , considering  $L(x) = ax^{2^{2t}-2}$ , where  $t$  is such that  $n = 2t + 1$ , we have  $\mathcal{G}_{1,L}(x) = a^2 x^{2^{(2^n-2)}} + ax^{2^{2t}}$ .

3. Let  $n = 2t + 1$  and consider the Welch function  $x^{2^t+3}$ . If  $t$  is odd, then consider  $i$  such that  $t = 2i - 1$ . With  $L(x) = ax^{2^i+2^{2i+1}}$  we obtain  $\mathcal{G}_{i,L}(x) = a^{2^i} x^{2^{2i}(2^{2i-1}+3)} + ax^{2^{i+2}}$ . If  $t$  is even, then consider  $i$  such that  $t = 2i$ . Using  $L(x) = ax^{2^{3i+1}+2^{3i+2}}$  we obtain  $\mathcal{G}_{i,L}(x) = a^{2^i} x^4 + ax^{2^{3i+1}(2^{2i}+3)}$ .

4. For  $n = 2t + 1$ , with  $t$  odd, let  $t = 2i - 1$ . Then, with  $L(x) = ax^{2^n-2^i}$  we obtain that

$$\begin{aligned} \mathcal{G}_{i,L}(x) &= a^{2^i} x^{2^i-2^{2i}+1} + ax = a^{2^i} x^{2^{2i}(2^{-i}+2^{-2i}-1)} + ax \\ &= a^{2^i} x^{2^{2i}(2^{3i-1}+2^{2i-1}-1)} + ax = a^{2^i} x^{2^{2i}(2^{(3t+1)/2}+2^t-1)} + ax \end{aligned}$$

is equivalent to the Niho function (indeed  $(3t + 1)/2 = (6i - 3 + 1)/2 = 3i - 1$ ). If  $t$  is even, let  $t = 2i$ . Then with  $L(x) = ax^{2^{n-i}+2^{n-i+1} \dots + 2^{n-1}}$

$$\begin{aligned} \mathcal{G}_{i,L}(x) &= a^{2^i} x^{2^i} + ax^{2^{n-i}+2^{n-i+1} \dots + 2^{n-1} + 2^i} \\ &= a^{2^i} x^{2^i} + ax^{2^{n-i}(1+2 \dots + 2^{i-1}+2^{2i})} \\ &= a^{2^i} x^{2^i} + ax^{2^{n-i}(2^i-1+2^{2i})} \end{aligned}$$

is equivalent to the Niho function.

5. Let  $n = 2i + 1$  and  $j$  be an integer such that  $\gcd(n, j) = 1$ . Then with  $L(x) = ax^{2^{i+j}-2^i}$

$$\begin{aligned} \mathcal{G}_{i,L}(x) &= a^{2^i} x^{2^{2i+j}-2^{2i}+1} + ax^{2^{i+j}} = a^{2^i} x^{2^{2i}(2^j+2^{-2i}-1)} + ax^{2^{i+j}} \\ &= a^{2^i} x^{2^{2i}(2^j+1)} + ax^{2^{i+j}} \end{aligned}$$

is equivalent to the Gold function with parameter  $j$ . □

**Remark 3.5** From computational results, for  $n$  even, it seems that it is not possible to obtain APN functions as the isotopic shifts of a Gold map by (non-linear) monomials. The search has been performed for  $n = 4, 6, 8, 10$ , considering also non-APN Gold exponents.

**Problem 3.1** Is it possible to obtain the Dobbertin function as an isotopic shift of a Gold function by a non-linear  $L$ ?

**Problem 3.2** Is it possible to obtain the same result for  $n$  an even integer and  $L$  a non-linear multinomial?

## 4 Conclusions

Starting from the work [6], we introduced some generalizations of the isotopic shift construction for the case when the initial function is a Gold power function. In particular, using a generalized form of the isotopic shift with  $\mathbb{F}_{2^m}$ -polynomials, we were able to construct a general family of quadratic APN functions. This allowed us to classify into a family some of the previously known unclassified examples of APN functions for  $n = 8, 9$ , and to provide new APN functions on  $\mathbb{F}_{2^9}$ . The computations performed were restricted to linear maps with coefficients in the subfield. We expect that, without such restriction, it is possible to find new APN functions.

We also investigated the case of constructing an isotopic shift with a nonlinear function. In this case, for any odd  $n$ , we can obtain all known power APN functions (except the Dobbertin ones) using a nonlinear monomial function.

**Acknowledgements** The research of this paper was supported by Trond Mohn Foundation.

**Funding** Open Access funding provided by University of Bergen.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Albert, A.A.: Finite division algebras and finite planes, *Combinatorial Analysis: Proceedings of the 10th Symposium in Applied Mathematics (Providence)*, Symposia in Applied Mathematics, vol. 10, American Mathematical Society, pp. 53–70 (1960).
2. Beth T., Ding C.: On almost perfect nonlinear permutations, *Advances in Cryptology-EUROCRYPT'93*, vol. 765. Lecture Notes in Computer Science. Springer, New York (1993).
3. Biham E., Shamir A.: Differential cryptanalysis of DES-like cryptosystems. *J. Cryptol.* **4**(1), 3–72 (1991).
4. Browning K.A., Dillon J.F., Kibler R.E., McQuistan M.T.: APN polynomials and related codes. *J. Comb. Inf. Syst. Sci.* **34**(1–4), 135–159 (2009).
5. Budaghyan L.: *Construction and Analysis of Cryptographic Functions*. Springer, Cham (2014).
6. Budaghyan L., Calderini M., Carlet C., Coulter R., Villa I.: Constructing APN functions through isotopic shifts. *IEEE Trans. Inform. Theory* (2020). <https://doi.org/10.1109/TIT.2020.2974471>.
7. Budaghyan, L., Calderini, M., Carlet, C., Coulter, R., Villa, I.: Construction On Isotopic Shift, for Planar Functions, : *IEEE International Symposium on Information Theory (ISIT)*. Paris, France **2019**, 2962–2966 (2019).

8. Budaghyan L., Calderini M., Villa I.: On relations between CCZ- and EA-equivalences. *Cryptogr. Commun.* **12**, 85–100 (2020).
9. Budaghyan L., Carlet C., Leander G.: Constructing New APN functions from known ones. *Finite Fields Appl.* **15**, 150–159 (2009).
10. Budaghyan, L., Carlet, C., Leander, G.: On a Construction of Quadratic APN Functions., *Proceedings of IEEE Information Theory workshop ITW'09*, pp. 374–378 (2009).
11. Carlet, C.: Vectorial Boolean Functions for Cryptography, *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, pp. 398–470 (2010).
12. Carlet C., Charpin P., Zinoviev V.: Bent functions and permutations suitable for DES-like cryptosystems. *Des. Codes Cryptogr.* **15**(2), 125–156 (1998).
13. Dobbertin H.: Almost perfect nonlinear power functions over  $GF(2^n)$ : the Welch case. *IEEE Trans. Inform. Theory* **45**, 1271–1275 (1999).
14. Dobbertin H.: Almost perfect nonlinear power functions over  $GF(2^n)$ : the Niho case. *Inform. Comput.* **151**, 57–72 (1999).
15. Dobbertin, H.: Almost perfect nonlinear power functions over  $GF(2^n)$ : a new case for  $n$  divisible by 5, *Proceedings of Finite Fields and Applications FQ5*, pp. 113–121 (2000).
16. Edel E., Pott A.: A new almost perfect nonlinear function which is not quadratic. *Adv. Math. Commun.* **3**, 59–81 (2009).
17. Göloğlu, F., Pavlu, J.: Search for APN permutations among known APN functions. Presented at BFA2019.
18. Gold R.: Maximal recursive sequences with 3-valued recursive cross-correlation functions. *IEEE Trans. Inform. Theory* **14**, 154–156 (1968).
19. Guobiao, W., Tan, Y., Gong G.: *On quadratic almost perfect nonlinear functions and their related algebraic object*. *Proceedings of Workshop on Coding and Cryptography 2013*.
20. Janwa, H., Wilson, R.: Hyperplane sections of Fermat varieties in  $P^3$  in char. 2 and some applications to cycle codes. In: *Proceedings of AAECC-10, LNCS*, vol. 673, Springer, Berlin, pp. 180–194 (1993).
21. Kasami T.: The weight enumerators for several classes of subcodes of the second order binary Reed-Muller codes. *Inform. Control* **18**, 369–394 (1971).
22. Nyberg, K.: Differentially uniform mappings for cryptography, *Advances in Cryptography, EURO-CRYPT'93. Lecture Notes in Computer Science* **765**, 55–64 (1994).
23. Yoshiara S.: Equivalences of quadratic APN functions. *J. Algebr. Comb.* **35**(3), 461–475 (2012).

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.