

# Norwegian Index for Emergency Medical Assistance

Studies on the Use and Precision of the Emergency Medical  
Dispatch Guidelines in Norway

**Eirin Nybø Ellensen**



Thesis for the degree of philosophiae doctor (PhD)  
at the University of Bergen

2017

Date of defence: 10.08.2017

© Copyright Eirin Nybø Ellensen

The material in this publication is protected by copyright law.

Year: 2017

Title: Norwegian Index for Emergency Medical Assistance  
Studies on the Use and Precision of the Emergency Medical Dispatch  
Guidelines in Norway

Author: Eirin Nybø Ellensen

Print: AiT Bjerch AS / University of Bergen



## Scientific environment

This project was carried out within the doctoral education program at the Faculty of Medicine and Dentistry and the Department of Global Public Health and Primary Care, University of Bergen, Norway.

While being employed by the Norwegian Air Ambulance Foundation, Department of Research, my everyday work has been carried out at the National Centre for Emergency Primary Health Care, Uni Research Health, in Bergen.



## Acknowledgements

First and foremost a huge thank you to my supervisors:

- **Erik Zakariassen** – Thank you for your patience and encouragement, and consistently thorough and constructive supervision throughout all those long years.
- **Torben Wisborg** – Thank you for keeping up my spirits, and for providing a slightly more emergency medical glance upon my work.
- **Steinar Hunsjørn** – Thank you for your ability to see the big picture, and in a magic way make me solve the impossible puzzles.

My co-workers at the National Air Ambulance Foundation and National Centre for Emergency Primary Health Care – Thank you for both scientific and good times!

I would like to thank the members supporting the Norwegian Air Ambulance Foundation for making our research possible. This project has also been made possible through funding by the Norwegian ExtraFoundation for Health and Rehabilitation. Thank you.

A huge thank you to all the EMCCs in Norway, including operators and management. Without you this project would never have been possible. A special thank you to Bergen EMCC, and Tone Alfsvåg Engelsen and Lars Myrnes.

Halvor Næss – Thank you for providing NorStroke data and making the last study on Index validity possible.

Vegard – From the outside, it may appear as though this thesis has been written despite you. We both know that's not entirely true. Thank you for never stopping to believe in me. And Brage, Marcus, Stella Olivia and August – thank you for being my precious ones.

## **Preface**

Handling an emergency medical situation by phone, and phone only, requires multiple simultaneous tasks: get a rapid overview of the situation and patient, determine the problem and dispatch an adequate response, alert prehospital doctors on call, casualty clinics or hospital, and instruct the caller on how to perform first aid. All while keeping cool.

The Norwegian Index for Emergency Medical Assistance (Index) provides the guidelines supporting the emergency medical dispatcher during an emergency 113 call handling. The Index was developed in 1994, and is used by all the 19 Emergency medical communication centres in Norway. It has also been translated, adapted and implemented in several European countries.

However, the Index has never been validated, and there has been no knowledge about its ability to identify the true patients and to provide these with the proper resources. It was considered well-functioning, and all complaints made to the authorities regarding dispatch issues were rejected as long as Index had been used during the call. Meanwhile, the actual use of the guidelines was never measured.

Medical triaging by phone is difficult. Nonetheless, it is of utmost importance for both the individual patient and the society as a whole that the triage is as expedient as possible. The acutely ill or severely injured patient needs acute response, highly-skilled resources and transport directly to regionalized hospitals for treatment, while the moderately ill patient may receive good care at the local casualty clinic. The prehospital resources are limited and need to be utilized properly.

This project was initiated to obtain the necessary data to clarify the use of the Index and start a validation process of its ability as dispatch guidelines.

## Abbreviations and terminology

113	The medical emergency telephone number. Toll free, and the public's access point to acute medical care. Answered by the EMCCs.
116 117	The medical urgency telephone number. Toll free, and the public's access point to urgent medical care. Answered by the LEMCs.
AMIS	The software program used by the EMCC to register and keep record of all contacts. Contains information on patient/situation, location, caller phone number, Index criteria code, urgency, response, resources utilized and time events.
Ambulance-/GP-alarm	Simultaneous notification of both ambulance and GP on call. Should be released as a rule for all acute responses.
Casualty clinic (Legevaktlokale)	The actual premise of the out-of-hours services.
CBD	Criteria based dispatch. An emergency medical dispatch system using lists of criteria/prompts to determine a response. As a contrast to algorithmic dispatch systems.
Dispatch	1. Term used specifically about sending out an ambulance, or other resources.

2. Term used in general for the whole process of receiving emergency call, triage, deciding upon a response, and then dispatching adequate resource.

EMCC (AMK-sentral)	Emergency medical communication centre (Akuttmedisinsk kommunikasjonsentral). Receives and handles emergency medical 113 calls from the public, health line calls from prehospital emergency primary health care. Coordinate and dispatch the ambulance fleet. Manned by operators and resource coordinators.
EMD	Emergency medical dispatch. Term used to describe the dispatch process.
EMD response interval (AMK reaksjonstid)	Term used to describe the time interval between answering the call and dispatching the adequate resource.
EMS	Emergency medical services. Term used to describe the system providing rapid and appropriate medical care to victims of sudden illness or injuries. Includes both prehospital and in-hospital resources.
GP (Fastlege)	General practitioner. The term used for physicians working at primary care level. Work on contract with the municipality as regular GPs in office hours, and take shifts on call in the out-of-hours services.
Health trusts	The central government owns the secondary health care services through four regional health trusts; North, Middle, West and South-East.

HEMS	Helicopter emergency medical services. Norway has six rescue helicopters and 13 ambulance helicopters, all staffed by anesthesiologists. Each helicopter base also has an emergency car for missions when use of helicopter is inappropriate: incident in close proximity to the base, poor weather conditions, no flight time left. HEMS is part of the secondary health care system, alongside the ambulance service.
ICD-10	International classification of diseases. A classification of diseases and other health problems, mainly used by the secondary health care system in Norway. Provided by the World Health Organization and last revised 1990.
Index	The Norwegian Index for Emergency Medical Assistance (Norsk indeks for medisinsk nødhjelp). The dispatch guidelines used in Norway since 1994. Revised in 2009, 3 <sup>rd</sup> edition.
LEMC (Legevaktsentral)	Local emergency medical communication centre. Receives and handles urgent medical 116 117 calls from the public. Usually co-localized with a casualty clinic.
NPV	Negative predictive value. The proportion of true negatives among all test negatives.
Operator	The person answering and handling the 113 call at the EMCC. Usually a registered nurse with extra EMCC training.
OR	Odds ratio. The ratio of the odds of an event occurring in

	one group to the odds of it occurring in another group.
OOH (Legevaktjenesten)	Term used for organized out-of-hours emergency primary care service in the municipalities. Staffed by nurses and GPs on call. Handles the majority of the urgent and non-urgent patients.
PPV	Positive predictive value. The proportion of true positives among all positives test outcomes.
Resource coordinator	The person coordinating and dispatching the ambulance fleet at the EMCC. Also handling health lines from prehospital emergency primary care services. Usually an ambulance worker with extra EMCC training.
Response	<p>Each Index criteria code has a pre-determined recommended response which includes urgency and resource:</p> <p>Acute response = Red response (Akutt). Presumably life-threatening situation. GP on call is to be notified and ambulance dispatched immediately. Ambulance may drive with lights and sirens. Consider notifying HEMS.</p> <p>Urgent response = Yellow response (Haster). Potentially life-threatening situation. Cooperation with LEMC. Consider ambulance dispatch.</p> <p>Non-urgent response = Green response (Vanlig). Transfer to LEMC.</p>

Sensitivity	The likelihood of a condition positive being identified as a positive in a test for the condition.
Specificity	The likelihood of a condition negative being identified as a negative in a test for the condition.



## Summary

The main aim of this thesis was to initiate a validation process of the Norwegian Index for Emergency Medical Assistance (Index). Before we could do that, however, we had to explore to which extent the Index actually is used by the Norwegian emergency medical communication centres (EMCCs).

The Index is the emergency medical dispatch guidelines used by operators at EMCC when they receive and handle calls at the emergency medical line. In Norway, 113 is the specific toll-free line for medical emergencies. The Index was developed in 1994, and is now available in the 3rd edition (2009). All EMCCs and a majority of the local emergency medical communication centres (LEMCCs) in Norway claim to use the Index upon receipt of 113 calls. In 2010, knowledge on the epidemiology of 113 calls was non-existing.

Criteria-based dispatch guidelines are designed to be used in close cooperation with experienced health educated EMCC operators, and the operators are intended to bring their knowledge, experience and skill into the assessment process. This has complicated past research, as it is difficult to determine whether the determined criteria code or dispatched response is due to the guidelines themselves, the operator's own assessment or a combination of both. In addition, the Index is still paper-based, so that its use remains unable to track.

After the initial mapping of the place, situation and the patient's vital functions found on the start page, the rest of the Index criteria cards are symptom-based. This makes it challenging to compare specific Index criteria codes with specific discharge diagnoses from hospitals, as a measure on Index validity. Criteria card "27 Altered levels of consciousness – paralysis", specifically the criteria codes A. 27.03–06, was considered comparable to the diagnosis of stroke, including intracerebral haemorrhage, acute ischemic stroke and transient ischemic attack.

The objectives of the individual studies were to gain more knowledge about:

- The 113 epidemiology in Norway
- Use of the Index by the EMCCs
- The Index's validity in identification and prediction of stroke

**Study 1** was a cross-sectional survey of all 113-enquiries to the 19 EMCCs during a 72-hour period in August 2011. The final material consisted of 2 298 printouts from the electronic EMCC records, Emergency medical information system (AMIS), and contained information about time, caller, patient, Index criteria code, response and resources allocation. The national mean 113 contact rate was 56/1 000 population per year, but the variation among the different EMCCs was from 33 to 114. The acute contact rate was 21/1 000 per year, with a variation of 5 to 31. Urgency distribution showed 37% acute, 34% urgent and 27% non-urgent contacts. The most frequently used Index criteria card was "06 Unclear problem", which was used in 20% of the contacts, with a variation of 10 to 42% among the EMCCs.

**Study 2** was a national questionnaire study to all EMCC operators, to explore self-reported use of the Index. The questionnaire contained questions about use of the Index, education, EMCC experience, and training in, and repetition on the use of the Index at their workplace. Response rate was 63.4%, and the typical operator among the respondents was a female registered nurse with six years of EMCC experience, who worked in rotation with either the emergency department or ambulance. Mean self-reported use of the Index was calculated to be 3.95, corresponding to the response format 4 = "often, > 75%". Rotation with ambulance reduced use of the Index, while experienced focus on use of the Index at workplace increased the use. The operators checked whether the patient was conscious or not in 93% of the calls, but claimed to use the start page in only 47%. Availability of the electronic AMIS was the most frequent stated reason for not using the start page or the Index as a whole.

**Study 3** was a review of the audio logs from the same time period as **Study 1**, to assess use of the Index in an objective manner, and then see if the guideline adherence

affected the emergency medical dispatch (EMD) response interval. Seven EMCCs participated, with a total of 299 randomized calls, 174 acute and 125 urgent. The EMCC selection was strategic, based on variation in size, geographical location, various health trusts, and the EMCCs' mean self-reported use of the Index from **Study 2**. The listening form measured whether and how fast the operator got various indicators on use of the Index confirmed. The indicators "location", "consciousness" and "criteria compliance" were combined in an overall guideline adherence variable, where 0 indicated no guideline adherence and 3 the maximum guideline adherence. Mean guideline adherence was 2.41, equivalent to 80% of the maximum score. EMD response interval increased with decreased guideline adherence score.

**Study 4** was a retrospective registry study comparing patients with EMCC stroke suspect criteria codes and patients with hospital stroke diagnoses. The material consisted of AMIS-printouts from Bergen EMCC, and patient data from Haukeland University Hospital, Haraldsplass Diaconal Hospital and Voss Hospital. Less than half of the confirmed stroke patients' initial EMCC contact was via 113 line, 52% of the patients were in contact with the primary health care services first. By combining patient data from the stroke database at Haukeland University Hospital and the 113-data from Bergen EMCC, we calculated sensitivity for stroke identification at EMCC contact of 57.9%, specificity of 99.1%, positive predictive value (PPV) of 45.7% and negative predictive value (NPV) of 99.4%. Stroke patients whom initial EMCC contact was via the 113 line had a higher proportion of acute responses and stroke suspect criteria codes. Although the majority of stroke patients who came in contact with the EMCC via primary health care services did not receive stroke suspect criteria codes, 85% of these calls had free-text notes addressing stroke suspicion.

**Main findings:**

- The national mean 113 contact rate was 56/1 000 a year, but the individual contact rates for the different EMCCs varied between 33 and 114. The urgency distribution was 37% acute, 34% urgent and 27% non-urgent contacts. Index criteria card “06 Unclear problem” was the most frequently used, used in between 10 and 42% of the contacts among the different EMCCs.
- Self-reported use of the Index by the operators was > 75%. Working in rotation, with shifts at both the EMCC and the ambulance service, was associated with reduced use of the Index, while focus on use of the Index at the EMCC increased use. AMIS was the most frequent stated reason for not using the Index.
- Measured mean guideline adherence was 80% of the maximum score. Low guideline adherence score correlated with longer EMD response interval.
- The Index’s ability to identify or predict stroke patients is modest, with a sensitivity of 57.9%, specificity of 99.1%, PPV of 45.7% and NPV of 99.4%. More than half of the stroke patients initially contacted primary health care services instead of 113.

## Norwegian summary – Sammendrag

Hovedmålet med avhandlingen var å sette i gang en valideringsprosess av Norsk indeks for medisinsk nødhjelp. For å kunne gjøre det måtte vi imidlertid først undersøke i hvilken grad Indeks brukes ved norske AMK-sentraler.

Norsk indeks for medisinsk nødhjelp (Indeks) er det beslutningsstøtteverktøyet operatørene ved AMK-sentralen støtter seg til når de mottar og håndterer medisinske nødsamtaler på 113. Indeks ble tatt i bruk i 1994, og foreligger nå i 3de utgave (2009). Samtlige AMK-sentraler og en majoritet av legevaktsentralene i Norge oppgir å bruke Indeks ved mottak av 113 henvendelser. I 2010 var kunnskap om 113 epidemiologien nasjonalt ikke-eksisterende.

Kriteriebaserte beslutningsstøtteverktøy er laget for å brukes i tett samarbeid med helsefaglært AMK-operatør, og operatøren er ment å skulle bruke sin erfaring og kunnskap inn i vurderingsprosessen. Dette har komplisert tidligere validitetsforskning, da det er vanskelig å si om hastegrad og respons utløst skyldes selve verktøyet eller operatørens egen vurdering. I tillegg er Indeks fortsatt papirbasert, slik at bruken ikke kan spores digitalt.

Etter den initiale kartleggingen av sted, situasjon og pasientens vitale funksjoner på Startkortet, er resten av Indeks oppslagene symptombasert. Det gjør det utfordrende å sammenligne spesifikke Indeks kriteriekoder med spesifikke utskrivelsesdiagnoser fra sykehusene, som et mål på hvor god validiteten til Indeks er. Oppslag ”27 Nedsatt bevissthet – lammelser”, nærmere bestemt kriteriekode A.27.03 – A.27.06 ble vurdert å kunne måles opp mot diagnosen hjerneslag, inkludert hjerneblødning, hjerneinfarkt og drypp.

Målet med de enkelte studiene var å få mer kunnskap om:

- 113 epidemiologien i Norge
- bruken av Indeks ved AMK
- Indeks' validitet med tanke på å identifisere og predikere hjerneslag

**Studie 1** var en tverrsnittsundersøkelse av alle 113-henvendelser til samtlige 19 AMK-sentraler gjennom en 72 timers periode i august 2011. Det endelige materialet bestod av 2 298 utskrifter fra det elektroniske journalsystemet ved AMK, Akuttmedisinsk Informasjonssystem (AMIS), og inneholdt informasjon om tid, innringer, pasient, Indeks kriteriekode, hastegrad og involverte ressurser. Samlet 113 kontaktrate var 56 /1 000 innbyggere per år, men variasjonen for de ulike AMK-sentralene var fra 33 til 114. Den akutte kontaktraten var 21/1 000 per år, med variasjon 5 til 31. Hastegradsfordelingen viste 37% akutte, 34% haster og 27% vanlige henvendelser. Det hyppigst brukte oppslaget var "6 – Uavklart problem", som ble brukt i 20% av henvendelsene, med en variasjon på 10 til 42% mellom AMK-sentralene.

**Studie 2** var en nasjonal spørreskjema studie til operatørene ved samtlige 19 AMK-sentraler, for å kartlegge selvrappert bruk av Indeks. Spørreskjemaet inneholdt spørsmål om bruk av Indeks, utdanning, AMK-erfaring, og opplæring i og repetisjon av bruk av Indeks ved sentralen. Svarprosenten var 63.4%, og den typiske operatøren var en kvinnelig sykepleier med seks års AMK-erfaring, og jobbet i rotasjon med enten akuttmottak eller ambulanse. Bruk av Indeks ble kalkulert til å være 3.95, svarende til responsformatet 4 = "ofte, > 75%". Det å jobbe både ved AMK og i ambulansetjenesten reduserte bruk av Indeks, mens opplevd fokus på bruk av Indeks ved arbeidsplassen økte bruken. Operatørene oppga å undersøke om pasienten var bevisst i 93% av henvendelsene, men brukte Startkortet i kun 47%. Bruk av AMIS var hyppigste oppgitte årsak til ikke å bruke Startkortet eller Indeks som helhet.

**Studie 3** var en strukturert gjennomgang av lydlogger fra samme tidsperiode som studie 1, for å måle bruk av Indeks objektivt, og deretter se om bruk av Indeks påvirket AMK-reaksjonstid. Syv strategisk utvalgte AMK-sentraler deltok, med tilsammen 299

randomiserte akutte og hastehenvendelser, 174 akutte og 125 haster. Utvelgelsen av sentraler var basert på spredning i størrelse, geografisk lokalisasjon, ulike helseforetak, og sentralenes gjennomsnittlige selvrapperte bruk av Indeks fra studie 2. Det ble utarbeidet et lytteskjema som målte om og hvor raskt operatøren fikk bekreftet ulike indikatorer på bruk av Indeks. Indikatorene lokalisasjon, bevissthet og strukturert gjennomgangen av kriterier på valgt oppslag ble satt sammen til en variabel, der 0 var minsteskår og 3 maksskår. Gjennomsnittlig bruk av Indeks var 2.41, svarende til 80% av maksskår. AMK-reaksjonstid økte med over 2 minutter for de henvendelsene med minsteskår på bruk av Indeks, sammenlignet med de med maksskår.

**Studie 4** var en retrospektiv registerstudie som sammenlignet pasienter med slagsuspekterte kriteriekoder fra AMK med pasienter med hjerneslagsdiagnoser fra sykehus. Materialet bestod av AMIS-utskrifter fra Bergen AMK, og pasientdata fra Haukeland Universitetssykehus, Haraldsplass diakonale sykehus og Voss sykehus. Mindre enn halvparten av de bekreftede slagpasientene kom i kontakt med AMK via 113, 52% var i kontakt med primærhelsetjenesten først. Ved å kombinere pasientdataene fra slagdatabasen ved Haukeland Universitetssykehus med 113-data fra Bergen AMK kalkulerte vi sensitivitet for å oppdage slagpasienter ved initial kontakt med AMK på 57.9%, spesifisitet på 99.1%, positiv prediktiv verdi (PPV) på 45.7% og negativ prediktiv verdi (NPV) på 99.4%. Slagpasientene som kom i kontakt med AMK via 113 hadde høyere andel akutte hastegrader og slagsuspekterte kriteriekoder enn de som først var i kontakt med primærhelsetjenesten. Selv om majoriteten av slagpasientene som gikk via primærhelsetjenesten ikke fikk slagsuspekterte kriteriekoder, hadde 85% av disse et fritekstnotat om mistanke om slag.

**Hovedfunn:**

- Den nasjonale kontaktrate for 113 var 56/1 000 per år, men varierte fra 33 til 114 mellom de ulike AMK-sentralene. Hastegradene fordelte seg med 37% akutte, 34% haster og 27% vanlige henvendelser. Det hyppigst brukte oppslaget var ”6 Uavklart problem”, som ble brukt i mellom 10 og 42% av henvendelsene ved de ulike AMK-sentralene.
- Selvrappert bruk av Indeks blant operatørene var > 75%. Jobbrotasjon mellom AMK-sentral og ambulanse var assosiert med redusert bruk av Indeks, mens opplevd fokus på bruk av Indeks ved sentralen økte bruken. AMIS var den hyppigst oppgitte årsaken for ikke å bruke Indeks.
- Målt bruk av Indeks var 80% av maksimalskåren. Det var en sammenheng mellom lav bruk av Indeks og lengre AMK-reaksjonstid.
- Indeks’ evne til å identifisere eller predikere slagpasienter var beskjedne, med sensitivitet på 57.9%, spesifisitet på 99.1%, PPV på 45.7% og NPV på 99.4% . Over halvparten av slagpasientene kontakter primærhelsetjenesten i stedet for 113 direkte.



## List of publications

The thesis is based on three papers, from four studies:

### Study 1 and 2

**Paper I: Variations in contact patterns and dispatch guideline adherence between Norwegian emergency medical communication centres – a cross-sectional study.**

Ellensen EN, Hunskaar S, Wisborg T, Zakariassen E

*Scand J Trauma Resusc Emerg Med.* 2014;22:2.

doi: [10.1186/1757-7241-22-2](https://doi.org/10.1186/1757-7241-22-2)

### Study 3

**Paper II: Dispatch guideline adherence and response interval – a study of emergency medical calls in Norway.**

Ellensen EN, Wisborg T, Hunskaar S, Zakariassen E

*BMC Emerg Med* 2016;16:40. doi: [10.1186/s12873-016-0105-2](https://doi.org/10.1186/s12873-016-0105-2)

### Study 4

**Paper III: Stroke identification by criteria-based dispatch.**

Ellensen EN, Naess H, Wisborg T, Hunskaar S, Zakariassen E

*Submitted December 2016*

The studies are referred to as **Studies 1-4**, and the papers as **Papers I-III** throughout the thesis. The papers are included in full text in the appendices. Reprints were made with permissions from Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine and BMC Emergency Medicine.

## Contents

<i>Scientific environment</i>	2
<i>Acknowledgements</i>	3
<i>Preface</i>	4
<i>Abbreviations and terminology</i>	5
<i>Summary</i>	10
<i>Norwegian summary – Sammendrag</i>	14
<i>List of publications</i>	18
<i>Table of contents</i>	19
<b>1 Background</b>	<b>22</b>
1.1 Prehospital emergency medical services	22
1.2 Prehospital emergency medicine in Norway	23
1.2.1 Emergency primary health care system	23
1.2.2 Emergency secondary health care system	24
1.3 The Emergency Medical Communication System	25
1.4 Emergency Medical Communication Centres	26
1.4.1 Acute Medical Information System	27
1.4.2 Audio log recordings	27
1.5 Emergency medical dispatch	27
1.6 The Norwegian Index for Emergency Medical Assistance	29
1.6.1 Start page	30
1.6.2 Criteria cards	30
1.6.3 Urgency and responses	31
1.7 Criteria card “27 Altered levels of consciousness – paralysis” and stroke	32
1.8 Summary of background	33
<b>2 Aim and objectives</b>	<b>34</b>
<b>3 Methods and material</b>	<b>35</b>
3.1 <b>Study 1 – 113 epidemiology (Paper I)</b>	35
3.1.1 Method and data collection	36
3.1.2 AMIS printouts	36
3.2 <b>Study 2 – Self-reported use of Index (Paper I)</b>	36

3.2.1	Method and data collection	36
3.2.2	Questionnaire	37
3.3	<b>Study 3 – Guideline adherence (Paper II)</b>	37
3.3.1	Study sample	38
3.3.2	Randomization of calls	38
3.2.3	Data collection	39
3.2.4	Listening form	39
3.2.5	Variables and outcome measures	40
3.4	<b>Study 4 – Stroke identification (Paper III)</b>	40
3.4.1	Method	40
3.4.2	Data collection	41
3.4.3	Material	42
3.5	Statistical analyses	44
3.6	Ethical considerations	44
<b>4</b>	<b>Main findings</b>	<b>46</b>
4.1	<b>Study 1 – 113 epidemiology</b>	46
4.2	<b>Study 2 – Self-reported use of the Index</b>	47
4.3	<b>Study 3 – Guideline adherence</b>	49
4.4	<b>Study 4 – Stroke identification</b>	50
4.4.1	Stroke identification	51
4.4.2	Stroke prediction	53
<b>5</b>	<b>Discussion</b>	<b>54</b>
5.1	Summary of main findings	54
5.2	Methodological considerations	54
5.2.1	Prehospital EMS research	55
5.2.2	AMIS as data source	55
5.2.3	Questionnaire and listening form	56
5.2.4	Strategical selection of EMCCs	57
5.2.5	Criteria cards	57
5.2.6	Missing data	58
5.3	Discussion of the results	59

5.3.1	Use of the Index	59
5.3.2	Stroke identification	64
<b>6</b>	<b>Conclusions</b>	<b>68</b>
<b>7</b>	<b>Future aspects</b>	<b>69</b>
7.1	Implications of the findings	69
7.2	Future research	70
	<i>References</i>	71
	<i>Appendices:</i>	79
1	Index: Table of contents	
2	Index: Start page	
3	Index: Criteria card 27	
4	Questionnaire	
5	Listening form	
<b>6</b>	<b>Papers I-III:</b>	
I	Variations in use of emergency medicine dispatch guidelines in Norway – a cross-sectional study	
II	Dispatch guideline adherence and response interval – a study of emergency medical calls in Norway	
III	Stroke identification by criteria-based dispatch	

# 1 Background

## 1.1 Prehospital emergency medical services

Emergency medical service (EMS) has been defined as a system which provides effective, coordinated and timely delivery of health care services to victims of sudden illness or injuries (1). Historically, prehospital EMS strategies have been described as dichotomous; basic level service versus advanced level service, or in other words “scoop and run” versus “stay and stabilize”. The “scoop and run” strategy aimed to bring the patient to the hospital as fast as possible, and was used to a great extent in EMS systems with open access to the emergency ward. The ambulance services were typically ground-based, and considered part of the public safety system, alongside fire and police departments. The “stay and stabilize” strategy aimed to bring the hospital to the patient, and utilized physician or paramedic-staffed vehicles, both cars and helicopters, with more advanced equipment, alongside regular ambulances (1-3).

The organization of EMS systems varies substantially throughout the world, as summarized in a series on EMS organization worldwide published in *Resuscitation* 2003–2009, utilizing a diversity of combinations of ambulance workers, paramedics, specialized nurses, general practitioners, anaesthesiologists, emergency physicians and physicians in residency (2-12). More recent publications on EMS organization are sparse, but there are reasons to believe that most EMSs today use a combination of both the “scoop and run” and the “stay and stabilize” strategy, depending on acuity, patient need, location and resources available.

This heterogeneity in EMS systems, together with lack of uniform data collections, often results in small and non-generalizable studies where the isolated outcome effect of the prehospital interventions can be difficult to measure (13).

A qualitative study published in 2014 found multiple key barriers for conducting prehospital clinical studies: funding, ethical approval, data collection, protocol training and compliance, randomizing and blinding interventions, obtaining patient outcomes,

adequate study staffing and EMS partnering (14). A meta-analysis of randomized trials published in a high impact factor emergency medicine journal revealed that only 8% of the studies were from the prehospital field (15). Overall, prehospital emergency medicine has been a neglected area of research for many years compared to in-hospital medical fields (13). A PubMed search reveal that although there is a 7.4 times increase in number of publications involving “prehospital” and “emergency medicine” from 1996–2000 to 2011–2015, the number of medical publications as a whole rises by 9.2 during the same period (16).

## **1.2 Prehospital emergency medicine in Norway**

The health care system in Norway is twofold, with a primary care level and a specialist/hospital care level. A general referral system serves as a strict gatekeeper towards the second level: the hospital emergency department can only be attended by patients admitted by physicians, or acute patients brought in directly by ambulances.

Although the majority of the 5 million population lives in urban areas, there are rural settlements along the entire coastline of 25 000 km. Scattered population, fjords, mountains, seasonal climate changes and poor road quality many places provide challenging conditions for the rescue and transport of patients (7).

### *1.2.1 Emergency primary health care system*

The principle that all issues should be dealt with at the lowest effective level of health care means that the large quantity of patients in need of non-urgent or urgent medical assistance are handled by the emergency primary health care system: GPs and the out-of-hours (OOH) service. The regular GPs are obligated to provide emergency medical assistance for their own patients during office hours. The out-of-hours medical assistance is provided by LEMCs and casualty clinics, manned by nurses and

physicians. In addition to regular GPs taking out-of-hours shifts on call, physicians in internship, hospital physicians and foreign stand-in physicians have played an important role in manning the casualty clinics, accounting for almost half of the shifts (17). In 2015, a new regulation on emergency medicine stated that the physician on call at the casualty clinics should be a GP (18). Besides handling urgent and non-urgent patients at the casualty clinics, the GPs on call are also supposed to be notified about all acute patients prehospital but choose themselves whether to attend the situation or not. Both notification and involvement of GPs with acute patients have proved to vary geographically (19,20).

The municipalities are responsible for the primary health care system, including the medical emergency parts. The number of casualty clinics is decreasing, indicating increasing inter-municipal cooperation. In 2014, 58% of the casualty clinics were inter-municipal and 42% municipal (21). The emergency primary health care system handled close to 1 950 000 contacts in 2015 (17).

### *1.2.2 Emergency secondary health care system*

The patients in need of acute medical assistance are mainly handled by the secondary health services: the ambulance service and EMCCs (7,22), but GPs on call are supposed to be alarmed and assist if necessary. The ambulance service include cars, boats, planes and helicopters. Cars are staffed by two ambulance workers, planes by specialized nurses and helicopters by anaesthesiologists and rescuers (18). Each helicopter base also has an emergency car for missions when use of helicopter is inappropriate: close proximity to the base or poor weather conditions.

In contrast to GPs being notified of all acute patients, the criteria for notifying helicopter emergency medical services (HEMS) are rather unspecific: “If appropriate, consider sending the nearest otherwise occupied ambulance; notifying nearest doctor not on call; notifying HEMS; recommend private patient transport.” Geographical

access, shortened transfer time and advanced prehospital critical care interventions have been found to be the three main benefits of primary prehospital HEMS (23-25), and HEMS availability has been used as a factor mitigating centralization of trauma care and specialized health services. This is based on the assumption that the EMCCs are able to dispatch HEMS to the patients who need it, and avoid unnecessary use for patients with minor injuries or no other indication for HEMS. Once notified by EMCC, the HEMS anaesthesiologist accepts or declines the mission based on medical indication (26,27), and the EMCC thus have no authority to dispatch the helicopter, in contrast to the rest of the ambulance fleet. In Western Norway more than one third of the HEMS dispatches from the EMCC were declined or aborted by the HEMS crew, due to diminished medical indication, bad weather conditions or competing missions (27).

The regional trusts are responsible for the secondary health care system, and all contacts are made through the EMCCs. In 2015 the EMCCs handled close to 850 000 contacts (26).

### **1.3 The emergency medical communication system**

When in need of acute medical assistance, the public has access to both primary and secondary health care level through the two toll-free emergency telephone numbers: The 116 117 line, to be used in an urgent medical emergency situation; and the 113 line when there is an acute, life-threatening situation.

The 116 117 line is answered by the nearest LEMC. They assist the caller with medical advice and referral to GPs and casualty clinics. If necessary, they can also request an ambulance through the EMCC. During the last decade, there has been a desired shift towards increased use of LEMC as the public's access point during out-of-hours, but some casualty clinics still allow direct attendance without calling beforehand. This is especially common in the larger cities.



The 113 line is answered by the nearest EMCC, which also coordinates and dispatches the ambulance fleet when needed. If deemed appropriate, the EMCC can transfer an obviously non-urgent call to LEMC, and vice versa.

In addition to assisting the public in need of acute medical help, the emergency medical communication system (113, EMCC and LEMC) also facilitates radio and telephone communication between all the different participants in the chain of pre-hospital emergency medical care; ambulance personnel, GPs on-call, casualty clinics and in-hospital specialists (18,28). From 2015, the emergency health services, police and fire and rescue services have used Nødnett, a separate radio network for inter-agency public safety communication during rescue missions, accidents and major situations (29).

#### **1.4 The Emergency Medical Communication Centres**

The EMCCs and the ambulance service are part of the secondary health care system, administered through four regional health trusts. There are now 16 EMCCs in Norway (2017); 4 in North, 4 in Middle, 4 in West, and 5 in South-East (30). They differ in size, with regard to the corresponding population (65 000–1 165 000 in 2011) (31), tasks and staffing. The EMCCs are normally located at the nearest hospital, although co-localization with the other emergency services, fire brigade and police, also exists. In addition to the public 113 line, the centres also receive direct lines from casualty clinics, in-hospital emergency lines, ambulance booking line, and admission lines from GPs. Some EMCCs are co-located with the nearby LEMC.

Although the personnel requirements of the regulations have been unspecified, most EMCCs are manned by registered nurses and ambulance workers with specific EMCC training (18). The EMCC operators, usually nurses, traditionally answer and triage phone calls made to the 113 line, and assist the caller with medical advice if necessary. The resource coordinators, usually ambulance workers, handle health lines and

dispatch and coordinate the entire ambulance fleet including cars, boats, planes and helicopters. At the smaller EMCCs these tasks can be handled by the same person.

National EMCC data has been available to some extent from 2012 (26,30,32,33).

#### *1.4.1 Acute Medical Information System*

The software program Acute Medical Information System (AMIS) is used by all EMCCs to register and document each contact. AMIS records contain information on patient identity, location, caller, time of all events, Index criteria, urgency level, response dispatched, and resources involved. It is a local record system for the EMCC alone, and not integrated with the hospitals' record systems.

#### *1.4.2 Audio log recordings*

Prior to the regulation of 2015 (18) there was no time limit for storage of audio log recordings. After the new regulation, all calls are taped and stored for a minimum of three years, then they are deleted. There is an exemption for deletion if the log has served as evidence or documentation in a complaint, case supervision or other case management.

### **1.5 Emergency medical dispatch**

Emergency medical dispatch is an important key in the prehospital chain of survival. Identification of patients in potential life-threatening situations is the vital first step toward providing them rapid and appropriate help. In the 1980s two different dispatch systems developed in the United States of America; the Medical Priority Dispatch System (MPDS) and the Criteria Based Dispatch (CBD) system. The MPDS is an

algorithm-based protocol, digitalized and used by non-health educated personnel with special MPDS training. The system has the advantage of easy monitoring on protocol compliance and other quality markers (34). This facilitates research in general and in particular validity studies (35).

The CBD guidelines use symptoms and prompts to determine the nature of the medical condition, the urgency and hence the appropriate level of response. It is supposed to be used by an experienced health-educated operator with special EMCC training. This allows for a dynamic approach to the situation at hand, but also complicates research on CBD (34,36). The complex interaction between the supporting guidelines and the operator introduces an unknown variance when studying guideline adherence and validity (37,38). The skill, knowledge and experience of the individual operator will interfere with how the guidelines are interpreted and modified to the situation at hand. This will influence the criteria set, and hence the urgency and response dispatched. Another factor complicating CBD research is the symptom-based approach, which makes comparison between initial dispatch criteria and final hospital discharge diagnoses as a measure of validity difficult.

A 2013 systematic review on adherence to prehospital guidelines and protocols failed to identify any eligible study in the field of emergency medical dispatch, neither CPD, nor MPDS-based (39). Nevertheless, a Norwegian study from 2005, evaluating dispatch in drug-related emergencies, found an operator guideline compliance of 99%, based on a multiple choice questionnaire, and a 64% compliance when going through log recordings (40). Research on CBD compliance internationally has been sparse, but in 2014 a study on cardiac arrest calls compared an American MPDS protocol to the Norwegian CBD guidelines (Index) and found both to have equally high levels of consciousness clarification (100% for MPDS and 97% for CBD) and detection of respiratory distress (100% and 98%, respectively) (41).



The flip-over Index in use by an operator at the EMCC (Photo: VS Ellensen).

## 1.6 The Norwegian Index for Emergency Medical Assistance

The Index comprises the criteria-based medical dispatch guidelines used in Norway (28). It was introduced in 1994, developed from the original American CBD guidelines (42). As the EMCC holds a key role in resource allocation in the prehospital setting, implementation of the Index at all EMCCs should secure a comprehensive and equitable service for all citizens regardless of location. In addition to being used by all the EMCCs when receiving medical emergency phone calls (113), it is also used by a majority of the LEMCs when receiving out-of-hours phone calls (116 117) (21).

It is designed to be used by health educated operators with special training, and aims to assist the operator throughout the dispatch process; initial survey of the situation at hand, dispatch of the adequate resources, and finally, provision of medical advice and instructions on first aid and care for the caller. Despite its many years in service, and

several exports abroad, Denmark and Croatia most recently (43,44), there was no knowledge in 2010 of how and to which degree it was used or how well it performed.

In 2016, the Index still exists as a flip-over in large paper format in Norway. It consists of a start page and 40 criteria cards (Appendix 1).

### *1.6.1 Start page*

The start page is algorithmic and clarifies vital information like location, phone number, vital functions of the patient and a brief questioning on the problem at hand (Appendix 2). The operator then decides if there is an immediate need of an ambulance, life-saving instructions, or if the time allows; moving on to the proper criteria card to explore the situation further.

### *1.6.2 Criteria cards*

The first three criteria cards are mainly instructional (“01 Unconscious [lifeless] adult”, “02 Unconscious [lifeless] child” and “03 Choking / foreign body in throat”), with a focus on cardiopulmonary resuscitation. There are five cards of a more administrative order (“04 Disaster – major accident”, “05 Transport reservations”, “24 Help in crisis”, “31 Cooperation with others” and “40 About Index”) and 32 cards based on symptoms and situations.

A symptom card is a list of criteria prompts in decreasing severity, and the operators start at the top and move downwards until one criterion is met. Each criterion has a predefined urgency and a recommended dispatch. The cards also include additional questions, caller advice and supporting instructions for health personnel on scene. “27 Altered levels of consciousness – paralysis” is enclosed as a criteria card example in appendix 3.

The administrative criteria card “05 Transport reservations” is used for a variety of ambulance transports, including primary responses, secondary transfers and return of patients. It includes all urgencies, and is used for both pre-planned and acute incidents. If the caller is a physician or another health personnel calling in to order an ambulance, the operator can choose between using criteria card 5 or using a more specific symptom card according to the patient at hand.

Criteria card “06 Unclear problem” is a collection of criteria covering mainly unclear situations, where the operator is unable to provide enough information to designate a proper criteria card. It also contains some more specific criteria that don’t fit into any other criteria card.

### *1.6.3 Urgency and responses*

The Index divides the responses into three urgency categories: acute (red), urgent (yellow) and non-urgent (green). Acute responses are when the situation is acute and life-threatening, and there is a need for immediate acute ambulance dispatch and alarm of the GP on-call. Urgent responses are potentially life-threatening situations where an ambulance is normally dispatched but without lights and sirens, and the need for the GP on-call is assessed. In non-urgent responses, there is no life-threatening situation and the patients are usually referred to the primary level of emergency care, the casualty clinics, unless there is an obvious need for ambulance transport.

## **1.7 Criteria card “27 Altered levels of consciousness – paralysis and stroke”**

The criteria card “27 Altered levels of consciousness – paralysis” is one of very few Index cards corresponding closely to a medical diagnosis, or diagnoses group, namely stroke (Appendix 3). Over the past decade, stroke has entered prehospital emergency medicine as a time-critical event in need of acute response, on the same acute level as heart attack and severe trauma. Current stroke treatment gives better outcomes, both in terms of survival and dependency, but the therapeutic window is time-limited. The benefits decrease proportionally with increasing time between symptom onset and treatment, disappearing between 4.5 and 6 hours (45,46).

Time from symptom onset to treatment relies on many factors: patient’s or bystander’s awareness of stroke symptoms and urgency for seeking medical assistance; dispatch recognition of stroke symptoms; prehospital time, including time to scene, on-scene and transport to hospital; and in-hospital time, including diagnostics (47,48). Identification of stroke at EMCC can reduce both prehospital and in-hospital time, through acute responses and pre-arrival notification of the hospital (49-51).

Although beneficial for the acute ischemic stroke (AIS), thrombolysis is potentially lethal to patients suffering from haemorrhagic stroke. Imaging diagnostics is therefore essential prior to treatment. Mobile stroke units, ambulances equipped with computed tomography (CT) scanners, are now emerging in the prehospital field (52-55), aiming to reduce time to treatment by eliminating in-hospital time spent on diagnostics. In the future, thrombolysis might be given prehospital as well, reducing the time from symptom onset to treatment substantially. But, this presupposes a valid stroke dispatch.

Studies on different dispatch protocols have found a sensitivity for stroke identification ranging from 41% to 83% (56-60). The ability of Index and criteria card “27 Altered levels of consciousness – paralysis” to identify stroke patients is unknown.

## **1.8 Summary of background**

Internationally the organization of emergency medical systems is diverse, both within countries, and between countries. In Norway, the emergency medical health care system is twofold: The emergency primary health care system includes the GPs, the casualty clinics and the LEMCs, with the public urgent emergency medical telephone line 116 117. The secondary level includes the ambulance service and the EMCCs with the public acute emergency medical telephone line 113.

The EMCCs are manned by operators handling the 113 calls, and resource coordinators dispatching and administering the ambulance fleet. Since 1994, all Norwegian EMCCs use the same CBD guidelines, the Index, but both validity of the guidelines and the actual use of it remained unknown.

Emergency medical dispatch is a key element in the chain of prehospital emergency medical care. The nature of CBD calls for close interaction between the operator and the guidelines, and prior CBD research has been sparse due to the difficulties of addressing these tangled factors. The criteria cards are symptom-based, which challenges the matching of initial dispatch criteria code and patient outcome. Criteria card “27 Altered levels of consciousness – paralysis” and stroke diagnosis are considered to be the closest match possible. Stroke is a highly time critical situation, as the treatment is time-dependent. Valid dispatch of the timely and appropriate resources is a presumption for proper prehospital resource allocation.



## **2 Aim and objectives**

The overall aim of this thesis is to contribute to increased knowledge and awareness about how and to which degree the Index is used, and to test its ability to identify stroke patients.

The four studies had the following specific objectives:

1. To investigate the epidemiology of 113 calls, focusing on possible differences among the EMCCs in urgency levels, Index criteria use and contact rates.
2. To determine self-reported use of the Index among operators, and explore possible factors influencing this use.
3. To measure guideline adherence in an objective manner, and explore a possible effect on response interval.
4. To explore sensitivity, specificity and predictive values of stroke identification at initial EMCC contact, and to evaluate possible factors associated with stroke identification and prediction.

### 3 Methods and material

AMIS records from the EMCCs were collected as a data basis for all four studies. The operators' self-reported use of the Index and measured guideline adherence were obtained through questionnaires and audio log recordings. In the last study, we compared AMIS data on Index criteria codes at dispatch with hospital data on diagnoses at submission.

The studies were performed 2011–2013. The first two studies were national, including all EMCCs in Norway. The third study included a strategic selection of EMCCs (7 out of 19), and the last study focused on one EMCC in particular.

During the study period, there were 19 EMCCs in Norway; 5 in the North, 4 in the Middle, 4 in the West and 6 in the South-East health trusts. They differed in size, with populations ranging from 65 000 to 1 165 000, calculated by the end of 2011 (31). All EMCCs officially used the Index as their only tool to prioritize and handle 113 calls, and the software program AMIS to register and document each contact.

We conducted a total of four studies (**Studies 1-4**), contributing to a total of three papers (**Papers I-III**). Findings from the first study were mainly reported in **Paper I**, but the material was also used in **Paper II**.

#### 3.1 Study 1 – 113 epidemiology (Paper I)

This was a national cross-sectional population-based epidemiologic study on 113 calls. The study period was 72 hours during late August 2011, including both weekdays and weekend. The purpose of the study was to investigate the epidemiology of 113 calls among the different EMCCs, with regards to urgency levels, Index criteria use and population contact rates.

### *3.1.1 Method and data collection*

All 19 EMCCs contributed with AMIS data for every medical emergency contact made to 113 during the study period, 2 298 in total. Calls from other direct lines (fire department, police and LEMCs), incidents with too little information (missing three or more of the following: gender, age, Index criteria, urgency or response), duplicates, maculated incidents and misdialling were excluded.

### *3.1.2 AMIS printouts*

The AMIS records were collected as printouts. They contained information on date, time, caller role, patient gender and age, set Index criteria, set urgency, response, and involvement of resources like ambulances, air ambulances, GPs and others.

## **3.2 Study 2 – Self-reported use of the Index (Paper I)**

This was a national questionnaire-based survey on the use of the Index and associated factors. All 19 EMCCs participated. The purpose of the study was to determine use of the Index as perceived by the operators themselves, and explore possible factors influencing their use.

### *3.2.1 Method and data collection*

A questionnaire on the use of the Index was sent to all 429 EMCC operators in Norway, distributed through EMCC management and returned directly to the research facility to secure anonymity for the individual respondent. Operators were defined as personnel whose primary occupation was to answer the 113 calls. Resource coordinators were defined as personnel whose primary occupation was to dispatch

resources and coordinate the ambulance fleet. Only answers from operators were included in the study.

### 3.2.2 *Questionnaire*

The questionnaire was developed in cooperation with Bergen EMCC and tested on their operators to avoid misinterpretations. It included different sections on individual use of the Index, education, EMCC work experience, and initial training and repetitions on use of the Index. It is enclosed in Appendix 4.

The eleven questions on the operators use of the Index were arranged as a Likert scale of five symmetric response formats: never, seldom (<25%), sometimes, often (>75%) and always. Each question was expressed in the same form: “During a real emergency call, how often do you...?” and covered different parts of the Index. In earlier editions of the Index, the question “Does the patient breathe?” was part of the start page. This was changed to “Is the patient able to talk?” in the 2009 3<sup>rd</sup> edition (28). This switch was used in the study to reveal whether the operator actually used the present start page, or if they used their own mental version of the former vital questions, by picking questions directly from the start page, like “... Check if the patient is awake,” and “... Is the patient able to talk?” It was possible to add explanations or points of view at the end of the questionnaire.

## 3.3 **Study 3 – Guideline adherence (Paper II)**

This was an observational cross-sectional study of acute and urgent 113 calls during the same 72-hour study period utilized in **Study 1**. The purpose of the study was to measure guideline adherence by listening through audio log recordings of real emergency calls, and to explore whether guideline adherence affects emergency medical dispatch (EMD) response interval.

### 3.3.1 Study sample

A strategic selection of nine EMCCs was invited to participate, and seven accepted. The selection was based on diversity in geographic area (all health regions represented), EMCC size, population density (5–210 inhabitants per km<sup>2</sup>), and findings from the first paper; contact rates (36–75 contacts per 1 000 inhabitants a year) and self-reported use of the Index at EMCC level (3.3–4.4). The EMCCs constituting the study sample were: Tromsø, Harstad, Trondheim, Bergen, Sørlandet, Buskerud and Oslo og Akershus.

In order to verify the sample validity, contact rates and mean self-reported use of the Index in the EMCC study sample were compared to the national findings reported in **Paper I** (61). The study sample had a contact rate of 57/1 000 inhabitants a year, corresponding to the national rate of 56/1 000 inhabitants a year. The mean self-reported use of the Index in the sample was slightly below the national findings, 3.86 (*SD* 0.40) and 3.95 (*SD* 0.39) respectively, but both corresponded to the response format “often, > 75%” (=4).

### 3.3.2 Randomization of calls

A random sample of 300 acute and urgent calls from the study period were included in the study. The five largest EMCCs contributed 50 calls each, while the two smallest EMCCs in the study sample, Tromsø and Harstad, did not have 50 acute or urgent calls during the study period, and hence contributed 30 and 20 calls instead. Guideline adherence during the call was evaluated using the developed listening form. The randomization of the calls was achieved through a number generator ([www.random.org](http://www.random.org)). Corresponding AMIS records for each call provided information on event times, set criteria and set urgency. One call was excluded due to missing AMIS data, resulting in 299 included calls.

### 3.3.3 Data collection

The call listening and evaluation sessions on Index adherence were done locally at each EMCC by the primary researcher personally.

### 3.3.4 *Listening form*

The listening form used in the study was based on an internal quality control form used by Bergen EMCC, but modified to fit our study. The form is enclosed in Appendix 5. Both information given spontaneously by the caller and information specifically asked for by the operator were registered separately, but later merged for analyses.

Each form measured six different indicators. Indicators 1–4 were derived from the start page and measured vital information: (1) confirmation of caller phone number, (2) location, (3) patient consciousness, (4) patient responsiveness. These were categorized into “within 1 minute”, “within 2 minutes” or “not at all”. Information gained after 2 minutes was interpreted as negative.

Indicator 5, criteria compliance, measured the number of criteria above the set criterion that were not accounted for during the call, either spontaneously or asked for by the operator. Hence an increasing criteria compliance value means the operator has overlooked some more urgent or vital criteria. The indicator was grouped in 0, 1 and 2 and more.

Indicator 6, communication problems, was registered positive if the operator failed to understand what the caller was trying to communicate, as assessed by the researcher.

### 3.3.5 Variables and outcome measures

The three variables confirmation of location, confirmation of consciousness and criteria compliance were combined to compose the main outcome variable “Overall guideline adherence”. Each variable was dichotomized, and information gained within 1 minute for the two first variables, and zero or one unchecked criteria above the set criterion were interpreted as positive. Information gained after 2 minutes and 2 and more criteria above set unchecked were both interpreted as negative. All three variables were weighted equally, composing a measured guideline adherence score of 0 (no positive) to 3 (all positive).

## 3.4 Study 4 – Stroke identification (Paper III)

This was a retrospective register-based study. The data period was 1st January 2011–31st January 2012. The purpose of the study was to explore the Index’s ability to identify and predict stroke at initial EMCC contact, and to explore possible factors associated with stroke identification and prediction.

### 3.4.1 Method

The data collection for this study was stepwise:

First step:

- a. Identify all criteria code 27 patients from Bergen EMCC.
- b. Identify all confirmed stroke patients at Haukeland University Hospital

Second step:

- a. Code 27 patients hospitalized at Haraldsplass Diaconal Hospital and Voss Hospital – stroke diagnoses or not?
- b. Confirmed stroke patients with no code 27 – which EMCC code?

A stroke suspect was defined as a patient receiving dispatch code A.27.03 – Sudden facial drooping, A.27.04 – Sudden impaired strength in an arm or a foot, A.27.05 – Sudden speech difficulties or A.27.06 – Increasing confusion/bluntness – suspicious for stroke at initial EMCC contact. These codes were chosen because they are marked in Index as stroke-suspicious, asking the operator to consider actions facilitating thrombolysis. Criteria card 27 is enclosed in the appendices (App 4).

A stroke patient was defined as a patient with a diagnosis of intracerebral haemorrhage (I61), acute ischemic stroke (I63), unspecified stroke (I64) or transient ischemic attack (G45) at hospital discharge, according to International Classification of Diseases, Tenth Revision (ICD-10).

### *3.4.2 Data collection*

AMIS records were collected from Bergen EMCC; data on stroke patients were retrieved from the Bergen study part of the Norwegian stroke registry (NORSTROKE) at Haukeland University Hospital. Supplementary diagnostic data on stroke suspects admitted elsewhere were retrieved from Haraldsplass Diaconal Hospital and Voss Hospital.


Cases were merged based on the unique civil registration numbers and date of EMCC contact/hospital admission.




### 3.4.3 Material

During the study period, there were 549 patients with confirmed stroke at Haukeland University Hospital. In the same period, there were 94 606 medical calls to Bergen EMCC, and 1 013 patients received a 27 criteria dispatch code. Of these, there were 709 patients with a stroke-suspect dispatch code of A.27.03–06, and 568 of these were hospitalized. In Step two, we collected dispatch codes for the stroke patients without a code 27, and diagnoses on the code 27 patients sent to Haraldsplass Diaconal Hospital or Voss Hospital. The material is illustrated in Figure 3.4.3 a and b.

**Figure 3.4.3a** Flow chart of stroke patients at Haukeland University Hospital and their initial EMCC contact line.



**Figure 3.4.3b** Flow chart of stroke suspects after initial contact with EMCC in the study period.



### 3.5 Statistical analyses

Results are presented in absolute numbers ( $N$ ), percentages (%) and odds ratios ( $OR$ ). Continuous data are presented as means (standard deviation) for symmetric data, and medians (interquartile range) for skewed data. Rates were calculated as contacts per 1 000 inhabitants per year. Response interval was analysed as a continuous variable (seconds). Standard error of the observed proportion was calculated using the normal approximation to the binominal distribution, providing 95%  $CI$  for sensitivity, specificity and predictive values (**Study 4**). Outcome variables were checked for normal distribution by Q-Q plots.

The Likert scale in **Study 2** was constituted from eleven questions and five response formats valued 1 to 5. The outcome variable “use of Index” was calculated for each operator as the mean Likert scale score. One-way analysis of variance (ANOVA) was used to compare the EMCCs with regards to variation in mean use of Index (**Study 2**) and guideline adherence (**Study 3**), while Mann –Whitney U test and Kruskal-Wallis tests were used to compare scores on response interval (**Study 3**). Different regression analyses were used to explore factors associated with use of Index (**Study 2**, univariate and multivariate linear), and with identification of Stroke (**Study 4**, binary logistic).

Statistical analyses were performed using Statistical Package for the Social Sciences (IBM SPSS version 20 for **Studies 1** and **2**, and version 23 for **Studies 3** and **4**) and STATA (Stata/IC 12.1 for **Studies 1** and **2**).  $P$ -values below 0.05 were considered statistically significant.

### 3.6 Ethical considerations

Asking patients and next of kin for consent can be difficult. Emergency medical situations are often particularly complex, as it may include both patient, caller and next of kin, complicating identifying all persons involved and collecting consent. The

situation can be dramatic and life-threatening, and many patients may no longer be alive at the time of the study. We applied for, and received, exemption of consent considering the importance of a complete material, the possible distress on patients and next of kin collecting consent, and that there was no intervention affecting the patients.

The first three studies were assessed by the Regional Committee for Medical and Health Research Ethics, and were deemed to be in no need of ethical approval (2011/756/REC West). The committee also exempted patients and EMCC operators' consent to pertaining to analysis of sensitive data. The studies were approved by the Data Protection Official for Research.

The fourth study was approved by the Regional Committee for Medical and Health Research Ethics (2013/982/REC West). The committee exempted patients' consent pertaining to analysis of sensitive data.

**Studies 1 and 3** included AMIS records with identifiable patient information in the form of civil registration numbers and addresses. The AMIS printouts have been kept locked up in archive cabinets, behind both key-locked and electronically locked doors. The questionnaires in **Study 2** were collected anonymously. The log recordings in **Study 3** were never extracted from the EMCCs, only listened through locally. Although no directly identifiable patient information was noted on the listening form, the information was indirectly identifiable as AMIS identification numbers were used as a link to the corresponding AMIS. Hence, the listening forms were kept locked up together with the AMIS printouts. The data from Haukeland University Hospital in **Study 4** contained civil registration numbers to enable linkage with AMIS, and was kept on an encrypted USB-memory-stick and locked up.

## 4 Main findings

This chapter includes a brief overview on the main findings from the four studies.


### 4.1 Study 1 – 113 epidemiology

These results were published in **Paper I**, together with the findings from **Study 2**.

The 72-hour study period yielded 3 294 AMIS records, of which 2 244 were included for analyses. The overall national 113 contact rate was 56/1 000 inhabitants per year (95% *CI* 53-57). Contact rates varied among the different EMCCs, from 33 (95% *CI* 21-45) to 114 (95% *CI* 87-141). There was variation in acute contact rates among the EMCCs, from 5 to 31/1 000 inhabitants per year, with an overall national acute contact rate of 21/1 000 per year. Assessment of the contact urgency category showed 37% acute, 34% urgent, and 27% non-urgent, while 1.5% of the contacts received no urgency assessment.

The most frequently used Index criterion, accounting for 20% of all contacts all over, was “6 – Unclear problem”. The variation in use of this criterion was 10–42%. Other criteria, like “5 – Ordered assignment” and “28 – Psychiatry – Suicide”, also varied much between different EMCCs, 0–22% and 0–19% respectively. Besides huge variation in frequency, the urgency assessments on Criteria 6 also differed between the EMCCs; acute 0–53%, urgent 0–67% and non-urgent 9–86% (Figure 4.1).

**Figure 4.1** Variation in urgency assessment and use of Index criteria card “06 Unclear problem” among the EMCCs.



Red = Acute, yellow = urgent, green = non-urgent.

## 4.2 Study 2 – Self-reported use of Index


These results were published in **Paper I**, together with the findings from **Study 1**.

All 19 EMCCs were represented in the material, consisting of 272 questionnaires after three reminders, a response rate of 63.4%. The typical operator was female, a registered nurse and worked in rotation with either the emergency ward or ambulances. The majority received initial training in the use of the Index when starting at the EMCC, but repetitions later on were rare. The median EMCC work experience was six years (Q1 3, 12).

Use of the Index ranged from 2.91 to 4.82, with an overall mean of 3.95 (*SD* 0.39) for all operators, corresponding closely to the questionnaire response format “often, > 75%” (=4). On the EMCC level, use of the Index varied from 3.7 (0.24) to 4.4 (0.39) ( $p < 0.001$ ) (Figure 4.2). When exploring possible factors associated with use of the

Index, rotation with ground ambulance, regular use of the Index and the perception of focus on use of the Index at the workplace all had a statistically significant effect on use of the Index ( $p < 0.005$ ). Adjusting for explanatory variables, only the negative effect of rotation with ambulances ( $\beta = -0.21, p < 0.002$ ) and the positive effect of focus on use of the Index at workplace ( $\beta = 0.46, p < 0.001$ ) were statistically significant.

**Figure 4.2** Variations in mean use of the Index among the different EMCCs,



The start page was used often or always (> 75% of real emergency calls) by 47% of the operators, but 93% always checked if the patient was awake and 46% if he was able to talk. “I find what I need in AMIS”, “I know it”, and “It takes too long” were the main reasons for not using the Index at all. The main reasons for not using the start page were “I know it”, “I prefer to go straight to the proper card” and “I find what I need in AMIS”.

Free text note explanations or supplements did not add further information relevant on the questions addressed in the study, and were hence not analysed.

### 4.3 Study 3 – Guideline adherence

These findings were published in **Paper II**. There were 174 acute and 125 urgent contacts. An ambulance was dispatched in 99% of the acute contacts ( $n=173$ ) and 87% of the urgent contacts ( $n=106$ ).

Guideline adherence was calculated based on six measured indicators: caller's phone number confirmed, location confirmed, patient's consciousness, patient's responsiveness, criteria compliance and communication. The mean overall guideline adherence was 2.41 (0.73), 80% of the maximum value of 3. Although the individual EMCCs varied in guideline adherence (75 to 89%), this variation was not statistically significant ( $p=0.073$ ).

Acute contacts had higher levels of overall guideline adherence, compared with urgent contacts ( $\rho=0.27$ ,  $p<0.0005$ ), but there was no correlation between overall guideline adherence and self-reported use of the Index, EMCC contact rate or population density.

Urgent contacts had a median response interval of 4:10 minutes compared to acute contacts with 2:01 minutes. This increase of 2:09 minutes was statistically significant ( $p<0.0005$ ). Increasing overall guideline adherence decreased EMD response interval ( $p<0.0005$ )

Criteria compliance was the only individual indicator with a statistically significant effect on EMD response ( $p=0.001$ ), with increased EMD response in the group with more criteria unchecked. EMD response interval was not influenced by confirmation



of location, telephone number or consciousness status, or by any of the communication challenges (language difficulties, caller non-cooperation or caller not on the scene).

#### 4.4 Study 4 - Stroke identification

These findings are submitted in **Paper III**.

Of the 519 stroke patients, 48% reached EMCC through the 113 line. Combining the population from Haukeland University Hospital and the 113 line criteria code A.27.03–06, we calculated a sensitivity for identifying a stroke patient at initial EMCC contact of 57.9% (95% *CI* 51.5, 64.1), a specificity of 99.1% (95% *CI* 98.9, 99.2), PPV of 45.7% (95% *CI* 40.1, 51.4) and a NPV of 99.4% (95% *CI* 99.3, 99.5) (Table 4.4). Figure 4.4 shows the overlap of stroke suspects, stroke patients, 113 line and health lines.


**Table 4.4** Cross tabulation of the 113 line: Patients with Index dispatch criteria code A.27.03–06 (stroke suspects) versus confirmed stroke diagnoses (stroke patients).

		Stroke patients <sup>1</sup>		
		Yes	No	Total
Stroke suspects <sup>2</sup>	Yes	143	170	313
	No	104	17 987	18 091
	Total	247	18 157	18 404

<sup>1</sup> True stroke condition: Confirmed stroke diagnoses at hospital discharge.

<sup>2</sup> Predicted stroke condition: Index dispatch criteria code A.27.03–06 at first EMCC contact.

**Figure 4.4** Illustration of the match between stroke suspects and stroke patients, 113 line and health lines.



#### 4.4.1 Stroke identification

Table 4.4.1 shows an overview of the Haukeland University Hospital stroke patients and their line of initial EMCC contact. The 113 line contacts are characterized by a high proportion of acute responses (84%) and stroke suspect dispatch codes (58%). The differences between the two populations were statistically significant with regards to urgency, dispatch codes and age groups ( $p < 0.0005$ ,  $p < 0.0005$  and  $p = 0.005$ , respectively).

**Table 4.4.1** Stroke patients and line of initial EMCC contact.

	Total	113 line		Health lines	
	<i>N</i>	<i>N</i>	%	<i>N</i>	%
<i>Total</i>	519	247	48	272	52
<i>Urgency</i>					
Acute		207	84	77	28
Urgent		31	13	73	27
Non-urgent		9	4	122	45
<i>Dispatch code</i>					
Stroke suspect		143	58	30	11
None		8	3	111	41
Dispatch code 5		25	10	113	42
Others		71	29	18	7
<i>Sex</i>					
Female		112	45	126	46
Male		135	55	146	54
<i>Stroke type</i>					
Infarction		192	78	224	82
Haemorrhage		36	15	22	8
TIA		19	8	26	10
<i>Age group</i>					
< 60		40	16	67	25
60–74		66	27	90	33
75–84		76	31	68	25
85 <		65	26	47	17

Further analyses of the 113 population showed that patients with haemorrhagic stroke were less likely to receive a stroke suspect code at initial EMCC contact, *OR* 0,4 (95% *CI* 0,2, 0,9,  $p=0.029$ ), while TIA patients were more likely to receive a stroke suspect code, *OR* 3.7 (95% *CI* 1.1, 13.2,  $p=0.042$ ).

In 85% of the 272 health line calls, there was a free text note mentioning stroke suspicion. The presence of stroke suspicion notes was associated with the dispatch code given; 97% of the stroke-suspect dispatch codes, 87% of the no dispatch codes, 86% of the “05 Transport reservations” dispatch codes and 44% of the other dispatch codes had free text notes mentioning stroke suspicion ( $p<0.0005$ ). Urgency was not associated with stroke suspicion notes.

#### 4.4.2 *Stroke prediction*

Of the 1 013 dispatch code 27 patients, 314 were true stroke patients, and 87.9% of these had received a stroke suspect code. The proportion of stroke patients among these four stroke suspect dispatch codes, A.27.03–07, ranged from 27.6 to 42.7%. Dispatch code A.27.04–“Sudden impaired strength in an arm or a foot” and admission to Voss hospital gave the highest *OR* for stroke in the multivariate regression analyses, 2.6 (95% *CI* 1.5, 4.5,  $p=0.001$ ) and 2.7 (95% *CI* 1.3, 5.8,  $p=0.011$ ) respectively. Age below 60 years had a reduced probability for stroke, *OR* 0.3 (95% *CI* 0.2, 0.5,  $p<0.0005$ ).

## 5 Discussion

### 5.1 Summary of main findings

The overall findings are that Index was used to a high degree by the operators, but that its ability to identify stroke patients was suboptimal.

The 113-epidemiology study showed a wide variation between the 19 EMCCs with regards to contact rates, specific acute rates and use of different Index criteria cards. The operators reported that they use the Index in more than 75% of the 113 calls, but there are variations in how the start page and other criteria cards are used between the EMCCs. The measured mean guideline adherence in the audio log study was 80%. Time to dispatch of resources decreased with higher levels of guideline adherence. In the stroke dispatch study criteria card 27 was tested for its ability to identify stroke patients: sensitivity of 57.9%, specificity of 98.7%, PPV of 37.1% and NPV of 99.4%. Only 48% of the stroke patients accessed emergency medical help through the 113 line.

### 5.2 Methodological considerations

The major methodological concern in the beginning of this project was the close interaction between the operator and the Index, and whether it was possible to distinguish the contribution of these two factors when looking at the dispatch outcome. As a criteria-based dispatch guideline rather than an algorithmic protocol, the Index was never meant to stand alone, so when **Studies 2** and **3** confirmed that the operators use the Index to a high degree, it supported the idea of considering the operator and the Index as a unity through the last study on its ability to identify and predict stroke.

### *5.2.1 Prehospital EMS research*

Most medical research faces challenges regarding patient ethics, but research on prehospital emergency medical services is complicated by yet another number of factors: critical time intervals, challenging environment, many different participants and lack of data conformity. The patients might be unconscious or suffering from supposedly life-threatening injuries or conditions, complicating patient consent to participate in studies. Time is critical, and data collection and interventions naturally receive less priority than patient care. Every situation provides different indoor and outdoor settings, and weather challenges, and a typical prehospital incident might include patient, next of kin, bystanders, GP on call, ambulance workers, rescuer and anaesthesiologist from HEMS. There is also a lack of uniformity, both in data collected and reported, but also in software solutions. This complicates not only the patient follow-through from initial location to final hospital discharge, but also comparison among different participants of the service.

### *5.2.2 AMIS as data source*

As the Index is a paper-based table fold-up generating no data on its use, AMIS data was used as a surrogate. **Studies 2** and **3** gave some conflicting views on the reliability of this data source as a surrogate on Index use. The Index criteria dispatch codes are all available directly in the AMIS display in a scroll-down menu, and this was stated to be the primary excuse when the Index was not used in **Study 2**. **Study 3**, on the other hand, found that the operators actually used the Index and its prompts to a high degree during the calls.

During registration of the AMIS entries, initially trying to log to which degree the GP on call had been involved in each case, it became clear that each EMCC uses AMIS coding differently.

In 2010 there was no national registry of AMIS data, 113 calls, Index criteria used or urgencies. Although these data were registered locally, extracting data electronically from the 19 different EMCCs seemed impossible for an external researcher. The resulting material of AMIS printouts collected for the epidemiology study (**Study 1**) was manually registered in an SPSS file, creating a local database with all information needed for the research project. This database later served as a background for the log recordings in **Study 3**.

A report on national 2012 EMCC data was published in 2013, providing overall data on population, contacts and response time, but no Index criteria data or urgency distributions for the 113 line separately (33). The 2013 report provided overall numbers on Index criteria cards and various presentations on 113 line calls, including urgency distribution (32). The Norwegian Directorate of Health has since then provided an interactive web page on EMCC activity, accessible for all to generate overall data extractions (26).

### *5.2.3 Questionnaire and listening form*

The questionnaire and listening form were both developed specifically for these studies (Appendices). The questionnaire was pretested among a small group of operators in an attempt to clarify possible misunderstandings, but not validated prior to the study. The main reason for this was the small overall operator population, and a desire to include all in the national study. The questionnaire study was vulnerable to recollection bias, asking the operators to recollect what they usually do. The “eagerness to please” bias, as the operators are expected to use the Index, was attempted minimized by providing anonymous return of the forms, directly to the researcher. All information and the reminders were distributed through the EMCC management, and this could be one of the reasons for the response rate of only 63.4%. This relatively low response rate opens for the possibility of a selection bias with regards to repliers versus non-repliers.

The listening form was developed from the quality control form used by Bergen EMCC, by simplifying the questions down to those necessary for the objectives of **Study 3**. Due to capacity issues, only one researcher visited the EMCCs participating in the study and listened to the included 113 calls. This could be regarded as both a strength, as all calls were assessed by the same investigator, and a weakness, as the assessments were not double-checked. This bias was presumed avoided by keeping the listening form simple, with yes/no/unnecessary and a time frame for each question, leaving little uncertainty about the findings. Whenever in doubt, a second researcher was asked for a second opinion.

#### *5.2.4 Strategic selection of EMCCs*

The invited EMCCs for **Study 3** were selected strategically in order to ensure that the findings would reflect the rest of the EMCCs. The selection was based on variation in size, geographic location, urban/rural, reputation and interest. All four health trusts were represented. The original idea was 6 centres with 50 calls each, acute and urgent. The AMIS material from **Study 1** was used as background data for the calls, and limited the number of acute and urgent calls available for selection. None of the EMCCs in the North health trust had 50 acute or urgent calls from this 72-hour period; hence two centres were included with 20 and 30 calls.

#### *5.2.5 Criteria cards*

The two most used criteria cards were “06 Unclear problem” and “05 Transport reservations”. From a researcher’s point of view, they are both deeply troublesome by providing material disturbances diminishing the validity of the data. To make things more complicated, the EMCCs don’t use the card similarly, as presented in **Study 4** on stroke patients. Some received dispatch codes from card 5, some from card 27.



Card 5 is used for organized ambulance transports, whether it's a planned non-urgent transport or an acute ride from a casualty clinic to the hospital. It's an administrative card, dealing with resource distribution alone in the form of orders.

Card 6 covers a whole range of different situations, from chaotic and unclear situations with too little information for the operator to designate a proper card, to well-defined situations where there just is no other criteria card that fits the situation. As shown in **Study 1**, the different EMCCs use this card very differently; 10–42% variation in frequency, and while some EMCCs use the acute criteria codes from this card most frequently, others use the non-urgent codes. The card seems to function as an easy solution if the operator doesn't solve the situation otherwise.

Criteria card 27 was selected for the validation study because it was the one card in the Index most closely associated to a diagnosis or diagnosis group. But, as the main card includes symptoms beyond those associated with stroke alone, the four dispatch codes A.27.03–06 were selected for the purpose of measuring stroke identification ability.

### *5.2.6 Missing data*

Missing data is never a good thing in research, as it reduces the validity of the findings. In connection with the Norwegian Directorate of Health undertaking the responsibility for reporting on national EMCC data in 2013, a manual for registration of EMCC data was developed (62). The intention was to secure homogeneous and comparable data, resulting in registration of urgency and criteria dispatch code becoming mandatory for all incidents registered in AMIS.

## 5.3 Discussion of the results

### 5.3.1 *Use of the Index*

The variation in 113 contact rates among the different EMCCs, from 33 to 114 calls/1 000 inhabitants a year, indicates that the population uses 113 differently in the different regions. Possible reasons for this could be varying thresholds for what is regarded as acute and what is urgent, or local preferences related to availability of the casualty clinic and LEMC. The overall 113 urgency distribution: acute 37%, urgent 34% and non-urgent 27%, taken together with the findings of a primary emergency health care study: acute 2.3%, urgent 21.1% and non-urgent 76.6%, show that the population as a whole knows which level to address depending on the degree of medical emergency (22).

The epidemiology study also reported differences in acute contact rates, which, assuming that the distribution of acute illnesses and accidents are equally spread nationwide, means that the EMCCs differ in determining urgency. The mean acute contact rate of 21/1 000 inhabitants a year can compare to the annual rate of 25, found among three Norwegian EMCCs in 2007 (19), and of 17 on a national level in Denmark (43). The Danish study also found a difference among their included areas: 13–21 acute ambulance dispatches a year.

There is also a large difference among the EMCCs in the use of criteria card “06 Unclear problem”, both with regards to the total use of this card and the variation of urgencies it was used with. This was the single most frequently used criteria card in 113 calls, as problematized in methodological considerations, used in 20% of the contacts. The same result was found in Copenhagen in 2013, with “Unclear problem” accounting for 19% of the calls (63). The criteria card covers a whole range of different situations, from unclear situations with little information to well-defined situations where no other criteria card fit.

The mean self-reported use of Index value, 3.95 corresponded closely to the response format “> 75% of the cases”, meaning the operators use the Index in a majority of the emergency calls. The variations were large also here, both on an individual operator level and on an EMCC level. The explanatory factors we investigated accounted for only 23% of this variation. The main factor associated with increased use of the Index was the perception by the operator that their EMCC focused on use of the Index. This increased use of the Index by 0.5 (10%). Although not significantly affecting use of the Index, the 19% reporting that their EMCC performed regular repetition on use of the Index was discouraging.

It was expected that use of the Index would diminish with experience, but this effect was unexpectedly small with only a 0.01 decrease in use of the Index per year working at the EMCC. Equally unexpected was the negligible effect the EMCC had on use of the Index, as this was thought to be the natural explanation for the variation found in the use of the Index.

The main factor associated with decreased use of the Index was rotational work at ground ambulance, reducing use of the Index by 0.2. This could be explained by these operators tending to use their own experience over the Index more often than operators without recent prehospital experience.

The > 75% use of Index found in **Study 2** corresponds well with the 80% guideline adherence found in **Study 3**. A prior Norwegian Index study from 2005 reported a self-reported Index compliance of 99% and a measured compliance of 64% (40). The consistency of our findings, and the fact that they both lay between the two 2005 findings, could indicate that our measures on guideline adherence are realistic. The reversed findings of our results, that the measured use is higher than the self-reported, could also indicate a positive development towards both higher actual use of the Index, and a more realistic reflection on own use. As CBD guidelines are developed to assist health care personnel in the decision process rather than to define the next step of action, the score on overall guideline adherence can never reach 100%. The optimal

level of guideline adherence is unknown, and the studies addressing CBD compliance internationally are non-existing.

Although the algorithmic dispatch systems facilitate high protocol compliance through electronic systems developed to monitor and increase compliance and other quality markers (38,56,64-67), less than 3% of the MPDS were Accredited Centers of Excellence in 2013 (68), which among other quality measurements require a minimum of 90–95% protocol compliance.

Less than half of the operators reported using the start page often or always. The reasons given for not using it were that they knew it by heart, and that they could find the key words they needed through AMIS. The low percentage of operators reporting actually asking if the patient is awake and able to talk, confirms that neither memory nor AMIS equals the start page. This question was changed from “awake and can breathe” to “awake and able to talk” in the 2009 revision of the Index (28). Confidence in AMIS providing the necessary key words to assess the situation was also the main reason for not using the Index as a whole. This raises concerns, as AMIS does not provide support in the decision making or give advice for the public or health personnel at scene. Discarding the guidelines completely in favour of memory, personal skills and experience constitutes a potential hazard of getting sidetracked or losing vital information due to unstructured call handling. This is one of the main risks highlighted by critics of CBD (34,65). Another given reason for not using the Index was the constant and cumbersome switching of hands and eyes between the electronic AMIS and the paper-based Index.

The relationship between clinical outcome and guidelines is known to be an important motivational factor for guideline adherence (39). Ambulance personnel arriving on scene can give the EMCC feedback on the patients, providing the operator the opportunity to adjust his/her perception of the situation initially formed by 113 call. The operators never receive information on the actual patient outcome though. This means that they have no possibility to adjust personal dispatch manners according to

patient outcome, unless the mistake is of such a fatal character that it ends up as a formal complaint.

Although an acceptable level of overall dispatch guideline adherence is unknown, it is not difficult to agree that start page compliance, including confirmation of patient location and consciousness, should be 100% in an emergency medical call. Despite the low reported use of start page, more than 90% of the operators in **Study 2** reported that they check if the patient is awake. **Study 3** found that the operators clarified consciousness within 1 minute in 83% of the cases. A 2014 study comparing an American MPDS with our CBD Index in cardiac arrest patients, reported a guideline adherence of 97% for successful clarifying level of consciousness (41). This difference can be explained partly by their material including cardiac arrest patients only, where the patients' loss of consciousness probably is the major reason for the caller to access 113 in the first place. Another explanation could be the wide variety between our EMCCs, ranging from 73 to 92%, when it comes to confirming consciousness. Although this variety was statistically insignificant in our study, the EMCC with the highest score corresponds to the EMCC participating in the 2014 study. Larger EMCCs allow for higher call processing rates, which is shown to increase dispatch performance: Operators processing more cardiac arrest calls recognized the cardiac arrest faster, dispatched first responding units faster, gave advice on cardiopulmonary resuscitation more often, and received bystander cooperation more often than operators processing few cardiac arrest calls (69). This supports the idea of fewer and larger EMCC units.

Consciousness and responsiveness is vital information for detecting possible life-threatening situations like cardiac arrest or breathing difficulties. These are situations where awareness and instructions from the operator might influence the outcome significantly. Failure to recognize cardiac arrest during the emergency call delayed both ambulance dispatch and arrival on the scene, and decreased three-month survival from 14 to 5% in the Netherlands (70). The Dutch dispatchers were supposed to clarify location, phone number, patient consciousness and breathing, and the study found

failure to ask whether and how the patient was breathing to be the primary reason for not recognizing the cardiac arrest.

Balancing over-triage and under-triage is vital with regards to resource allocation. A certain over-triage is generally accepted in the prehospital EMS, especially in dispatch where there are restricted possibilities to explore and assess the situation properly. Under-triage on the other hand, holds a potential risk for unnecessary time loss and suboptimal level of care, and is less accepted; especially for time-critical conditions like cardiac arrest, respiratory failure, trauma, acute coronary syndromes or stroke (the first hour quintet) (71). Low criteria compliance is a risk factor for under-triage, because important information might remain unrevealed. Experiences from Norwegian GPs participating in prehospital emergency situations show that the GPs downgrade 42% and upgrade 11% of the patients when triaging them on the scene compared to the initial EMCC dispatch (72). Acute abdominal cases were at highest risk of initial under-triage.

Patients initially coded as non-urgent ending up as fatalities have been reported in both Italy and Denmark (73,74). The Italian study found that these calls were predominantly made by next of kin reporting no life-threatening symptoms, but they were also shorter than other emergency calls and characterized by inadequate collection of vital information by the operators. The Danish under-triage study found that 152 (0.16% of the low-acuity) patients died the same day as the 112 call, and 12% of these were considered potentially preventable if the call had been assessed as acute. A Swedish study comparing CBD-based priority codes from the dispatch centre, with triage by the ambulance on scene, found that 4.8% of the patients were under-triaged (75). A similar finding of 4.6% under-triaged patients was recently found in Switzerland (76).

While the importance of not missing out on vital information is undebatable, the dispatch guidelines effect on time has been debated for years, with supporters claiming it saves time and critics claiming that it adds time to the dispatch process. The findings

from **Study 3** support that the Index could save time if used properly. Criteria compliance was the only Index indicator out of six explored in **Study 3** that had a statistically significant effect on response interval. The main effect on the response interval occurred between 1 and 2 unconfirmed criteria, with an increase of 1 minute and 31 seconds from 0 unconfirmed criteria to 2+ unconfirmed criteria. This indicates that the margins are small. The combined outcome variable “overall guideline adherence” showed an increase response interval of 2 minutes and 16 seconds between the lowest and highest value group.

### *5.3.2 Stroke identification*

Ideally, all patients experiencing a stroke should get in contact with 113 as soon as possible, and be identified at the EMCC as a stroke patient. Figure 4.4 shows that in reality, this is not the case.

Although validation studies of CBD guidelines have been problematized due to the unknown factor of operator–guideline interaction, several studies have aimed to assess stroke dispatch validity in CBD guidelines recent years. In Sweden, sensitivity for stroke recognition was measured to be 64% (59), while a Danish study found a sensitivity of 66.2% and a PPV of 30.2% (60). These results are comparable to our findings, except from our sensitivity being somewhat lower. As our dispatch guidelines are quite similar, in particular the Norwegian and Danish Index, this difference might reflect a difference in use of guidelines, confirming concerns raised earlier about the CBD being vulnerable to individual EMDs and EMCCs (37,38).

Training operators in stroke recognition has been shown to raise the proportion of identified stroke patients in algorithmic dispatch protocol systems (77). Such training could be expected to be even more effective in the CBD, as this dispatch system utilizes the operator’s individual skills and competence to a wider extent than the algorithmic systems. With regards to Index, increased stroke awareness may lead the

operator to criteria card 27 earlier, and hence decrease EMD response time further as well as potentially identifying more stroke patients.

The importance of a valid stroke dispatch is twofold. First and foremost, stroke recognition leads to a faster dispatch response and improves time to hospital arrival (49-51,78). This is confirmed in our findings; 91.9% of the patients dispatched as stroke suspects received an acute response, while the stroke patients with other dispatch codes received acute responses in 55.1% and urgent responses in 39.2%.

Secondly, an increasing focus on earlier prehospital diagnostics, and in the future maybe even prehospital thrombolysis, through dedicated stroke ambulances equipped with computed tomography scanners (52-54), presupposes a valid stroke dispatch.

The four dispatch codes selected for **Study 4**, A.27.03–06, are closely associated with the FAST stroke symptoms. The 2015 study on stroke patients from Sweden found that FAST symptoms were presented in 80% of the calls dispatched as stroke, compared to only 35% in those with other dispatch codes (49). Studies from the United Kingdom and Australia found quite opposite numbers, with FAST symptoms reported in less than 5% of the calls in the UK (79) and FAST symptoms reported spontaneously in 11% (facial droop) to 41 % (speech problems) in Australia (48). Our findings of nearly 90% identification of the stroke patients by these four dispatch codes compares to the Swedish numbers. Direct comparison with the other studies is difficult, as the dispatch systems are different. Fall is not mentioned as a criterion in Index, but is reported in the literature to be a known predictor of stroke (17–38%), as well as the word “stroke” mentioned by the caller (20–49%) (48,59,79-82).

**Study 4** revealed that less than half of the stroke patients accessed the EMCC through the public emergency medical 113 line. Similar results was found in a Dutch study on different prehospital paths where almost half of the stroke patients contacted their GP initially (47). In the Netherlands, GP involvement increased the median time between seeking medical advice and arrival at hospital, compared to ambulance involvement



alone. Whether the same result would be found in Norway, where GP involvement also includes GP call-outs to the patient scene, is unknown. Also in the UK, a majority of the stroke patients' first medical access point was a general practitioner (83). GPs have also been found to have lower precision level of stroke prediction than paramedics and emergency physicians (84,85), indicating that GPs might represent a potential delay to treatment for stroke patients. The majority of public stroke warning campaigns report positive intervention effects, but are generally limited by methodological weaknesses (86). A Japanese study of a two year long stroke education campaign, found an increased association between knowledge on stroke symptoms and intention to call an ambulance (87). There is generally no tradition for studying the effect of public awareness campaigns in Norway, and stroke awareness campaigns are no exemption.

The majority of the stroke patients whose initial EMCC contact was through primary care personnel received either criteria card "05 Transport reservations" dispatch codes, or no dispatch codes at all. The assessments of a GP or out-of-hours doctor on scene with the patient are rarely overturned by the operator, thus the Index is not used as it would have been the with a 113 call. A free text note mentioning stroke suspicion was listed in 85% of these cases, giving the GPs a far better prediction level than previously reported from other countries (84,85). The majority of these patients received urgent or non-urgent response, assumedly due to a time delay beyond the thrombolysis treatment window of four hours from symptom onset.

Studies on patient behaviour as a reason for the delays reveal that although there is a good awareness of stroke of the commonest symptoms and the need for emergency response, only half the patients recognize it themselves, according to a 2010 systematic review (83). Severity of the symptoms, along with the support of family and bystanders both affect patient delay, which can vary from an hour to several days (88). A recent British qualitative study on TIA patients and how primary health care services contributed to time delay, found two factors associated with delay in general: availability of same day appointments and the role of the receptionist in identifying

urgent cases (89). Whether the delay in Norway is due to an initial patient delay, or to accessing the wrong level of emergency medical care is yet unknown. Most likely, it is a combination of both.

## 6 Conclusions

This thesis has contributed to increased knowledge about the use of the Norwegian Index for Emergency Medical Assistance, and about the precision level of one of the criteria cards. We were able to answer our aim and objectives:

1. There was a large variation between the EMCCs with regard to both contact patterns and use of the Index.
2. Self-reported use of the Index was relatively high, but there were large individual differences. A clear focus on use of Index at the EMCC was associated with increased use.
3. Measured guideline adherence was higher than self-reported use of the Index. Low overall guideline adherence increased the EMD response intervals.
4. The ability of the Index to identify or predict stroke patients was modest. The 113 line was the primary EMCC access point for less than half of the stroke patients.

## **7 Future aspects**

### **7.1 Implications of the findings**

It seems that the criteria-based model of dispatch guidelines works well with the EMCC operators, allowing them to utilize their education, experience and skills yet still use the Index. The finding that focus at the workplace is a factor for improved use of the Index, might encourage management and personnel at the EMCCs to focus further on the use of Index through repetition, case reviews and other reminders.

AMIS was identified as a factor decreasing use of the Index, as it provides easy access to the criteria codes without having to manually look them up in the paper-based Index version. An obvious and necessary improvement of the Index in the future is the electronic version that has been “just around the corner” since 2010. This will not only simplify the work flow for the individual operators, but also secure valid data on how the Index is actually used and enable larger validity studies of higher quality. It is likely to believe that this will improve use of the Index as well.

There is an upcoming revision of the Index, which includes changes to both the start page and criteria card 27, among others. Ability of the present Index to identify stroke patients is suboptimal, and it is of great importance that the new version of the Index is evaluated with regards to how it identifies these patients. Increased stroke awareness among the operators might lower the threshold for using criteria card 27.

The studies have shown that there is an overall and immediate need for improved data quality. Work has already been done during the last couple of years to homogenize the use of AMIS codes, and determining urgency and dispatch code has been made mandatory. Both steps will strengthen future EMCC-based research.

The national stroke registry was implemented in 2013, the cardiac arrest registry in 2015, and the national trauma registry commenced data registration as of January 2015. Use of such databases will provide easier access to data and enable larger studies.

## **7.2 Future research**

Future research should strive to utilize registry resources and explore the effect of the imminent Index changes; complete numbers on urgencies and dispatch codes, a revised Index and hopefully in the near future, an electronic version.

Our stroke study was only the beginning of an Index validation process. With the use of electronic Index data and diagnosis registries, it would be possible to validate the Index as a whole.

Some interesting questions raised by this project, but not within its scope, are why the majority of stroke patients or their next of kin choose not to access the 113 line directly, and the time delays of different prehospital paths for stroke patients in Norway.

## References:

1. Al-Shaqsi S. Models of international emergency medical service (EMS) systems. *Oman Med J*. 2010;25:320–3.
2. Pozner CN, Zane R, Nelson SJ, Levine M. International EMS systems: The United States: past, present, and future. *Resuscitation*. 2004;60:239–44.
3. Trevithick S, Flabouris A, Tall G, Webber CF. International EMS systems: New South Wales, Australia. *Resuscitation*. 2003;59:165–70.
4. Lockey D. International EMS systems: Geographical lottery and diversity but many common challenges. *Resuscitation*. 2009;80:722.
5. Hung KKC, Cheung CSK, Rainer TH, Graham CA. EMS systems in China. *Resuscitation*. 2009;80:732–5.
6. Black JJM, Davies GD. International EMS systems: United Kingdom. *Resuscitation*. 2005;64:21–9.
7. Langhelle A, Lossius HM, Silfvast T, Björnsson HM, Lippert FK, Ersson A, et al. International EMS systems: The Nordic countries. *Resuscitation*. 2004;61:9–21.
8. Adnet F, Lapostolle F. International EMS systems: France. *Resuscitation*. 2004;63:7–9.
9. Papaspyrou E, Setzis D, Grosomanidis V, Manikis D, Boutlis D, Ressos C. International EMS systems: Greece. *Resuscitation*. 2004;63:255–9.
10. Symons P, Shuster M. International EMS systems: Canada. *Resuscitation*. 2004;63:119–22.
11. Gomes E, Araújo R, Soares-Oliveira M, Pereira N. International EMS systems: Portugal. *Resuscitation*. 2004;62:257–60.
12. MacFarlane C, van Loggerenberg C, Kloeck W. International EMS systems in South Africa – past, present, and future. *Resuscitation*. 2005;64:145–8.
13. Callaham M. Quantifying the scanty science of prehospital emergency care. *Ann Emerg Med*. 1997;30:785–90.
14. Lerner EB, Weik T, Edgerton EA. Research in prehospital care: Overcoming the barriers to success. *Prehosp Emerg Care*. 2016;20:448–53.
15. Jones CW, Hunold KM, Isaacs CG, Platts-Mills TF. Randomized trials in emergency medicine journals, 2008 to 2011. *Am J Emerg Med*. 2013;31:231–5.

16. PubMed [Internet]. [Cited 2016 Dec 5]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/>
17. Sandvik H, Hunskår S. Årsstatistikk fra legevakt 2015. (Annual statistics from Out-of-hours services 2015). National Centre for Emergency Primary Health Care, Uni Research Health. 2016. Available from: <https://bora.uib.no/handle/1956/11953>. Norwegian.
18. Ministry of Health and Care Services. Akuttmedisinforordningen. (Emergency Medicine Regulation.) 2015. Available from: [https://www.regjeringen.no/contentassets/bde3821703d44038a98b80923a1083ef/rundskriv\\_akuttmedisinforordningen\\_300415.pdf](https://www.regjeringen.no/contentassets/bde3821703d44038a98b80923a1083ef/rundskriv_akuttmedisinforordningen_300415.pdf). Norwegian.
19. Zakariassen E, Hunskaar S. Involvement in emergency situations by primary care doctors on-call in Norway – a prospective population-based observational study. *BMC Emerg Med*. 2010;10:5.
20. Zakariassen E, Sandvik H, Hunskaar S. Norwegian regular general practitioners' experiences with out-of-hours emergency situations and procedures. *Emerg Med J*. 2008;25:528–33.
21. Morken T, Midtbø V, Zachariassen S. Legevaktorganisering i Norge. Rapport fra Nasjonalt legevaktregister 2014. (Organization of Out-of-hour services in Norway 2014). National Centre for Emergency Primary Health Care, Uni Research Health. 2014. Available from: <https://bora.uib.no/handle/1956/8352>. Norwegian.
22. Zakariassen E, Hansen EH, Hunskaar S. Incidence of emergency contacts (red responses) to Norwegian emergency primary healthcare services in 2007 – a prospective observational study. *Scand J Trauma Resusc Emerg Med*. 2009;17.
23. Littlewood N, Parker A, Hearn S, Corfield A. The UK helicopter ambulance tasking study. *Injury*. 2010;41:27–9.
24. McQueen C, Smyth M, Fisher J, Perkins G. Does the use of dedicated dispatch criteria by Emergency Medical Services optimise appropriate allocation of advanced care resources in cases of high severity trauma? A systematic review. *Injury*. 2015;46:1197–206.
25. Gordon K, Swain A, Thirkell C, Bailey M, Greenberg D. The Wellington Life Flight Helicopter Emergency Medical Service (HEMS): a retrospective audit against new Ministry of Health criteria. *N Z Med J*. 2014;127:30–42.
26. Emergency Medical Communication Centres – Activity data [Internet]. [Cited 2016 Nov 8]. Available from: <https://amk.beekeeper.no/dashboard>. Norwegian.
27. Østerås Ø, Brattebø G, Heltne J-K. Helicopter-based emergency medical

- services for a sparsely populated region: A study of 42,500 dispatches. *Acta Anaesthesiol Scand*. 2016;60:659–67.
28. Norwegian Medical Association. Norsk Indeks for Medisinsk Nødhjelp. (Norwegian Index for Emergency Medical Assistance). 3rd ed. Stavanger: Laerdal Medical A/S; 2009. Norwegian.
  29. Directorate for Emergency Communication [Internet]. [Cited 2016 Dec 5]. Available from: <http://www.dinkom.no/en/>
  30. Ministry of Health and Care Services. Sentrale elementer vedrørende organisering av AMK-sentralene. (Key elements regarding the organization of the EMCCs.) 2016. Available from: [https://www.regjeringen.no/contentassets/b332572e94304549b8b6db04ad617f76/amk\\_utredning\\_150916.pdf](https://www.regjeringen.no/contentassets/b332572e94304549b8b6db04ad617f76/amk_utredning_150916.pdf). Norwegian.
  31. Statistics Norway [Internet]. [Cited 2016 Oct 18]. Available from: [www.ssb.no](http://www.ssb.no). Norwegian.
  32. Norwegian Directorate of Health. Årsrapport 2013: Akuttmedisinske kommunikasjonsentraler – AMK. (Emergency Medical Communication Centres 2013.) 2014. Available from: <https://helsedirektoratet.no/Lists/Publikasjoner/Attachments/54/Akuttmedisinske-kommunikasjonssentraler-AMK-arsrapport-2013-IS-2198.pdf>. Norwegian.
  33. Norwegian Directorate of Health. Nasjonale data fra akuttmedisinske kommunikasjonsentraler. Årsrapport 2012. (Emergency Medical Communication Centres – National data 2012.) 2013. Norwegian.
  34. Clawson JJ, Martin RL, Hauert SA. Protocols vs. guidelines. Choosing a medical-dispatch program. *Emerg Med Serv*. 1994;23:52–60.
  35. Gardett I, Clawson J, Scott G, Barron T, Patterson B, Olola C. Past, present, and future of emergency dispatch research: A systematic literature review. *Ann Emerg Disp Resp* 2013;1:29–42.
  36. Clawson J, Olola CHO, Heward A, Scott G, Patterson B. Accuracy of emergency medical dispatchers' subjective ability to identify when higher dispatch levels are warranted over a Medical Priority Dispatch System automated protocol's recommended coding based on paramedic outcome data. *Emerg Med J*. 2007;24:560–3.
  37. Grusd E, Kramer-Johansen J. Does the Norwegian emergency medical dispatch classification as non-urgent predict no need for pre-hospital medical treatment? An observational study. *Scand J Trauma Resusc Emerg Med*. 2016;24:65.



38. Clawson JJ, Cady GA, Martin RL, Sinclair R. Effect of a comprehensive quality management process on compliance with protocol in an emergency medical dispatch center. *Ann Emerg Med.* 1998;32:578–84.
39. Ebben RHA, Vloet LCM, Verhofstad MHJ, Meijer S, Mintjes-de Groot JAJ, van Achterberg T. Adherence to guidelines and protocols in the prehospital and emergency care setting: A systematic review. *Scand J Trauma Resusc Emerg Med.* 2013;21:9.
40. Lorem T, Elisabeth S, Lars W. Handling of drug-related emergencies: an evaluation of emergency medical dispatch. *Eur J Emerg Med.* 2009;16:37–42.
41. Hardeland C, Olasveengen TM, Lawrence R, Garrison D, Lorem T, Farstad G, et al. Comparison of Medical Priority Dispatch (MPD) and Criteria Based Dispatch (CBD) relating to cardiac arrest calls. *Resuscitation.* 2014;85:612–6.
42. Culley LL, Henwood DK, Clark JJ, Eisenberg MS, Horton C. Increasing the efficiency of emergency medical services by using criteria based dispatch. *Ann Emerg Med.* 1994;24:867–72.
43. Andersen MS, Johnsen SP, Sørensen JN, Jepsen SB, Hansen JB, Christensen EF. Implementing a nationwide criteria-based emergency medical dispatch system: A register-based follow-up study. *Scand J Trauma Resusc Emerg Med.* 2013;21:53.
44. Matic I, Golub M. Implementation of Croatian Emergency Call Reception Index modified according to Norwegian Index of Medical Emergencies. *Sestrinski glasnik.* 2014;19:63–5.
45. Emberson J, Lees KR, Lyden P, Blackwell L, Albers G, Bluhmki E, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: A meta-analysis of individual patient data from randomised trials. *Lancet.* 2014;384:1929–35.
46. Wardlaw JM, Murray V, Berge E, del Zoppo GJ. Thrombolysis for acute ischemic stroke, Update August 2014. *Stroke.* 2014;45:e222–5.
47. Doggen CJM, Zwerink M, Droste HM, Brouwers PJAM, van Houwelingen GK, van Eenennaam FL, et al. Prehospital paths and hospital arrival time of patients with acute coronary syndrome or stroke, a prospective observational study. *BMC Emerg Med.* 2016;16:3.
48. Mosley I, Nicol M, Donnan G, Patrick I, Dewey H. Stroke symptoms and the decision to call for an ambulance. *Stroke.* 2007;38:361–6.
49. Berglund A, Svensson L, Sjostrand C, Arbin von M, Euler von M,

- Wahlgren N, et al. Higher prehospital priority level of stroke improves thrombolysis frequency and time to stroke unit: The Hyper Acute STroke Alarm (HASTA) Study. *Stroke*. 2012;43:2666–70.
50. Puolakka T, Strbian D, Harve H, Kuisma M, Lindsberg PJ. Prehospital phase of the stroke chain of survival: A prospective observational study. *J Am Heart Assoc*. 2016;5:e002808–9.
51. Caceres JA, Adil MM, Jadhav V, Chaudhry SA, Pawar S, Rodriguez GJ, et al. Diagnosis of stroke by emergency medical dispatchers and its impact on the prehospital care of patients. *J Stroke Cerebrovasc Dis*. 2013;22:e610–4.
52. Wendt M, Ebinger M, Kunz A, Rozanski M, Waldschmidt C, Weber JE, et al. Improved prehospital triage of patients with stroke in a specialized stroke ambulance: Results of the pre-hospital acute neurological therapy and optimization of medical care in stroke study. *Stroke*. 2015;46:740–5.
53. Kostopoulos P, Walter S, Haass A, Papanagiotou P, Roth C, Yilmaz U, et al. Mobile stroke unit for diagnosis-based triage of persons with suspected stroke. *Neurology*. 2012;78:1849–52.
54. Weber JE, Ebinger M, Rozanski M, Waldschmidt C, Wendt M, Winter B, et al. Prehospital thrombolysis in acute stroke: results of the PHANTOM-S pilot study. *Neurology*. 2013;80:163–8.
55. Parker SA, Bowry R, Wu T-C, Noser EA, Jackson K, Richardson L, et al. Establishing the first mobile stroke unit in the United States. *Stroke*. 2015;46:1384–91.
56. Ramanujam P, Guluma KZ, Castillo EM, Chacon M, Jensen MB, Patel E, et al. Accuracy of stroke recognition by emergency medical dispatchers and paramedics –San Diego experience. *Prehosp Emerg Care*. 2008;12:307–13.
57. Buck BH, Starkman S, Eckstein M, Kidwell CS, Haines J, Huang R, et al. Dispatcher recognition of stroke using the National Academy Medical Priority Dispatch System. *Stroke*. 2009;40:2027–30.
58. Deakin CD, Alasaad M, King P, Thompson F. Is ambulance telephone triage using advanced medical priority dispatch protocols able to identify patients with acute stroke correctly? *Emerg Med J*. 2009;26:442–5.
59. Berglund A, Euler von M, Schenck-Gustafsson K, Castrén M, Bohm K. Identification of stroke during the emergency call: A descriptive study of callers' presentation of stroke. *BMJ Open*. 2015;5:e007661.
60. Viereck S, Møller TP, Iversen HK, Christensen H, Lippert F. Medical dispatchers recognise substantial amount of acute stroke during emergency calls. *Scand J Trauma Resusc Emerg Med*. 2016;24:89.

61. Ellensen EN, Hunskaar S, Wisborg T, Zakariassen E. Variations in contact patterns and dispatch guideline adherence between Norwegian emergency medical communication centres – A cross-sectional study. *Scand J Trauma Resusc Emerg Med.* 2014;22:2.
62. Norwegian Directorate of Health. Registreringsveileder: Opplysninger fra AMK-sentralene. (Registration manual – EMCC data.) 2013. Available from: <https://helsedirektoratet.no/Lists/Publikasjoner/Attachments/635/Registreringsveileder-opplysninger-fra-amk-sentralene-rapportering-til-norsk-pasientregister-npr-IS-2042.pdf>. Norwegian.
63. Møller TP, Ersbøll AK, Tolstrup JS, Østergaard D, Viereck S, Overton J, et al. Why and when citizens call for emergency help: An observational study of 211,193 medical emergency calls. *Scand J Trauma Resusc Emerg Med.* 2015;23:88.
64. Hinchey P, Myers B, Zalkin J, Lewis R, Garner D. Low acuity EMS dispatch criteria can reliably identify patients without high-acuity illness or injury. *Prehosp Emerg Care.* 2007;11:42–8.
65. Heward A, Damiani M, Hartley-Sharpe C. Does the use of the Advanced Medical Priority Dispatch System affect cardiac arrest detection? *Emerg Med J.* 2004;21:115–8.
66. Clawson J, Olola C, Heward A, Patterson B, Scott G. Ability of the Medical Priority Dispatch System Protocol to predict the acuity of “unknown problem” dispatch response levels. *Prehosp Emerg Care.* 2009;12:290–6.
67. International Academies of Emergency Dispatch: Accreditation [Internet]. [Cited 2016 Nov 11]. Available from: <http://www.emergencydispatch.org/Accred20PointsMedical>
68. Heward A. Dispatch research: Someone, somewhere needs to know. *Ann Emerg Dispatch Response.* 2013;1:5–8.
69. Kuisma M, Boyd J, Väyrynen T, Repo J, Nousila-Wiik M, Holmström P. Emergency call processing and survival from out-of-hospital ventricular fibrillation. *Resuscitation.* 2005;67:89–93.
70. Berdowski J, Beekhuis F, Zwinderman AH, Tijssen JGP, Koster RW. Importance of the first link: Description and recognition of an out-of-hospital cardiac arrest in an emergency call. *Circulation.* 2009;119:2096–102.
71. Krafft T, Castrillo-Riesgo LG, Edwards S, Fischer M, Overton J, Robertson-Steel I, et al. European Emergency Data Project (EED Project): EMS data-based Health Surveillance System. *Eur J Public Health.*

- 2003;13:85–90.
72. Rørtveit S, Meland E, Hunskaar S. Changes of triage by GPs during the course of prehospital emergency situations in a Norwegian rural community. *Scand J Trauma Resusc Emerg Med.* 2013;21:89.
73. Palma E, Antonaci D, Coli A, Cicolini G. Analysis of emergency medical services triage and dispatch errors by registered nurses in Italy. *J Emerg Nurs.* 2014;40:476–83.
74. Andersen MS, Johnsen SP, Hansen AE, Skjaereth E, Hansen CM, Sørensen JN, et al. Preventable deaths following emergency medical dispatch – An audit study. *Scand J Trauma Resusc Emerg Med.* 2014;22:74.
75. Ek B, Edström P, Toutin A, Svedlund M. Reliability of a Swedish pre-hospital dispatch system in prioritizing patients. *Int Emerg Nurs.* 2013;21:143–9.
76. Dami F, Golay C, Pasquier M, Fuchs V, Carron P-N, Hugli O. Prehospital triage accuracy in a criteria based dispatch centre. *BMC Emerg Med.* 2015;15:32.
77. Watkins CL, Leathley MJ, Jones SP, Ford GA, Quinn T, Sutton CJ, et al. Training emergency services’ dispatchers to recognise stroke: An interrupted time-series analysis. *BMC Health Serv Res.* 2013;13:318.
78. Watkins CL, Jones SP, Leathley MJ, Ford GA, Quinn T, McAdam JJ, et al. Emergency Stroke Calls: Obtaining Rapid Telephone Triage (ESCORTT) – A programme of research to facilitate recognition of stroke by emergency medical dispatchers. Programme Grants for Applied Research. Southampton (UK): NIHR Journals Library; 2014;2:1–124.
79. Jones SP, Carter B, Ford GA, Gibson JME, Leathley MJ, McAdam JJ, et al. The identification of acute stroke: An analysis of emergency calls. *Int J Stroke.* 2013;8:408–12.
80. Handschu R, Poppe R, Rauss J, Neundorfer B, Erbguth F. Emergency calls in acute stroke. *Stroke.* 2003;34:1005–9.
81. Leathley MJ, Jones SP, Gibson JME, Ford GA, McAdam JJ, Quinn T, et al. “Can you send an ambulance please?”: A comparison of callers’ requests for emergency medical dispatch in non-stroke and stroke calls. *Emerg Med J.* 2014;31:e25–8.
82. Krebes S, Ebinger M, Baumann AM, Kellner PA, Rozanski M, Doepp F, et al. Development and validation of a dispatcher identification algorithm for stroke emergencies. *Stroke.* 2012;43:776–81.

83. Lecouturier J, Murtagh MJ, Thomson RG, Ford GA, White M, Eccles M, et al. Response to symptoms of stroke in the UK: A systematic review. *BMC Health Serv Res.* 2010;10:355.
84. Harbison J, Hossain O, Jenkinson D, Davis J, Louw SJ, Ford GA. Diagnostic accuracy of stroke referrals from primary care, emergency room physicians, and ambulance staff using the face arm speech test. *Stroke.* 2003;34:71–6.
85. Karliński M, Gluszkiewicz M, Członkowska A. The accuracy of prehospital diagnosis of acute cerebrovascular accidents: An observational study. *Arch Med Sci.* 2015;11:530–5.
86. Mellon L, Doyle F, Rohde D, Williams D, Hickey A. Stroke warning campaigns: Delivering better patient outcomes? A systematic review. *Patient Relat Outcome Meas.* 2015;6:61–73.
87. Nishikawa T, Okamura T, Nakayama H, Miyamatsu N, Morimoto A, Toyoda K, et al. Effects of a public education campaign on the association between knowledge of early stroke symptoms and intention to call an ambulance at stroke onset: The Acquisition of Stroke Knowledge (ASK) Study. *J Epidemiol.* 2016;26:115–22.
88. Mc Sharry J, Baxter A, Wallace LM, Kenton A, Turner A, French DP. Delay in seeking medical help following transient ischemic attack (TIA) or “mini-stroke”: A qualitative study. Ozakinci G, editor. *PLoS One.* 2014;9:e104434.
89. Wilson A, Coleby D, Regen E, Phelps K, Windridge K, Willars J, et al. Service factors causing delay in specialist assessment for TIA and minor stroke: A qualitative study of GP and patient perspectives. *BMJ Open.* 2016;6:e011654.


# INNHOOLD

---

- |  |  |
|--|--|
| <b>01</b> Bevisstløs voksen            | <b>21</b> Hypotermi – hypertermi       |
| <b>02</b> Bevisstløst barn             | <b>22</b> Kjemikalier – gasser         |
| <b>03</b> Fremmedleg. i luftveiene     | <b>23</b> Krampeanfoll                 |
| <b>04</b> Katastrofe – stor ulykke     | <b>24</b> Krisehjelp                   |
| <b>05</b> Bestilt oppdrag              | <b>25</b> Magesmerter – ryggsmertter   |
| <b>06</b> Uavklart problem             | <b>26</b> Mulig dødsfall – krybbedød   |
| <b>07</b> Allergisk reaksjon           | <b>27</b> Neds. bevissthet – lammelser |
| <b>08</b> Blødning – ikke traumatisk   | <b>28</b> Psykiatri – suicid           |
| <b>09</b> Brannskade – elektrisk skade | <b>29</b> Pustevansker                 |
| <b>10</b> Brystsmertter – hjertesykdom | <b>30</b> Rus – forgiftning – overdose |
| <b>11</b> Diabetes                     | <b>31</b> Samarbeid med andre          |
| <b>12</b> Drukning                     | <b>32</b> Sykt barn                    |
| <b>13</b> Dykkerulykke                 | <b>33</b> Sår – brudd – småskader      |
| <b>14</b> Dyrebitt – insektstikk       | <b>34</b> Trafikkulykke                |
| <b>15</b> Feber                        | <b>35</b> Ulykker                      |
| <b>16</b> Forgiftning hos barn         | <b>36</b> Urinveier                    |
| <b>17</b> Fødsel                       | <b>37</b> Vold – mishandling           |
| <b>18</b> Gynekologi – svangerskap     | <b>38</b> Øre – nese – hals            |
| <b>19</b> Hodepine                     | <b>39</b> Øye                          |
| <b>20</b> Hud – utslett                | <b>40</b> Om Norsk indeks              |



# Start





# 27 Nedsatt bevissthet - lammelser

	KRITERIER	RÅD	RESPONS
	<p><b>Blødning:</b> Se 08 Blødning -ikke traumatisk.  <b>Sukkersyke:</b> Se 11 Diabetes.  <b>Krampeanfall:</b> Se 23 Krampeanfall.  <b>Rus eller overdose:</b> Se 30 Rus - overdose.  <b>Sykt barn:</b> Se 32 Sykt barn.  <b>Skader:</b> Se 35 Ulykker.</p>		
RØD - akutt	A.27.01 Reagerer ikke på tilrop og risting.	6	<p><b>LVS</b></p> <ol style="list-style-type: none"> <li>Hold forbindelsen.</li> <li>Alarmer ambulanse og LV-lege hvis ikke annet er forhåndsavtalt med AMK.</li> <li>Kople inn AMK og bistå med lokalkunnskap.</li> </ol> <p><b>AMK</b></p> <ol style="list-style-type: none"> <li>Alarmer ambulanse og LV-lege.</li> <li>Hvis hhnsiktsmessig, vurder også å                     <ul style="list-style-type: none"> <li>sende nærmeste amb. (selv med pas. om bord)</li> <li>varsle nærmeste lege (selv om ikke i vakt)</li> <li>varsle luftamb./legebil/spesialambulanse</li> <li>anbefale pasienttransport i privatbil</li> </ul> </li> <li>Gå gjennom resterende kriterier.</li> <li>Still relevante tillegsspørsmål.</li> <li>Gi relevante råd til innringer.</li> <li>Hold fortsatt forbindelsen (hvis mulig).</li> <li><b>Ved mistanke om slag:</b>  <b>Følg lokale prosedyrer for å rådspørre lege ved slagenhet om hastegrad.</b></li> <li>Overvåk aksjonen og oppdater alle (inkl. LVS)</li> <li>Tilby tips til helsepersonell.</li> </ol>
	A.27.02 Pustevansker.	1.4.5.10	
	A.27.03 Plutselig skjevhet i ansiktet.*	2.4.5	
	A.27.04 Plutselig nedsatt kraft i en arm eller en fot.*	2.4.5	
	A.27.05 Plutselig vansker med å snakke.*	2.4.5	
	A.27.06 Tiltagende uklar/sløv - mistanke om hjerneslag.*	1.4.5	
	A.27.07 Plutselig intens og uvanlig hodepine.	1.2.4.5	
	A.27.08 Fortsatt våken, men holder på å besvime.	1.3.4.5	
	A.27.09 Blek og klam.	1.3.4.5	
	A.27.10 Plutselig sterke magesmerter.	3.4.5.9	
	A.27.11 Plutselig hurtig puls og virker medtatt.	3.4.5	
	<p>* Hvis tegn og symptomer gir mistanke om hjerneslag, kan det være indikasjon for rask trombolyse selv flere timer etter symptomdebut. Hvis &lt; 4,5 timer siden symptomdebut: Still relevante tillegsspørsmål og kontakt straks vakthavende lege ved nærmeste slagenhet (eller annen avd. som tilbyr trombolytisk behandling) for råd om hastegrad.</p>		
GUL - haster	H.27.01 Har hatt krampeanfall, har kjent epilepsi og er fortsatt bevisstløs.	1.4.5.8	
	H.27.02 Har hatt krampeanfall, og er fortsatt like sløv/forvirret etter mer enn 20 min.	1.4.5.8	
	H.27.03 Plutselig uklar/sløv uten åpenbar forklaring.	1.3.4.5	
	H.27.04 Vedvarende sløv eller forvirret.	1.4.5	
	H.27.05 Plutselig lammelse som raskt har blitt bedre.	2.4.5	
	H.27.06 Mistet bevisstheten flere ganger siste døgn.	4.5	
	H.27.07 Plutselig synstap på det ene øyet.	2.4.5	
GRØNN - vanlig	V.27.01 Besvimte, men våken og OK nå.	3.5	
	V.27.02 Besvimte etter å ha tatt nitroglyserin, men er våken nå.	3.5.7	
	V.27.03 Har kjent epilepsi. Er i ferd med i ferd med å våkne etter et anfall.	1.5.8	
			<p>Lokal tilpasning</p> <p>Dato _____ Ansvarlig lege _____</p> <p><b>AMK / LVS</b></p> <ol style="list-style-type: none"> <li>Gå gjennom resterende kriterier.</li> <li>Still relevante tillegsspørsmål.</li> <li>Gi relevante råd til innringer.</li> <li>Gjør ett av følgende i hht. lokal instruks:                     <ul style="list-style-type: none"> <li>Sett innringer i kontakt med LV-lege.</li> <li>Informert LV-lege og kontakt innringer igjen.</li> <li>Send en ambulanse.</li> <li>Be pas. komme til legevakt/poliklinikk.</li> </ul> </li> <li>Be innringer ta ny kontakt straks hvis forverring.</li> </ol> <p><b>AMK</b></p> <ol style="list-style-type: none"> <li>Informert LVS.</li> </ol> <p><b>LVS</b></p> <ol style="list-style-type: none"> <li>Still relevante tillegsspørsmål.</li> <li>Gi relevante råd til innringer.</li> <li>Følg lokal instruks for videre håndtering, f.eks:                     <ul style="list-style-type: none"> <li>Informert LV-lege.</li> <li>Be innringer selv kontakte LV-lege på tlf. ....</li> <li>Be pas. komme til legevakten kl. ....</li> <li>Avtal ny telefonkontakt med innringer.</li> <li>Informert evt. hjemmesykepleien.</li> <li>Be pas. kontakte sin fastlege neste virkedag.</li> <li>Informert pasientens fastlege neste virkedag.</li> <li>Avslutt samtalen hvis opplagt ikke behov for hjelp.</li> </ul> </li> <li>Be innringer ta ny kontakt straks hvis forverring.</li> </ol> <p><b>AMK</b></p> <ol style="list-style-type: none"> <li>Gjør ett av følgende i hht. lokal instruks:                     <ul style="list-style-type: none"> <li>Formidle henvendelser til LVS.</li> <li>Be innringer selv kontakte LVS.</li> </ul> </li> </ol>





## Studie om medisinske operatørers bruk av "Norsk indeks for medisinsk nødhjelp" ved mottak av 113-samtaler

### Invitasjon til å delta i et forskningsprosjekt

Studien inngår i forskningsprosjektet "Norsk indeks for medisinsk nødhjelp - bruk og validering", et doktorgradsprosjekt ved Institutt for samfunnsmedisinske fag ved Universitetet i Bergen. Bakgrunnen for forskningsprosjektet er at Indeks nå har vært i bruk i 17 år og at den refereres til som en fasit for håndtering av medisinske nødanrop. Det har imidlertid aldri blitt undersøkt i hvilken grad Indeks faktisk hjelper den medisinske operatøren til å fatte en riktig beslutning på basis av primærmeldingen fra innringer, blant annet hvilken hastegrad som gjelder og hvilke ressurser som skal settes inn.

Før vi kan se nærmere på om Indeks fungerer må vi vite noe om hvordan og i hvilken grad den brukes. Indeksbruk ved AMK-sentraler er så langt lite undersøkt, men i en undersøkelse gjort av Nasjonalt kompetansesenter for prehospital akuttmedisin (NAKOS) i 2004 fant man ved hjelp av lydlogger at Indeks kun ble brukt i ca 64 % av henvendelser om bevisstløse pasienter, mens operatørene selv oppga bruken å være betraktelig høyere (99 %).

Invitasjonen til å delta i studien går ut til alle medisinske operatører ved AMK-sentralene i Norge. Du skal fylle ut et spørreskjema om din bruk av Indeks, og dette vil ta 15 – 20 minutter. Det er frivillig å delta, og du kan på hvilket som helst tidspunkt trekke deg fra studien. Vi spør ikke om direkte personidentifiserende opplysninger som navn eller personnummer, men noen faktorer som alder og kjønn kan være indirekte personidentifiserbare. Derfor vil kun jeg og min hovedveileder, Erik Zakariassen, ha tilgang til datamaterialet. Alle data blir lagt inn i og bearbejdet i et statistikkprogram, og ingen personer vil være mulig å identifisere ved presentasjon av funn. Doktorgradsprosjektet forventes å være avsluttet i løpet av 2014 og alle skjema vil da bli destruert.

Da vi ønsker din personlige besvarelse ber vi om at utfyllelse skjer uten samarbeid med kolleger eller leder. Skjemaet returneres oss i vedlagt ferdigfrankert konvolutt.

Denne undersøkelsen vil sammen med tre andre undersøkelser (AMIS-skjemaer, lydlogger og fokusgrupper) gi oss en god oversikt over hvordan og i hvilken grad Indeks brukes, og med det legge grunnlaget for del to av prosjektet som skal se nærmere på Indeks validitet. Resultatene vil bli publisert i internasjonale tidsskrift og gjort tilgjengelig for det nasjonale fagmiljøet. Prosjektet finansieres av Stiftelsen norsk luftambulans og Nasjonalt kompetansesenter for legevaktmedisin.

Undersøkelsen er godkjent av Regional etisk komité og tilrådd av Personvernombudet for forskning. Utfylling og innsending av skjemaet regnes som et samtykke til å delta.

På forhånd takk!

Med vennlig hilsen

Eirin Nybø Ellensen

Ph.d.- stipendiat

Institutt for samfunnsmedisinske fag, Universitetet i Bergen

E-post: [eirin.ellensen@isf.uib.no](mailto:eirin.ellensen@isf.uib.no),

Tlf.: 55 58 85 04 / 404 75 039





## Spørreundersøkelse om medisinske operatørers bruk av "Norsk indeks for medisinsk nødhjelp" ved mottak av 113-samtaler

### 1. Hvordan bruker du Indeks?

Ved mottak av en reell nødsamtale, angi hvor ofte du gjør følgende:	Aldri	Sjelden (<25 %)	Av og til	Ofte (> 75 %)	Alltid
Jeg slår opp i bordutgaven under samtalen					
Jeg slår opp på Start-kortet i bordutgaven					
Jeg bestemmer meg for hastegrad før jeg går inn på det aktuelle oppslaget					
Jeg stiller tilleggsspørsmål når jeg har avklart respons					
Jeg avklarer om pasienten er våken, dersom innringer og pasient ikke er samme person					
Jeg starter på toppen av det aktuelle oppslaget og jobber meg nedover					
Dersom jeg ikke bruker bordutgaven underveis, sjekker jeg etterpå at jeg har fått med meg alt					
Jeg gir råd til helsepersonell på stedet, hvis aktuelt					
Jeg setter kriteriekode uten å slå opp i Indeks					
Jeg spør om pasienten kan snakke, dersom innringer og pasient ikke er samme person					
Jeg gir råd til innringer eller pasient, hvis aktuelt					

I de tilfellene du ikke bruker bordutgaven, hva er grunnen til at du ikke gjør det? (du kan krysse av for flere alternativer)	
Det tar for lang tid	
Jeg er trygg på at jeg kan den	
Det er liten hjelp i Indeks	
Det blir tungvint å flytte hendene mellom tastatur og bordutgaven	
Det har blitt en vane å ikke slå opp	
Det forventes at jeg skal kunne Indeks	
Jeg er ikke opplært til det	
Jeg finner kriteriekodene i AMIS	
Annet?	

<b>I de tilfellene du ikke blar opp på Startkortet, hva er grunnen til at du ikke gjør det?</b> (du kan krysse av for flere alternativer)	
Det blir tungvint å bla opp flere ulike steder i bordutgaven	
Det tar for lang tid	
Jeg kan det	
Jeg foretrekker å gå rett til antatt aktuelt oppslag	
AMIS dekker startkortet	
Annet?	

<b>Angi i hvilken grad du er enig i følgende påstander:</b>	Svært uenig	Uenig	Litt uenig	Litt enig	Enig	Svært enig
Det er få som faktisk bruker Indeks						
Jeg blir sløv av å bruke Indeks i stedet for mitt eget hode						
Det fungerer bedre for meg å snakke fritt med innringer fremfor å følge Indeks						

<b>Vil følgende faktorer påvirke sannsynligheten for at du slår opp i bordutgaven?</b>	Øke	Senke	Ingen påvirkning	Ikke aktuelt
Sjelden problemstilling				
Alvorlig problemstilling				
Usikker hastegrad				
Kommunikasjonsproblemer med innringer (stress, språk, barn...)				
Uavklart problemstilling				
Dårlig magefølelse				
Aktuelt å varsle luftambulansen				
Aktuelt å varsle legevaktlegen				

**Hender det at du overstyrer Indeks i forhold til anbefalt respons?**

- Ja       Nei

<b>Dersom ja, i hvilke situasjoner skjer dette og hva gjør du?</b> (kryss bare av for det som passer)	Oppgraderer	Nedgraderer
Ambulansen er langt unna		
Kommunikasjonsproblemer med innringer (stress, språk, barn...)		
Dårlig magefølelse		
Gjengangere		
Annet?		

<b>Angi omtrentlig hvor ofte du bruker de enkelte oppslagene ved mottak av nødsamtaler (både bruk av bordutgave og bruk av kriterienummer i AMIS):</b>	Hver vakt	Ukentlig	Månedlig	Sjeldnere/aldri
01 Bevisstløs voksen				
02 Bevisstløst barn				
03 Fremmedlegeme i luftveiene				
04 Katastrofe - stor ulykke				
05 Bestilt oppdrag				
06 Uavklart problem				
07 Allergisk reaksjon				
08 Blødning - ikke traumatisk				
09 Brannskade				
10 Brystmerter - hjertesykdom				
11 Diabetes				
12 Drukning				
13 Dykkerulykke				
14 Dyrebitt - insektstikk				
15 Feber				
16 Forgiftning hos barn				
17 Fødsel				
18 Gynekologi - svangerskap				
19 Hodepine				
20 Hud - utslett				
21 Hypotermi - hypertermi				
22 Kjemikalier - gasser				
23 Krampeanfoll				
24 Krisehjelp				
25 Magesmerter - ryggmerter				
26 Mulig dødsfall - krybbedød				
27 Nedsatt bevissthet - lammelser				
28 Psykiatri - suicid				
29 Pustevansker				
30 Rus - forgiftning - overdose				
31 Samarbeid med andre				
32 Sykt barn				
33 Sår - brudd - småskader				
34 Trafikkulykke				
35 Ulykker				
36 Urinveier				
37 Vold - mishandling				
38 Øre - nese - hals				
39 Øye				

## 2. Alarmering av legevaktlege

Hvor ofte alarmerer du legevaktlegen? (ikke legevakten eller legevaktsentralen)	Aldri	Sjelden (< 25%)	Av og til	Ofte (> 75%)	Alltid
Ved grønn respons - via radio					
- via telefon					
Ved gul respons - via radio					
- via telefon					
Ved rød respons - via radio					
- via telefon					

Ved rød respons: angi i hvilken grad du er enig/uenig i følgende påstander:	Uenig	Litt uenig	Verken eller	Litt enig	Enig
Legevaktlegen skal alltid alarmeres ved rød respons					
Ambulansearbeiderne håndterer akutsituasjoner bedre enn legevaktlegen					
Jeg foretrekker luftambulanselege fremfor legevaktlege					
Legevaktlegene svarer uansett ikke på radio					
Legevaktlegene rykker uansett ikke ut					
Pasientene fraktes uansett til legevakten om det ikke er behov for akutt innleggelse					
Ambulansepersonellet klarer selv å vurdere om pasienten må innlegges eller ikke					
Legevaktlegen forsinker arbeidet					
Legevaktlegen går i veien for ambulansepersonellet					
Jeg alarmerer om ambulansen er langt unna					
Jeg vurderer alarmering på basis av hvor alvorlig jeg tror den røde responsen er					
Jeg alarmerer avhengig av hvilken lege som er på vakt og om alarmering ønskes					
Terskelen for å alarmere legevaktlegen er lavere ved traumer					
Terskelen for å alarmere legevaktlegen er høyere ved psykiatri					
Terskelen for å alarmere legevaktlegen er høyere ved hjerte- og lungeproblematikk					
Jeg alarmerer ikke til kommuner hvor jeg av erfaring vet at de ikke bruker radio					
Jeg alarmerer legevaktlegen fordi det er fast praksis ved min sentral					
Jeg alarmerer legevaktlegen i henhold til fastlagte lokale prosedyrer					
Jeg alarmerer til kommuner der det er avtale om å alarmere					
Jeg alarmerer legevaktlegen i henhold til Indeks					

### Hvordan foregår alarmering av legevaktlege ved rød respons vanligvis ved din sentral?

- Jeg alarmerer legevaktlegen
- Jeg avgjør om legevaktlegen skal alarmeres, men det er ressurskoordinator som alarmerer
- Ressurskoordinator tar seg alene av alarmeringen ut fra medlytt

### **3. Alarmering av luftambulanse (helikoptertjeneste) ved primæroppdrag**

**Er din AMK en LA-sentral (luftambulansesentral)?**

- Ja     Nei

**Dersom nei – hvordan foregår vanligvis kontakten med luftambulansesentralen:**

- Vi overlater hele oppdraget til LA-sentralen  
 LA-sentralen setter oss i forbindelse med luftambulansesentralen  
 Vi tar kontakt med luftambulansen selv

**Hvilke luftambulansesentraler forholder din sentral seg oftest til?**

- 1.
- 2.
- 3.
- 4.

**Kjenner du til om din AMK-sentral har en klart definert avtale med luftambulansesentralen/(-e) om hvilke hendelser de skal varsles på**

- Ja, skriftlige                       Ja, muntlige                       Nei

**Dersom ja, hvilke kriterier skal de varsles på?**

**Hvor ofte bestemmer du deg for å alarmere luftambulansen ved primæroppdrag?**

- Daglig (hver vakt)                       Ukentlig                       Månedlig  
 Årlig                       Sjeldnere/aldri gjort

**Når du alarmerer luftambulansen (enten direkte eller via LA-sentral) – hva er vanligste scenario:**

- De rykker umiddelbart ut gitt ok flyforhold
- Du og anestesilege diskuterer problemstillingen og tar en felles beslutning
- Anestesilegen avgjør alene hvorvidt de tar oppdraget eller ikke

**Dersom noen ber om luftambulanse, hender det du avviser forespørselen uten å snakke med vaktcrew?**

- Ja
- Nei

Hva er grunnen dersom du avviser forespørselen?	Ofte	Av og til	Sjelden	Aldri
Det er ikke flyvær				
Helikopteret er ute av drift				
De er opptatt med annet oppdrag				
Det er etter min vurdering ikke behov for luftambulanse				
Annet?				

**4. Din bakgrunn**

- Kvinne
- Mann

**Utdannelse:**

- Ambulansearbeider
- Sykepleier
- Annet: \_\_\_\_\_
- Evt videreutdanning: \_\_\_\_\_

**Arbeidserfaring:**

Hvor lenge har du arbeidet ved AMK? \_\_\_ år el. \_\_\_ mnd

Stillingsprosent ved AMK: \_\_\_ %



- Jobber du i rotasjon?  Nei
- Ja, i kombinasjon med ambulanse
- Ja, i kombinasjon med akuttmottak
- Ja, i kombinasjon med \_\_\_\_\_

Annen relevant bijobb (helsefaglig): \_\_\_\_\_ %

Tidligere relevant arbeid:  ambulansearbeider (utover utdannelsen)

sykepleier  annet: \_\_\_\_\_

## **5. AMK-sentral og Indeks**

### **Har du fått opplæring i bruk av Indeks?**

- Ja, i form av kurs da jeg begynte som operatør i AMK  Ja, blitt vist litt av kolleger
- Ja, i form av kurs etter ca \_\_\_\_\_ mnd  Nei

### **Repeteres bruk av Indeks for de som har jobbet ved AMK en stund?**

- Ja, regelmessig etter fastlagt plan Hvor ofte: \_\_\_\_\_
- Ja, med ujevne mellomrom
- Nei, ikke i det hele tatt

### **Synes du det er fokus på bruk av Indeks i ditt arbeidsmiljø?**

- Ja  Noe  Nei

## **6. Evt kommentarer?**

(Er det noe ved bruk av Indeks som vi ikke har fått frem ved dette spørreskjemaet? Noe du synes er viktig å få frem? Bruk evt baksiden ved behov.)

# NORSK INDEKS FOR MEDISINSK NØDHHJELP – BRUK AV INDEKS LYDLOGGPROSJEKT

Sentral: \_\_\_\_\_ AMIS: \_\_\_\_\_ Dato: \_\_\_\_\_ Klokken: \_\_\_\_\_  
 Hovedproblemi: \_\_\_\_\_  
 Innringer: \_\_\_\_\_ (pasient, helsepersonell...)  Tredjehåndsinopplysninger  
 Innringer samarbeider: Ja  Nei

## STARTKORT

HVOR – Hvor er pasienten/ulykken?

	Ja	Nei	Tid til avklart (</> 1 min)	Score
Adresse/stedsbeskrivelse verifisert?				
Tlf.nr verifisert?				

HVA – Hva er problemet?

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

VÅKEN – Er pas. våken og i stand til å snakke?

Spm. stillt/registerert	Ja	Åpenb./unødv	Nei	Tid til avklart ( 1 min)	Score
Er pasienten våken?					
Er pasienten i stand til å snakke?					
Reagerer på tilrop og risting?					
Alder?					
Kjønn?					
Navn?					
Full ID?					
Fastlege?					

## STARTKORT

TILSTAND/HASTEGRAD	RESPONS
	RÅD/VEILEDNING/INSTRUKSJON
TOTALT:	

## TILSTAND / HASTEGRAD

Satt kriterie \_\_\_\_\_ Satt hastegrad \_\_\_\_\_ Enig? \_\_\_\_\_

A	Ja	Åpenb./unødv	Nei	H	Ja	Åpenb./unødv	Nei	V	Ja	Åpenb./unødv	Nei
01				01				01			
02				02				02			
03				03				03			
04				04				04			
05				05				05			
06				06				06			
07				07				07			
08				08				08			
09				09				09			
10				10				10			
11				11				11			
12				12				12			
13				13				13			
14				14				14			

Vurdert kriterie \_\_\_\_\_

Vurdert hastegrad \_\_\_\_\_

## RESPONS

	Ja	Åpenb./unødv	Nei	Score	Kommentar
Tiltak iht hastegrad/kriterie?					
Legevaktlege alarmert?					

## RÅD/VEILEDNING/INSTRUKSJON

	Ja	Åpenb./unødv.	Nei	Score	Kommentar
Riktig å tilby RVI?					
Mulig å tilby RVI?					
RVI tilbudt?					
RVI korrekt?					
Instruksjon akseptert?					





ORIGINAL RESEARCH

Open Access

# Variations in contact patterns and dispatch guideline adherence between Norwegian emergency medical communication centres - a cross-sectional study

Eirin N Ellensen<sup>1,2\*</sup>, Steinar Hunskaar<sup>2,3</sup>, Torben Wisborg<sup>4,5</sup> and Erik Zakariassen<sup>1,2,3</sup>

## Abstract

**Background:** The 19 Norwegian Emergency medical communication centres (EMCCs) use Norwegian Index for medical emergency assistance (Index) as dispatch guidelines. Little is known about the use of Index, nor its validity. We aimed to document the epidemiology of contacts made to the public emergency medical phone number and the operators' self-reported use of Index as a first step towards a validation study.

**Methods:** We registered all medical emergency calls to the EMCCs during a 72 h period in a national cross sectional study. We subsequently sent a questionnaire to all EMCC operators in Norway, asking how they use Index. A combined outcome variable "use of Index" was computed through a Likert scale, range 1–5. Regression models were used to examine factors influencing use.

**Results:** 2 298 contacts were included. National contact rate was 56/1 000 inhabitants per year, range between EMCCs 34 – 119. Acute contact (life-threatening situations) rate was 21/1 000 per year, range between EMCCs 5 – 31. Index criteria 6 – 'Unresolved problem' accounts for 20% of the 113 contacts, range between EMCCs 10 – 42%. The mean use of Index was 3.95 (SD 0.39), corresponding to "more than 75% of emergency calls". There were differences in use of Index on EMCC level, range 3.7 – 4.4, and a multi regression model explained 23.4% of the variation in use. Operators working rotation with ground ambulance services reported reduced use of Index compared to operators not working in rotation, while distinct EMCC focus on Index increased use of Index compared to EMCCs with no focus on Index. Use of electronic records and operators experience were the main reasons given for not using Index.

**Conclusions:** There is a large variation between the EMCCs with regard to both contact patterns and use of Index. There is a relatively high overall self-reported use of Index by the operators, with variations on both individual and EMCC level.

**Keywords:** Criteria based dispatch, Emergency medical dispatch, Emergency medical services, Emergency medical communication systems, Triage, Guideline adherence

\* Correspondence: eirin.nybo.ellensen@norskluftambulanse.no

<sup>1</sup>Department of Research, Norwegian Air Ambulance Foundation, PB 94, Drøbak 1441, Norway

<sup>2</sup>Department of Global Public Health and Primary Care, University of Bergen, 5050, Bergen, PB 7800, Norway

Full list of author information is available at the end of the article



## Introduction

Norway has a dedicated medical emergency phone number 113, which is answered and handled by an operator at the nearest Emergency medical communication centre (EMCC). The EMCCs dispatch the ambulance fleet, consisting of cars, boats, helicopters and planes, and facilitates radio and telephone communication amongst all the different participants in the chain of pre-hospital emergency medical care, from ambulance personnel through general practitioners on call to in-hospital specialists, and with other emergency authorities like police and fire brigades [1].

Norwegian Index for Medical Emergency Assistance (Index) [2] is the dispatch guidelines used by the EMCCs. It was developed from the Criteria Based Dispatch (CBD) guidelines in King County, Washington, USA [3], and introduced in Norway in 1994. Today Index is implemented as the only dispatch tool in all 19 EMCCs, serving an important role in securing equal high quality of the emergency health communication system throughout Norway [2,4,5], as well as assisting the individual operator when handling an emergency call. Despite its importance on both the individual level (the patient in acute need of medical assistance) and on the systemic level (adequate resource allocation), Index has not been evaluated or validated during its nearly twenty years in use.

The CBD guidelines, and thus the Index, use symptom criteria to determine the urgency of the medical condition, and the appropriate level of response. This allows for a dynamic approach to the dispatch situation, compared to the more strict protocol compliance required by algorithm based dispatch systems. CBD and Index hence require an operator with a certain level of medical education and experience. In Norway, this is normally a registered nurse with experience from an emergency ward or an intensive care unit. The interpretation and modification of the formal guidelines according to individual skills will influence the criteria set, introduce an unknown variance in guideline adherence and hence lower the validity of any study regarding dispatch [6]. Guideline adherence is defined as "conformity in fulfilling or following official, recognized, or institutional requirements, guidelines, recommendations, protocols, pathways, or other standards" [7]. The algorithm based dispatch systems, such as Medical Priority Dispatch System (MPDS), traditionally use the term 'compliance' when referring to protocol adherence [6,8,9]. MPDS monitor compliance frequently along with many other quality markers [9], facilitating validity studies.

To our knowledge there are no international studies addressing CBD adherence. A recent systematic review on adherence to guidelines and protocols in the pre-hospital setting was not able to identify any eligible

study in the emergency medical dispatch setting, neither CBD nor MPDS based [10]. However, a Norwegian study from 2005 evaluated dispatch in drug-related emergencies, and found an Index adherence of 99% based on multiple choice questionnaire, but only 64% guideline adherence based on log recordings [11].

A study on the epidemiology of medical emergency contacts made to three Norwegian EMCCs during a three month period in 2007 [12], focused on acute life threatening situations coded as acute only. The study showed statistically significant differences in the acute response rates among the EMCCs, but gave no explanation with regards to whether these differences could indicate variations in EMCC triaging or differences in the populations. National data on different urgency levels, use of Index criteria codes and contact rates to explore possible variations on a national level or study the triaging component further was not available in 2011 when we started the present study, but is now available as an annual report [13].

Before starting a validation study we needed to document the epidemiology of the contacts made to 113 and the emergency medical dispatch guideline adherence. The first aim of this study was to document possible differences in urgency levels, Index criteria and contact rates among the EMCCs. The second aim was to document the operators' self-reported use of Index by a questionnaire, and determine factors influencing the use.

## Methods and materials

### Setting

Autumn 2011 there were 19 EMCCs in Norway, belonging to four regional health trusts. The EMCCs differ in size, covering from 65 000 to 1 165 000 inhabitants, with nearly 5 million inhabitants in total [14,15]. They are traditionally manned by registered nurse operators who answer and handle the 113 calls, and ambulance educated resource coordinators who dispatch and coordinate the ambulance fleet.

In addition to using the same dispatch guideline to prioritize and handle contacts [2], all EMCCs use the same software program Acute medical information system (AMIS) to register information and document each contact. AMIS contains information on Index criteria used, urgency level, and response dispatched.

### 113 Epidemiology – cross sectional population based survey

All 19 EMCCs contributed with data from every medical emergency contact made to the emergency medical phone number 113, during a 72 hour set period in August 2011. The data were collected in form of AMIS printouts, and contained information about date, time,

the caller role, the patients sex and age, Index criteria used, urgency, response, and involvement of ambulances, air ambulances, GPs and others for each contact. Only calls registered to the 113 telephone line was included, excluding calls made by direct lines from fire brigades, police and Local emergency communication centers (LEMCS). Duplicates, maculated incidents and misdialling were also excluded (Figure 1).

#### Norwegian index for medical emergency assistance (index)

Index is a flip-over in large format consisting of a Start page, 36 symptom cards and 4 administrative cards. The Start page is built like an algorithm and clarifies important information as location, phone number, vital functions and a brief questioning on the patient's problem. The operator then moves on to the proper symptom card, determined by the problem presented. The symptom cards display a list of different criteria in decreasing severity, from the acute symptoms at the top and all the way down to the non-urgent symptoms at the bottom. The operator starts at the top and moves downwards through specific criteria, until one criterion is finally met. This criterion then determines the response dispatched. Index divides the response into three categories; red, yellow and green. Red response is when the situation is acute and life threatening, and calls for immediate "blue lights and sirens" ambulance dispatch and GP on-call alarm. Yellow response is an urgent and potentially life-threatening situation where ambulance and GP alarm is constantly assessed. Green responses are non-urgent situations, where the patients often are referred to the primary level of emergency care, unless there is an obvious need of ambulance transport.

#### Self-reported use of index – questionnaire based survey

There were no validated questionnaires available to address our aim, so we developed a questionnaire in cooperation with one of the EMCCs and tested it on operators there to avoid possible misunderstandings.

Questions on their use of Index were arranged as a Likert scale of five symmetric response formats; never, seldom (<25%), sometimes, often (>75%) and always. The eleven different questions were all expressed in the same form: "During a real emergency call, how often do you...?" and covered different aspects of Index (Table 1). In addition to asking directly about use of Start page, we also asked specific questions from the Start page, like "... clarify if the patient is awake?" and "... if the patient can talk?" to unmask actual use of Start page and not just flipping of the page. The operators were asked about their educational level, EMCC work experience, initial training and later repetitions with regards to use of Index. We also asked if they thought there was focus on use of Index at their EMCC. The operators were given the possibility to add explanations or points of view at the end of the questionnaire. The questionnaire was distributed in paper through EMCC management, but returned directly to the research facility to secure anonymity for the individual operator.

#### Population

There were 429 persons working as operators in the EMCCs when our study started. An operator was defined as the person for whom the primary occupation is to answer the emergency medical phone call. A resource coordinator was defined as a person for whom the primary occupation is to coordinate the ambulance fleet.




Figure 1 Excluded AMIS forms.

**Table 1 Questions used to determine use of Index**

	N	Mean	SD
<b>When receiving a true emergency medical call, please specify how often you do the following:</b>			
I use the table format Index during the conversation	270	4.02	0.82
I use Start page	272	3.36	1.23
I clarify if the patient is awake	270	4.93	0.26
I clarify whether the patient can talk or not	270	4.32	0.73
I do not decide upon urgency before criteria (reversed in questionnaire)	272	3.02	0.87
I do not determine criteria code without using Index (reversed in questionnaire)	269	3.80	0.94
I start on top of the symptom card and work my way downwards	267	4.01	0.86
I ask further questions	269	4.09	0.74
I give advices to the caller/patient, if needed	270	4.56	0.56
I give advices to health personnel on scene, if needed	268	3.70	1.00
If Index is not used during the call, I check afterwards whether I have covered everything	263	3.63	1.02
Total (Likert score)		3.95	0.39

1 = Never, 2 = Seldom (<25%), 3 = Sometimes, 4 = Often (>75%), 5 = Always. N = 272 operators.

Only questionnaires from operators were included in the study.

#### Statistical analysis and ethical approvals

Continuous data are presented as mean (SD) or median (quartiles) for symmetric and skewed data, respectively. Rates are calculated as contacts per 1 000 inhabitants per year. The five response formats on the Likert scale were valued 1 to 5, and the outcome variable “use of Index” was calculated for each operator as the mean score from all eleven questions constituting the Likert scale. Q-Q plots were used to check the outcome variable for normal distribution. We explored the variation in mean use of Index with respect to each of the explanatory variables by one-way analysis of variance (ANOVA) and univariate linear regression. All explanatory variables were then entered in a full multiple linear regression model. Effect estimates are reported with 95% confidence intervals (CI). A possible hierarchical effect of EMCC central was assessed by adjusting for central in a mixed model. There were only marginal differences in effect estimates when adjusting for the mixed effect, and we therefore consequently report results from the ANOVA and linear regression models.

The statistical analyses were performed using Statistical Package for the Social Sciences (IBM SPSS Statistics 20) and STATA (Stata/IC 12.1). P-values below 0.05 were considered statistically significant.

Both studies were evaluated by the Regional committees for medical and health research ethics and considered not to be in need of ethical assessment. Exemption for consent was given by the Regional committees for medical and health research ethics in terms of receiving and analyzing sensitive data. The Data Protection Official for Research approved the study.

## Results

### Epidemiology

We collected 3 294 AMIS forms during a 72 hour period, of which 2 298 were included for analyses. This gives a calculated overall national emergency medical contact rate of 56/1 000 inhabitants per year (95% CI 54–59), ranging from 34 (95% CI 22–46) to 119 (95% CI 91–146) among the different EMCCs. Of the 2 298 contacts 37% were assessed to be acute, 34% urgent, and 27% non-urgent, while 1.8% of the contacts had no urgency assessment. Looking at acute contacts separately, the rates varied significantly from 5 to 31/1 000 inhabitants per year (Table 2).

The most frequently used Index criteria was “6 - Unresolved issue” (Table 3), which accounted for 20% of all contacts. The use of this criteria differed from 10% at one EMCC to 42% at another. The use of other criteria like “5 - Ordered assignment” and “28 - Psychiatry - Suicide” also varied much between different EMCCs, from 0 to 22% and 19% respectively (Table 3). The urgency assessments on criteria 6 differed between the EMCCs; acute 0 - 53%, urgent 0 - 67% and non-urgent 9 - 86% (Figure 2).

### Use of Index

We received 275 questionnaires after three reminders, of which 272 questionnaires were included in the study, representing all 19 EMCCs (response rate 63.4%). The operators were mainly female, registered nurses, of whom many worked in rotation at either emergency rooms (ER) or ambulances (Table 4). The majority received training in use of Index when starting to work at the EMCC, but later repetitions were rare (Table 4). The EMCC work experience ranged from 1 month to 36 years, with a median (quartile) time of 6 (3, 12) years.



**Table 2 Acute contact rates between different Emergency medical communication centres**

EMCC	Acute contacts		95% CI
	N	Rate	
1	19	31	(17–46)
2	14	14	(7–22)
3	11	21	(8–33)
4	13	12	(5–18)
5	3	5	(0–10)
6	17	15	(8–22)
7	46	19	(13–24)
8	12	13	(6–21)
9	18	15	(8–22)
10	12	14	(6–21)
11	87	25	(20–31)
12	23	16	(10–23)
13	34	12	(8–16)
14	56	24	(17–30)
15	78	26	(20–32)
16	38	15	(11–20)
17	242	25	(22–28)
18	76	24	(19–30)
19	42	18	(13–24)
<b>Total</b>	841	21	(19–22)

Contacts made to all the EMCCs, and assessed by the operators to be of the most urgent of three urgency categories. N = number of acute contacts during a data collection period of 72 h. Rates calculated per 1 000 inhabitants per year.

Use of Index ranged from 2.91 to 4.82, with overall mean (SD) for all operators being 3.95 (0.39), corresponding closely to the response format “often, > 75%” (=4) in the questionnaire. The individual questions on use of Index ranged from 4.93 (0.26) for “I clarify if the patient is awake” to 3.02 (0.87) for “I do not decide upon urgency before criteria” (Table 1). ANOVA showed significant differences in use of Index among the EMCCs ( $p < 0.001$ ), with the EMCCs ranging from 3.7 (0.24) to 4.4 (0.39). Figure 3 shows the spread between the EMCCs. Possible explanatory variables for use of Index were analyzed by univariate and multiple linear regression models (Table 5). Time, the effect of rotation with ambulance and a clear focus on use of Index at workplace on use of Index were statistically significant, with focus being by far the strongest component ( $\beta = 0.46, p < 0.001$ ) (Table 5).

For use of Start page separately, we found that 47% reported using the Start page often or always (> 75% of real emergency calls), while 93% always checked if the patient was awake and 46% if he could talk. The reasons given for not using Index are shown in Table 6.

The question “work percentage at EMCC” was misinterpreted by many operators with regard to whether

include rotatory work at ER/ambulances or not, and was hence excluded from analysis. Explanations offered in free text did not add any relevant information on the questions addressed, and were not analyzed.

## Discussion

The main finding of the epidemiological part of the study was the wide spread between the different EMCCs, regarding both overall contact rates, specific acute rates and use of different Index criteria when assessing the situations. The questionnaire study showed a relatively high overall use of Index reported by the operators. It also showed large unexplained variations in the use of Start page and several symptom cards.

## Strengths and weaknesses of the studies

The epidemiology study brings new knowledge about contacts made to 113, while the self-reported use of Index represents the first step towards determining use of Index. Both studies include all 19 EMCCs and hence are representative on a national level.

The short study period of 72 h limits the validity of extrapolation, and was mainly due to capacity issues in both ends, as data had to be exchanged in paper format at the time being. The questionnaire was not validated, as we wanted the main study population as large as possible, in terms of both individuals and centers.

Self-reporting allows for over- and underestimation. This was a calculated risk as we aimed to document how the operators themselves thought they used Index, and the results must be interpreted as subjective. We will address the problem with subjective contra objective use of Index in future audio-log study. The response rate of 63.4% could represent a selection bias, but the large range in ‘use of Index’ score, both on an individual level and on EMCC level, together with all EMCCs being represented, indicates that we have a representative material. The response rate can to some extent be explained by all information, distribution and reminders depending on the management at each EMCC. The official policy by the EMCC managements on always using Index, combined with the questionnaires and reminders being provided by the management, raises the possibilities of an “eagerness to please” bias. This was attempted opposed by ensuring anonymity and allowing the filled out questionnaires to return directly to the main investigator, through prepaid preaddressed envelopes following each single questionnaire. Recall biases are likely to occur when trying to recall what one usually does in certain situations. The use of a non-validated questionnaire led to misinterpretation of some questions. Never and always are narrow categories, and the results on use of Index might have been more correct if the categories had been more evenly distributed.

**Table 3 Top ten Index criteria**


EMCC	Contacts N	Criteria number*											Missing %	Total %
		6 %	10 %	5 %	30 %	33 %	29 %	35 %	25 %	28 %	27 %	All other %		
1	72	14	3	3	6	4	6	4	4	12	9	13	25	100
2	44	20	9	5	0	9	7	5	5	5	5	20	11	100
3	40	22	8	8	0	14	0	11	8	8	3	8	18	100
4	56	32	11	4	11	11	5	5	2	2	2	14	2	100
5	32	27	13	0	3	7	7	0	0	13	3	19	13	100
6	53	19	6	6	6	2	6	4	4	19	4	19	8	100
7	89	26	6	7	2	4	6	11	11	0	4	22	0	100
8	46	15	7	11	4	2	9	4	9	15	9	9	7	100
9	67	22	5	3	2	2	0	5	5	10	3	15	33	100
10	30	30	10	10	10	7	3	0	3	0	10	17	0	100
11	240	10	10	4	10	5	4	4	3	1	4	15	30	100
12	65	25	2	6	11	10	3	5	5	10	0	25	2	100
13	117	30	13	2	11	7	10	4	4	4	6	12	1	100
14	114	14	7	8	4	4	4	4	7	5	5	20	18	100
15	168	23	14	8	5	5	7	7	7	4	7	13	1	100
16	119	31	8	10	5	5	3	6	4	6	4	18	0	100
17	605	14	10	4	6	6	7	5	5	3	5	18	19	100
18	228	23	10	22	4	4	4	6	4	5	0	18	0	100
19	113	42	10	4	3	7	5	3	6	6	3	12	0	100
Total	2298	20	9	7	6	6	5	5	5	5	5	16	12	100

The most frequently used criteria in Index (%), for each of the 19 EMCC. All degrees of urgency included. N = total number of contacts during the study period of 72 h.  
 \*Criteria number explanation: 5 – Commissioned assignments (by health personnel), 6 – Unresolved issue, 10 – Chest pain, heart disease, 25 – Stomach pain, back pain, 27 – Reduced consciousness, paralyses, 28 – Psychiatry, suicide, 29 – Breathing issues, 30 – Intoxication, overdose, 33 – Wounds, fractures, minor injuries, 35 – Accidents.

**Epidemiology**

There is a wide spread in annual contact rates among the EMCCs, from 34 to 119/1 000 inhabitants. We have no national data on acute disease distribution to compare with, but one would not expect a similar spread in

distribution. Accidents are shown to disperse a geographical pattern, a study from 2013 comparing a rural county with an urban/rural county found a higher accident rate in urban areas, but a higher mortality rate per accident in the rural areas [16]. This dispersion of accidents



**Figure 2 Use of Index criteria “6 – Unresolved issue” between the EMCCs.** Differences in urgency assessment, and hence use of Index criteria «6 – Unresolved issue» between the different EMCCs. Red = acute, yellow = urgent, green = non-urgent contacts.

**Table 4 Operator characteristics**

		N	%
Sex	Female	178	65
	Male	85	32
	Missing	9	3
		272	100
Education	Ambulance worker	50	18
	Registered nurse	230	85
	Ambulance worker and nurse	16	69
	Other	11	4
	Further education	86	32
Work rotation	None	73	27
	With emergency room	116	43
	With ambulance	44	16
	Other	19	7
	Missing	20	7
		272	100
Work experience	≤ 1 year	27	10
	>1 ≤ 5 years	95	35
	>5 ≤ 10 years	60	22
	> 10 years	78	29
	Missing	12	4
		272	100
Training in use of Index	Formal training from the start	234	86
	Formal training eventually	16	6
	Taught by colleagues	8	3
	None	2	1
	Missing	12	4
		272	100
Repetition of use of Index	Regularly	51	19
	Irregularly	129	47
	None	71	26
	Missing	21	8
		272	100

Background information on the operators' (N = 272) educational and work experience, together with experienced initial training in use of Index and later repetitions.

does not account for the wide spread in contact rates, and this could indicate that the population uses 113 differently depending on location. A previous study of geographic variations in alerting, dispatch and response found that while severity of illness/trauma had no effect on the use of 113, the use was lower in rural areas compared with urban areas [17]. This could be due to local differences in organisation of LEMCs and casualty clinics, and cultural differences for when to access the different levels of emergency health services.

When separating contacts assessed to be acute, we also find a large spread in rates between the different EMCCs. The overall acute contact rate of 21 is slightly lower than reported by a previous study based on data from three EMCCs. They found a three month acute rate of 6.2/1 000 inhabitants in 2007, giving an annual rate of 25/1 000 [18]. As our study shows that the EMCCs differ with regards to both urgency distributions and contact rates, differences in inclusion of EMCCs will affect these outcomes.


A recently published study from Denmark studied Danish Index' ability to triage patients according to severity [19], following the implementation of the CBD guidelines in Denmark. They found a national acute (emergency level A) rate of 17 ambulance dispatches/1 000 inhabitants per year, which is somewhat lower than our 21 acute contacts/1 000 inhabitants per year. There were also differences between their included areas: with 13, 17 and 21 acute ambulance dispatches/1 000 per year. One possible explanation could be the large differences in EMCC population size, with five centres in Denmark and 19 in Norway covering 5.5 and 5 million inhabitants respectively. This difference could influence use of Index, and hence incidence of acute criteria codes. Cultural differences in what emergency medical level to access and differences in pre-hospital emergency medicine organization could influence this.

On primary health care level, a study of 85 000 contacts from 2007 investigated the distribution of urgency levels in the Norwegian emergency primary healthcare services: acute 2.3%, urgent 21.1% and non-urgent 76.6% [20]. Compared to our findings (acute 37%, urgent 35%, non-urgent 26%), these differences show that the population as a whole know what level to address, depending on degree of medical emergency.

The most frequently used Index criteria, all urgencies included, was "6 - Unresolved issue". This criterion covers a whole range of situations; from the unclear situations where the operator gets too little information to choose another criterion to well-defined situations where no other criteria match. The large differences between the EMCCs, both in terms of total use of this criterion and the variation in urgencies as it was used in, clearly indicates a variation in use of Index from center to center.

#### Use of Index

The operators reported a relatively high overall use of Index, corresponding to "in over 75% of real emergency calls". The variation among the operators was quite large though. Although we found some factors associated with positive and negative effects on use of Index, we were only able to explain 23.4% of this variation, indicating that there are factors influencing use of Index that we



**Figure 3 Use of Index between the EMCCs.** Box plot of mean Likert-score values, displaying use of Index between the different EMCCs. The box represent the first, second (median) and third quartile of the data. The whiskers show the lowest datum within 1.5 interquartile range of the first quartile, and the highest datum within 1.5 interquartile range of the third quartile. Outliers are shown as dots.

were not able to uncover. The main reason given for not using Index was AMIS, and the confidence that this software program provided the necessary key words to properly assess the acute medical emergency situation. The operators' background was to great extent as expected. It was an experienced group of employees, with median working time at the EMCC of 6 years. The results indicated decent routines for training new operators in use of Index, but not for repetition.

Our result of approximately 75% use of Index lies between the divergent findings from the 2005 study, with 99% self-reported Index guideline adherence and 64% adherence based on log-recordings [11]. To our knowledge there are no other studies addressing adherence to criteria based guidelines. As algorithm based dispatch systems require a much stricter adherence to protocol they have systems developed to monitor and increase protocol compliance, among other quality markers [6,8,21-24].

**Table 5 Explanatory variables for the variation in use of Index**

	Univariate linear regression			Multiple linear regression		
	Estimate	(95% CI)	P-value	Estimate	(95% CI)	P-value
Working experience	-0.011	(-0.018, -0.003)	0.004	-0.012	(-0.019, -0.004)	0.002
Rotation						
With ground ambulance	-0.218	(-0.360, -0.076)	0.003	-0.213	(-0.348, -0.078)	0.002
With emergency room	0.054	(-0.057, 0.165)	0.337	0.033	(-0.074, 0.139)	0.545
With others	-0.093	(-0.283, 0.098)	0.339	-0.046	(-0.229, 0.137)	0.621
Repetition of use of Index						
Regularly	0.214	(0.074, 0.354)	0.003	0.062	(-0.080, 0.205)	0.390
Irregularly	0.090	(-0.023, 0.203)	0.118	-0.002	(-0.116, 0.111)	0.969
Focus on use of Index						
Clear focus	0.522	(0.312, 0.732)	<0.001	0.459	(0.240, 0.679)	<0.001
Some focus	0.293	(0.073, 0.512)	0.009	0.207	(-0.013, 0.426)	0.065

Exploration of possible explanations for the variation in use of Index, by univariate and multiple linear regression models. Working experience = effect per year. P-value < 0.05 was regarded as statistically significant. Explained variation, R<sup>2</sup> = 0.234 for the multiple linear regression model.

**Table 6 Reasons for not using Index or Start page**

	N	%
<b>Question: When not using the Index flip over, why?</b>		
I find the criteria in AMIS	149	55
I know it	91	33
It takes too long	55	20
It is cumbersome to move my hands between Index and the keyboard	44	16
It has become a habit	24	9
I am expected to know it	23	8
It is of little help	13	5
I am not trained to use it	0	0
<b>Question: When not using the Start page, why?</b>		
I know it	117	43
I prefer to go straight to proper card	108	40
I find what I need in AMIS	57	21
It is cumbersome to move the hands between flip over and keyboard	38	14
It takes too long	27	10

Given reasons for not using Index/Start page, checked off by the operators.  
 Total number of operators = 272.

Although compliance is highlighted as such an important feature of MPDS, less than 3% of the registered users are Accredited Centers of Excellence [25], which among other quality measurements include a minimum of 90–95% compliance with different parts of the protocol [8].

Given the purpose of CBD guidelines, to assist health care personnel in decision making rather than defining specific questions or actions to be taken [3], use of Index was not expected to reach full score. Looking at the different explanatory variables though, it was rather unexpected that time working at an EMCC had so little effect on use of Index. It was equally unexpected that the effect of what EMCC the operator works at was so negligible, as we held this to be the natural explanation for the variations in use of Index.

Working in rotation with ground ambulance is the variable strongest associated with decreased use of Index. This is not associated with educational level, as half the operators rotating with ambulance are registered nurses and there is no difference in use of Index between the different educational levels among those rotating with ambulance. Previous research has found that individual experience and professional autonomy affects guideline adherence [10]. Our study finds no effect on use of Index for operators rotating with ER. A Dutch study on protocol adherence among emergency and ambulance nurses found the opposite result; ambulance nurses are more likely to hold protocol over experience while emergency nurses are more likely to hold experience over protocol [26].

The low use of Start page is reflected in the reasons given for not using it. They know it, and find the key words they need in AMIS screen. The Start page (3<sup>rd</sup> edition) includes the phrase: “Is the patient awake and can talk?” [2]. The low percentage reporting to actually ask about this latter, confirms that neither AMIS nor memory equals Start page, and one may speculate that the reason for including this question is not fully understood among those using the tool.

The main reason for not using Index appears to be that the operators choose to use the AMIS screen above Index, and a belief that AMIS provides the necessary keys to assist them in properly assessing the situation. This might be of some concern, since AMIS does not supply any support in neither decision making nor advices for the public or health personnel at scene, and hence cannot replace Index. Furthermore, discarding the guidelines in favor of memory and own experience is a potential hazard of losing vital information or getting sidetracked due to unstructured interrogation. This risk is strongly advocated by critics of criteria based dispatch [9,22].

Based on our findings we would recommend increased focus on use of Index at each EMCC. The systematic review on guideline and protocol adherence in the pre-hospital and emergency care setting [10] finds that tailored strategies towards identified barriers improve professional practice, as well as strategies aimed at influencing factors improve guideline adherence. Evidence based recommendations and a relationship between guideline adherence and clinical outcome are also mentioned as important motivational factors for guideline adherence.

In this study we estimated a mean value for use of Index among operators in Norway, based on self-report, which obviously represents a subjective view. Investigating the real practice based on objective data is therefore a natural next step in documenting use of Index.

## Conclusion

The EMCCs varies regarding both contact rates and use of Index. The operators report use of Index in > 75% of real emergency situations, but there are large individual differences. Clear focus on use of Index at the EMCC was the strongest predictor for increased use, and work rotation with ground ambulance services was associated with decreased use.

## Abbreviations

AMIS: Acute medical information system; ANOVA: Analysis of variance; CBD: Criteria based dispatch; CI: Confidence interval; EMCC: Emergency medical communication centre; ER: Emergency room; GP: General practitioners; LEMC: Local emergency medical communication centre; MPDS: Medical priority dispatch system; SD: Standard deviation; SPSS: Statistical Package for the Social Sciences.

## Competing interests

The authors declare no competing interests. The funding foundation had no participation in conducting the study or writing the manuscript.

#### Authors' contributions

All authors contributed in designing the studies. ENE collected and analysed the data, and drafted the manuscript. All authors contributed substantially in the interpretation of the findings and rewriting the manuscript. All authors read and approved this final version.

#### Acknowledgements

We would like to thank Jo Røislien at Norwegian Air Ambulance Foundation for excellent help on statistical issues. We also want to thank operators and management at all the 19 EMCCs in Norway for participating in the studies, and Tone Alfsvåg Engelsen at Bergen AMK for contributing in the planning process. The project was supported financially by the Norwegian ExtraFoundation for Health and Rehabilitation through EXTRA funds. Last but not least, we would like to thank all members of the Norwegian Air Ambulance Foundation for the financial support that makes this research possible.

#### Author details

<sup>1</sup>Department of Research, Norwegian Air Ambulance Foundation, P.O. Box 94, N-1441, Drøbak, Norway. <sup>2</sup>Department of Global Public Health and Primary Care, University of Bergen, P.O. Box 7804, N-5020 Bergen, Norway. <sup>3</sup>National Centre for Emergency Primary Health Care, Uni Health, Uni Research, Kalfarv. 31, N-5018 Bergen, Norway. <sup>4</sup>Anaesthesia and Critical Care Research Group, Faculty of Health Sciences, University of Tromsø, N-9037 Tromsø, Norway. <sup>5</sup>Norwegian Trauma Competency Service, Oslo University Hospital, N-0450 Oslo, Norway.

Received: 22 August 2013 Accepted: 31 December 2013

Published: 8 January 2014

#### References

1. Norwegian Ministry of Health and Care services: *Forskrift Om Krav til Akuttmedisinske Tjenester Utenfor sykehus*. (Regulation on pre-hospital emergency medicine services) [http://www.lovdata.no/for/si/ho/so-20050318-0252.html]
2. Norwegian Medical Association: *Norsk indeks for Medisinsk Nødhjelp (Noregian Index for Medical Emergency Assistance)*. 3rd edition. Stavanger: Laerdal Medical A/S; 2009.
3. Cully LL, Henwood DK, Clark JJ, Eisenberg MS, Horton C: **Increasing the efficiency of emergency medical services by using criteria based dispatch**. *Ann Emerg Med* 1994, **24**:867–872.
4. Norwegian Medical Association: *Medisinsk Nødmeldetjeneste, Standard for Prehospital Akuttmedisin*. (Emergency health communication system) [http://www.kokom.no/hurtigvalg/nodmeldetjenesten.htm]
5. Norwegian Ministry of Health and Care Services: *Stortingsmelding 43 (1999–2000) Om Akuttmedisinsk Beredskap*. (About emergency preparedness) [http://www.regjeringen.no/Rpub/STM/19992000/043/PDFA/STM19992000043000DDPDF.pdf]
6. Clawson JJ, Cady GA, Martin RL, Sinclair R: **Effect of a comprehensive quality management process on compliance with protocol in an emergency medical dispatch center**. *Ann Emerg Med* 1998, **32**:578–584.
7. The National Centre for Biotechnology Information (NCBI): *Medical Subject Headings (MeSH)*. [http://www.ncbi.nlm.nih.gov/mesh/?term=guideline+adherence]
8. International Academies of Emergency Dispatch: *Accreditation*. [http://www.emergencydispatch.org/Accred20PointsMedical]
9. Clawson JJ, Martin RL, Hauert SA: **Protocols vs. guidelines. Choosing a medical-dispatch program**. *Emerg Med Serv* 1994, **23**:52–60.
10. Ebben RH, Vloet LC, Verhofstad MH, Meijer S, Mintjes-de Groot JA, van Achterberg T: **Adherence to guidelines and protocols in the prehospital and emergency care setting: a systematic review**. *Scand J Trauma Resusc Emerg Med* 2013, **21**:9.
11. Lorem T, Saether E, Wik L: **Handling of drug-related emergencies: an evaluation of emergency medical dispatch**. *Eur J Emerg Med* 2009, **16**:37–42.
12. Zakariassen E, Burman RA, Hunskaar S: **The epidemiology of medical emergency contacts outside hospitals in Norway—a prospective population based study**. *Scand J Trauma Resusc Emerg Med* 2010, **18**:9.
13. The Norwegian Directorate of Health: *Nasjonale data Fra Akuttmedisinske Kommunikasjonssentraler. Årsrapport 2012*. (National data from Emergency Medical Communication Centers 2012) [http://helseidirektoratet.no/

- publikasjoner/nasjonale-data-fra-akuttmedisinske-kommunikasjonssentraler-arsrapport-2012]
14. *Statistics Norway*. [www.ssb.no]
  15. *National Centre on Emergency Communication in Health*. [www.kokom.no]
  16. Bakke HK, Hansen IS, Bendixen AB, Morild I, Lilleg PK, Wisborg T: **Fatal injury as a function of rurality – a tale of two Norwegian counties**. *Scand J Trauma Resusc Emerg Med* 2013, **21**:14.
  17. Vaardal B, Lossius HM, Steen PA, Johnsen R: **Have the implementation of a new specialized emergency medical service influenced the pattern of general practitioners involvement in pre-hospital medical emergencies? A study of geographic variations in alerting, dispatch, and response**. *Emerg Med J* 2005, **22**:216–219.
  18. Zakariassen E, Hunskaar S: **Involvement in emergency situations by primary care doctors on-call in Norway—a prospective population-based observational study**. *BMC Emerg Med* 2010, **10**:5.
  19. Andersen MS, Johnsen SP, Sørensen JN, Jepsen SB, Hansen JB, Christensen EF: **Implementing a nationwide criteria-based emergency medical dispatch system: a register-based follow-up study**. *Scand J Trauma Resusc Emerg Med* 2013, **21**:53.
  20. Zakariassen E, Hansen EH, Hunskaar S: **Incidence of emergency contacts (red responses) to Norwegian emergency primary healthcare services in 2007—a prospective observational study**. *Scand J Trauma Resusc Emerg Med* 2009, **17**:30.
  21. Hinchey P, Myers B, Zalkin J, Lewis R, Garner D: **Low acuity EMS dispatch criteria can reliably identify patients without high-acuity illness or injury**. *Prehosp Emerg Care* 2007, **11**:42–48.
  22. Heward A, Damiani M, Hartley-Sharpe C: **Does the use of the advanced medical priority dispatch system affect cardiac arrest detection?** *Emerg Med J* 2004, **21**:115–118.
  23. Ramanujam P, Guluma KZ, Castillo EM, Chacon M, Jensen MB, Patel E, Linnick W, Dunford JV: **Accuracy of stroke recognition by emergency medical dispatchers and paramedics—San Diego experience**. *Prehosp Emerg Care* 2008, **12**:307–313.
  24. Clawson J, Olola C, Heward A, Patterson B, Scott G: **Ability of the medical priority dispatch system protocol to predict the acuity of “unknown problem” dispatch response levels**. *Prehosp Emerg Care* 2008, **12**:290–296.
  25. Heward A: **Dispatch research: Someone, somewhere needs to know**. *Ann Emerg Dispatch Response* 2013, **1**:5–8.
  26. Ebben RHA, Vloet LCM, Schalk DMU, Mintjes-de Groot JAJA, van Achterberg T: **An exploration of factors influencing ambulance and emergency nurses' protocol adherence in the Netherlands**. *J Emerg Nurs* 2012, [Epub ahead of print].

doi:10.1186/1757-7241-22-2

Cite this article as: Ellensen et al.: Variations in contact patterns and dispatch guideline adherence between Norwegian emergency medical communication centres – a cross-sectional study. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* 2014 **22**:2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit









RESEARCH ARTICLE

Open Access



# Dispatch guideline adherence and response interval—a study of emergency medical calls in Norway

Eirin N. Ellensen<sup>1,2\*</sup>, Torben Wisborg<sup>3,4,5</sup>, Steinar Hunskaar<sup>2,6</sup> and Erik Zakariassen<sup>1,2,6</sup>

## Abstract

**Background:** The Emergency Medical Communication Centre (EMCC) operators in Norway report using the Norwegian Index for Medical Emergency Assistance (Index), a criteria-based dispatch guideline, in about 75 % of medical emergency calls. The main purpose of a dispatch guideline is to assist the operator in securing a correct response as quickly as possible. The effect of using the guideline on EMCC response interval is as yet unknown. We wanted to ascertain an objective measure of guideline adherence, and explore a possible effect on emergency medical dispatch (EMD) response interval.

**Methods:** Observational cross-sectional study based on digital telephone recordings and EMCC records; 299 random calls ending in acute and urgent responses from seven strategically selected EMCCs were included. Ability to confirm location and patient consciousness within an acceptable time interval and structural use of criteria cards were indicators used to create an overall guideline adherence variable. We then explored the relationship between different levels of guideline adherence and EMD response interval.

**Results:** The overall guideline adherence was 80 %. Location and patient consciousness were confirmed within 1 min in 83 % of the calls. The criteria cards were used systematically as intended in 64 % of the cases. Total median response interval was 2:28, with 2:01 for acute calls and 4:10 for urgent calls ( $p < 0.0005$ ). Lower guideline adherence was associated with higher EMD response interval ( $p < 0.0005$ ).

**Conclusion:** The measured guideline adherence was higher than previously reported by the operators themselves. Patient consciousness was rapidly confirmed in the majority of cases. Failure to use Index criteria as intended result in delayed ambulance dispatch and a potential risk of undertriage.

**Keywords:** Telephone recordings, Guideline adherence, Dispatch, Emergency medicine

## Background

The Emergency Medical Communication Centres (EMCCs) are important links in the chain of prehospital emergency medicine in Norway. They are responsible for receiving and triaging phone calls made to the toll-free public emergency medical telephone number 113, dispatching and administering the entire ambulance fleet including the air ambulance service, alerting primary care doctors on-call, and facilitating the communication between prehospital and in-hospital

health personnel in acute medical situations. The EMCCs are manned by registered nurses who answer the 113 calls, and ambulance personnel who coordinate the ambulance fleet. Norwegian Index for Medical Emergency Assistance (Index) is a criteria-based dispatch (CBD) guideline that supplements the medical knowledge and experience of the individual telephone operator during the dispatch process [1]. The criteria based approach to medical dispatch allows for a more open and dynamic interaction between caller and operator than stricter algorithm-based protocols like the Medical Priority Dispatch System (MPDS). CBDs have been criticized because it is difficult to monitor the dispatch process and determine guideline

\* Correspondence: eirin.ellensen@uib.no

<sup>1</sup>Department of Research, Norwegian Air Ambulance Foundation, PB 94, 1441 Drøbak, Norway

<sup>2</sup>Department of Global Public Health and Primary Care, University of Bergen, PB 7800, 5020 Bergen, Norway

Full list of author information is available at the end of the article



adherence, thereby complicating both quality improvement processes and research [2–4].

A 2013 systematic review found that there is an lack of studies on an international basis addressing the two principal systems concerning guideline adherence and protocol compliance [5]. A 2014 study compared cardiac arrest calls processed by a MPDS protocol in Richmond, USA, and the Norwegian CBD guidelines [6]. Indicators for protocol compliance/guidelines adherence were clarification of consciousness (100 and 97 % for MPDS and CBD, respectively), and detection of respiratory arrest (100 and 98 %, respectively). The generalization of these results to the protocol and guidelines as a whole is limited, given the narrow patient groups on which the study focused. A national questionnaire survey among the Norwegian EMCC operators showed a wide range in self-reported use of the Index, not only on the level of individual operators, but also on the EMCC level [2]. The mean self-reported use of the Index found in that study corresponded to operators' use of Index in approximately 75 % of all medical emergency phone calls. Another Norwegian study from 2008 focused on impaired consciousness and intoxication [7]. The study revealed a discrepancy between the use of the Index obtained through operators' questionnaires and recorded phone calls. Some 99 % of the operators claimed to use the Index, while the recordings showed that the level of consciousness was clarified in only 64 % of the contacts.

The aim of this study was to measure guideline adherence objectively through evaluation of real telephone recordings. Secondly, we wanted to investigate a possible association between use of the Index and the emergency medical dispatch (EMD) response interval.

## Method

### The Norwegian Index for Medical Emergency Assistance (Index)

The Index presupposes medical knowledge and is to be used by healthcare staff who are educated and experienced. It consists of a Start page and 36 symptom-based cards. The operator begins at the Start page to obtain important information on location, the state of consciousness and the situation at hand before deciding on immediate acute ambulance dispatch, life-saving instructions, or if the time allows; the most suitable symptom card to continue exploring the situation. The first three cards (Unconscious, Unconscious child and Obstruction of airways) are mainly instructional, with a focus on cardiopulmonary resuscitation (CPR). Each of the remaining cards consists of a list of criteria, corresponding responses, additional questions, caller advices, and supporting instructions for health personnel at scene. The guidelines use three different colour categories according to urgency: red, yellow and green. The colour red is for acute and

life-threatening situations, yellow is for urgent and potentially life-threatening situations, and green is for non-urgent situations. The list of criteria decline in urgency from the top down, and the operators are instructed to start from the top at the most acute criterion, reformulate into questions and work their way down the page until one criterion is finally met. As each criterion is pre-defined as acute, urgent or non-urgent, confirmation of a criterion automatically releases a correlated response.

### Study design and study sample

We performed an observational cross-sectional study of acute and urgent 113 calls during a 72-h period in late August 2011. Nine of Norway's 19 EMCCs were invited to participate; one did not have recordings from the inclusion period, one did not respond and seven agreed to join the study. The selection was strategically based on diversity in population density (5–210 inhabitants/km<sup>2</sup>), contact rates (36–75 contacts/1 000 inhabitants/year), geographic area (all regions represented) and self-reported use of Index at EMCC level (3.7–4.4). "Self-reported use of Index" originated from a previously published questionnaire study [2]. The EMCC operators were given a Likert scale of 11 questions on how often they used different parts of Index, with 5 response formats (1 = never, 2 = seldom, <25 %, 3 = sometimes, 4 = often, >75 %, 5 = always), giving each operator and EMCC a mean value on use of Index between 1 (=never) and 5 (=always) (2). The selected study sample had a contact rate of 57/1 000 inhabitants per year, compared to the national contact rate of 56/1 000 [2]. The mean self-reported use of Index in the sample was 3.86 (SD 0.40), slightly below the national value of 3.95 (SD 0.39) and corresponded to the response format "often, >75 %" (=4) [2].

### Data collection

We collected a random sample of 50 digitalized call recordings from each of the five largest EMCCs, and 20 and 30 from the two smallest EMCCs. Randomization was achieved through a number generator ([www.random.org](http://www.random.org)). Time records, set dispatch criteria and urgency were obtained from Acute Medical Information System (AMIS), the software program used to register and document each contact. One call recording was excluded because of missing AMIS information.

We used a pre-developed form when analysing the call recordings [Additional file 1], based on an internal form used for guidelines adherence and quality control by one of the participating EMCCs. Each call recording was analysed on the basis of six different indicators; (1) confirmation of caller phone number, (2) location, (3) patient consciousness, (4) patient responsiveness, (5) criteria compliance, and (6) communication problems. Both information given spontaneously by the caller and information

asked specifically for by the operator were registered, but merged for analyses.

Indicators 1–4 were from the Start page. This information was categorized into “within 1 min”, “within 2 min” or “not at all”. Information gained after 2 min was interpreted as negative. Criteria compliance (indicator 5) was measured by the number of criteria above the set criterion not accounted for during the call. Hence a criteria compliance value of 0 means that all criteria above the one set by the operator were accounted for during the call, whereas increasing criteria compliance values mean that the operator has overlooked some more urgent criteria. The variable “Criteria compliance” was grouped in 0, 1 and 2 and more. Indicator 6 (communication problems) was registered as positive if the operator failed to understand what the caller was trying to communicate, as assessed by the researcher.

The main outcome variable “Overall guideline adherence” was composed from the three variables confirmation of location, confirmation of consciousness and criteria compliance. Each variable was dichotomized, and information gained within 1 min for the two first variables, and zero or one unchecked criteria above the set criterion were interpreted as positive. The variables were weighted equally, composing a measured guideline adherence score of 0 (no positive) to 3 (all positive). The secondary outcome variable “EMD response interval” was defined as time from the call was answered until an ambulance was dispatched [8]. Time intervals are reported as minutes:seconds (mm:ss). The data are available as an additional file [Additional file 2].

### Statistical analysis

Continuous data are presented as means (SD) or medians (interquartile range, IQR) for symmetric and skewed data respectively. Rates are calculated as contacts per 1 000 inhabitants per year. Response interval was analysed as a continuous variable (seconds). Q-Q plots were used to check the outcome variables for normal distribution. Spearman rho was used to investigate correlations between the main variable and secondary variables. One-way analysis of variance (ANOVA) was used to compare mean scores on the main outcome variable between EMCC centres, and Mann-Whitney U test and Kruskal-Wallis tests to compare scores on the secondary outcome variable EMD response interval.

We performed statistical analyses using Statistical Package for the Social Sciences (IBM SPSS Statistics 23). *P*-values < 0.05 were considered statistically significant.

### Results

Of the 299 call recordings included in the study, 279 resulted in an ambulance dispatch; 99 % of the 174 acute contacts resulted in an ambulance dispatch ( $n = 173$ ),

whereas 87 % of the 125 urgent contacts released an ambulance dispatch ( $n = 106$ ). Table 1 shows an overview of the six Index indicators measured in the study.

### Adherence to guideline

The mean overall guideline adherence was 2.41 (0.73), 80 % of maximum value of 3. The EMCCs differed in guideline adherence from 75 to 89 % (Table 2), but one-way ANOVA showed no statistically significant differences between the centres ( $p = 0.073$ ).

There was a low, but statistically significant correlation, between overall guideline adherence and urgency ( $\rho = 0.27$ ,  $p < 0.0005$ ), revealing that acute contacts had higher levels of overall guideline adherence, compared with urgent contacts. There was no statistically significant correlation between overall guideline adherence and self-reported use of the Index, EMCC contact rate or population density.

### EMD response interval

Location, telephone number or consciousness status confirmations within 1, 2 min or not at all did not affect the EMD response interval to any statistically significant extent. Decreasing criteria compliance increased the response interval ( $p = 0.001$ ) (Table 1). Difficulties in communication due to language, non-cooperation or caller not on the scene did not affect EMD response interval.

The set level of urgency had a strong effect on the response interval, with urgent contacts having a median response interval of 4:10 min compared to acute contacts with 2:01 min. This increase of 2:09 min was statistically significant ( $p < 0.0005$ ). Increasing overall guideline adherence decreased EMD response interval ( $p < 0.0005$ ) (Table 3).

### Discussion

In this study we were able to measure actual use of the Index, and found a higher mean guideline adherence value compared to the previously published self-reported use of Index value [2]. Higher overall guideline adherence led to a statistically, significantly lower EMD response interval.

Considering that the guidelines are designed to function in cooperation with the operators’ skills and experience, it is difficult to designate a limit for acceptable use of the guidelines as a whole, and thus assess whether the mean guideline adherence of 80 % is acceptable or not. Start Page compliance, especially confirmation of patient location and consciousness, should be 100 % in an emergency medical call. The ideal compliance level with regards to the criteria cards will however be somewhat lower, due to the nature of the guidelines. The fact that the Index is still being used in a paper version in an otherwise electronic work environment can probably account for some of the situations where it is not used. Except occasional feedback

**Table 1** The six Index indicators and their effect on EMD response interval<sup>a</sup>

	All calls (N = 299)		Ambulance dispatches (N = 279)		EMD response interval (min:sec) <sup>a</sup>		P-value
	N	%	N	%	Median	IQR <sup>b</sup>	
Caller's phone number confirmed							0.973
Within 1 min	59	20	56	20	2:49	2:01	
1–2 min	19	6	18	6	2:21	2:31	
After 2 min/none	221	74	205	73	2:28	2:54	
Location confirmed							0.225
Within 1 min	247	83	237	85	2:23	2:36	
1–2 min	23	8	22	8	3:00	2:52	
After 2 min/none	29	10	20	7	3:11	4:27	
Patient's consciousness							0.429
Within 1 min	249	83	231	83	2:23	2:32	
1–2 min	24	8	23	8	3:12	4:27	
After 2 min/none	26	9	25	9	2:39	3:00	
Patient's responsiveness							0.086
Within 1 min	240	80	224	80	2:23	2:33	
1–2 min	29	10	26	9	3:57	3:56	
After 2 min/none	30	10	29	10	2:03	2:26	
Criteria compliance <sup>c</sup>							0.001
All criteria confirmed	191	64	184	66	2:17	2:17	
1 criterion unconfirmed	33	11	29	10	2:39	2:04	
2+ criteria unconfirmed	75	25	66	24	3:48	4:59	
Communication							0.691
Good	256	86	241	86	2:33	2:39	
Difficult <sup>d</sup>	43	14	38	14	2:22	2:48	

<sup>a</sup>Emergency Medical Dispatch response interval: Time from 113 call is answered until an ambulance is dispatched

<sup>b</sup>IQR Interquartile range

<sup>c</sup>How many criteria above the one set by the operator is not answered for during the call

<sup>d</sup>Difficult communication includes language difficulties, non cooperative caller and caller not on scene (third hand information)

from ambulance personnel on the scene, the EMCC operators never receive information on patient outcome. The relationship between guidelines and clinical outcome is known to be an important motivational factor for guideline adherence [5].

Although the 2009 study on impaired consciousness also found a discrepancy between self-reported use of Index (99 %) and measured use of Index (64 % confirmed consciousness) [7] the discrepancy is not only much wider, but also the opposite of our findings. This could indicate a

**Table 2** Overall guideline adherence distribution among the EMCCs

EMCC no	Guideline adherence value distribution (%)				Mean value (SD)	% of maximum
	0	1	2	3		
1	3	7	40	50	2.37 (0.77)	79
2	5	0	35	60	2.50 (0.76)	83
3	0	10	46	44	2.34 (0.66)	78
4	2	16	32	50	2.30 (0.81)	77
5	0	22	31	47	2.24 (0.80)	75
6	2	10	28	60	2.46 (0.76)	82
7	0	2	28	70	2.68 (0.51)	89
All	1	11	34	54	2.41 (0.73)	80

**Table 3** Overall guideline adherence and effect on EMD response interval<sup>a</sup>

	All calls (N = 299)		Ambulance dispatches (N = 279)		EMD response interval (min:sec) <sup>a</sup>		p-value
	N	%	N	%	EMD response interval (min:sec) <sup>a</sup>		
					Median	IQR <sup>b</sup>	
Overall guideline adherence							<0.0005
3	162	54	157	56	2:12	2:03	
2	101	34	92	33	3:19	3:56	
1	32	11	26	9	3:03	4:03	
0	4	1	4	1	4:28	10:36	0.225

<sup>a</sup>Emergency Medical Dispatch response interval: Time from 113 call is answered until an ambulance is dispatched

<sup>b</sup>IQR Interquartile range

positive development towards both higher actual use of Index and a more realistic view on own use.

The 2014 study comparing medical priority dispatch system (MPDS) and criteria-based dispatch (CBD) in cardiac arrest patients reported a guideline adherence of 97 % for successful clarifying level of consciousness [6]. This is higher than our finding that consciousness was clarified within 1 min in 83 % of the cases. One obvious explanation for this difference is their selection of highly acute contacts only, while our material includes all acute and urgent contacts. Furthermore, although not statistically significant, our study shows a wide variety between the EMCCs when it comes to percentage confirmed consciousness, ranging from 73 to 92 % for all acute and urgent cases in total. It has been shown that higher call processing rates increase dispatch performance. Kuisma et al. found in 2005 that operators processing >9 ventricular fibrillation cardiac arrest calls during the study period recognised cardiac arrest faster, dispatched first responding unit faster, gave cardiopulmonary resuscitation instructions and received bystander cooperation more often, and had a higher rate of patients discharged alive than dispatchers processing <4 VF calls during the study period [9].

Since the introduction of Index in Norway, it has been debated whether it saves time or adds time to the dispatch process. Of the six Index indicators investigated in our study, criteria compliance was the only one with a statistically significant effect on EMD response interval. The main effect on EMD response interval occurred between 1 and 2 unconfirmed criteria, with an increase of 1:31 (minutes:seconds) from 0 unconfirmed criteria to 2+ unconfirmed criteria. When looking at the combined outcome variable *overall guideline adherence*, we found a difference of 2:16 between the lowest and highest value group.

Failure to check vital information like consciousness or responsiveness obstruct detection of life-threatening situations like cardiac arrest or breathing difficulties where instructions from the operator might influence the outcome significantly. Berdowski et al. found in a study

published in 2009 that failure to recognize cardiac arrest during the emergency call delayed both ambulance dispatch and arrival on the scene, and decreased 3-month survival from 14 to 5 % [10]. Using a similar protocol to our Start page, the Dutch dispatchers were supposed to clarify location, phone number, patient consciousness and breathing, and the study showed that failure to ask whether and how the patient was breathing was the primary reason for not recognizing the cardiac arrest. Previous versions of Index included breathing as a key question on the Start page; this was replaced by patient responsiveness in 2009 [1].

Low criteria compliance might result in a lower acuity than actually needed because important information is unrevealed, the patient might lose important time due to undertriage. A certain over triage is generally accepted in the prehospital emergency system, especially considering the difficulties of assessing a situation solely over the telephone, but undertriage may put the patient at unnecessary risk. An Italian study on dispatch errors focusing on patients initially coded as non-urgent and ending up as fatalities found that these calls were predominantly made by next of kin reporting not life-threatening symptoms, but they were also shorter and characterized by inadequate collection of vital information by the operators [11]. Andersen et al. studied a group of patients not identified as acute but who died the same day they were reported via a 112 call. The researchers found that although none of the deaths were considered definitively preventable, 12 % were considered potentially preventable if the call had been assessed as acute [12]. A Swedish study published in 2011 compared priority codes between the dispatch centre (CBD guidelines) and the ambulance on scene (Medical Emergency Triage and Treatment System—A, METTS-A) and found that although the Swedish index has a high sensitivity for identifying acute patients, 4.8 % of the patients were undertriaged [13]. A similar finding of 4.6 % undertriage was found in Switzerland in a study published in 2015 [14]. In Norway, experiences of general practitioners (GPs) participating in prehospital emergency situations show that 42 % of the

patients are downgraded between initial dispatch centre triaging and on-the-scene triaging by a GP, while 11 % were upgraded, acute abdominal cases having the highest risk of being initially undertriaged [15].

### Strengths and limitations

Although only seven EMCCs took part in this study, they were strategically selected to be representative of the rest of the EMCCs and the results are therefore thought to be generalizable on a national level. This study thus provides valuable information on how and to what extent Index is actually used and how this affects the response interval.

The outcome variable *overall guideline adherence* derived from three of the six primary indicators, excluding telephone number confirmation, patient responsiveness and caller cooperation. Since the implementation of automatic number tracking of incoming calls, there seems to be a cultural acceptance at the EMCCs for only asking for the telephone number when the number is unregistered. Including this variable would have given a marked decrease in mean overall guideline adherence. The variable patient responsiveness provided little information beyond that provided by consciousness, and the numbers of uncooperative callers or language issues were very small. The cut-off value for the Start page indicators was chosen to be 'within one minute', as these are time-critical questions essential for further optimal handling of the situation. The cut-off for criteria compliance was set between one and two criteria not accounted for during the call, based on the effect on EMD response interval.

Our findings suggest higher guideline adherence compared to previous studies. Given the nature of CBD guidelines as an aid to the operator in the dispatch process, rather than the operator carrying out the dispatch process itself, one might question whether the present level of Index adherence is acceptable. However, further increased confirmation of consciousness and proper use of the criteria cards may reduce the risk of undertriage. Documenting overall guideline adherence is a further step towards exploring the validity of Index.

### Conclusion

The mean overall guideline adherence for acute and urgent emergency calls was 80 %. The results confirm previously self-reported use of Index findings. Low overall guideline adherence increases the EMD response interval and may increase the risk of undertriage.

### Additional files

**Additional file 1:** Index-113-registration, a call recordings registration form on adherence to different parts of index during the emergency call. (PDF 212 kb)

**Additional file 2:** Index-113calls work sheet, a data file supporting the conclusions of the article. (XLSX 64 kb)

### Abbreviations

AMIS: Acute Medical Information System; ANOVA: Analysis of variance; CBD: Criteria-based dispatch; CPR: Cardiopulmonary resuscitation; EMCC: Emergency Medical Communication Centre; EMD: Emergency medical dispatch; GP: General practitioner; IQR: Interquartile range; METTS-A: Medical Emergency Triage and Treatment System—A; MPDS: Medical Priority Dispatch System; SD: Standard deviation; The Index: The Norwegian Index for Medical Emergency Assistance

### Acknowledgements

We would like to thank both management and operators at Bergen EMCC, Buskerud EMCC, Harstad EMCC, Oslo and Akershus EMCC, Sørlandet EMCC, Sør-Trøndelag EMCC, and Tromsø EMCC for their cooperation in providing data material for this study.

We would also like to thank all supporting members of the Norwegian Air Ambulance Foundation, for making the emergency medicine research possible.

### Funding

This project has been made possible by the Norwegian ExtraFoundation for Health and Rehabilitation and the Norwegian Air Ambulance Foundation.

### Availability of data and materials

The pre-developed form used to analyse the call recordings and the dataset supporting the conclusions of this article are included as Additional files 1 and 2.

### Authors' contribution

All authors participated in planning the study design. EE collected the data, performed the statistical analyses and drafted the manuscript. TW, SH and EZ participated in critical revision of the manuscript. All authors read and approved the final manuscript.

### Competing interests

The authors declare no competing interests.

### Consent for publication

Not applicable.

### Ethical approval and consent to participate

The study was assessed by the Regional Committee for Medical and Health Research Ethics, and was deemed to be in no need of ethical approval (2011/756/REK vest). The committee also exempted EMCC operators and patients' consent pertaining to analysis of sensitive data.

### Author details

<sup>1</sup>Department of Research, Norwegian Air Ambulance Foundation, PB 94, 1441 Drøbak, Norway. <sup>2</sup>Department of Global Public Health and Primary Care, University of Bergen, PB 7800, 5020 Bergen, Norway. <sup>3</sup>Anaesthesia and Critical Care Research Group, Faculty of Health Sciences, University of Tromsø, Tromsø 9037, Norway. <sup>4</sup>Norwegian National Advisory Unit on Trauma, Oslo University Hospital, Oslo 0450, Norway. <sup>5</sup>Department of Anesthesia and Intensive Care, Hammerfest Hospital, Finnmark Hospital Trust, Hammerfest 9613, Norway. <sup>6</sup>National Centre for Emergency Primary Health Care, Uni Research Health, Kalfarveien 31, Bergen 5018, Norway.

Received: 15 June 2016 Accepted: 24 September 2016

Published online: 13 October 2016

### References

1. Norwegian Medical Association. Norsk indeks for medisinsk nødhjelp (Norwegian Index for Medical Emergency Assistance). 3rd ed. Stavanger: Laerdal Medical A/S; 2009.
2. Ellensen EN, Hunnskaar S, Wisborg T, Zakariassen E. Variations in contact patterns and dispatch guideline adherence between Norwegian emergency medical communication centres - a cross-sectional study. *Scand J Trauma Resusc Emerg Med.* 2014;22:2.

3. Clawson JJ, Martin RL, Hauert SA. Protocols vs. guidelines. Choosing a medical-dispatch program. *Emerg Med Serv.* 1994;23:52–60.
4. Clawson J, Olola CHO, Heward A, Scott G, Patterson B. Accuracy of emergency medical dispatchers' subjective ability to identify when higher dispatch levels are warranted over a Medical Priority Dispatch System automated protocol's recommended coding based on paramedic outcome data. *Emerg Med J.* 2007;24:560–3.
5. Ebben RHA, Vloet LCM, Verhofstad MHJ, Meijer S, Mintjes-de Groot JAJ, van Achterberg T. Adherence to guidelines and protocols in the prehospital and emergency care setting: a systematic review. *Scand J Trauma Resusc Emerg Med.* 2013;21:9.
6. Hardeland C, Olasveengen TM, Lawrence R, Garrison D, Lorem T, Farstad G, Wik L. Comparison of Medical Priority Dispatch (MPD) and Criteria Based Dispatch (CBD) relating to cardiac arrest calls. *Resuscitation.* 2014;85:612–6.
7. Tonje L, Elisabeth S, Lars W. Handling of drug-related emergencies: an evaluation of emergency medical dispatch. *Eur J Emerg Med.* 2009;16:37–42.
8. Castrén M, Karlsten R, Lippert F, Christensen EF, Bovim E, Kvam AM, Robertson-Steel I, Overton J, Kraft T, Engerstrom L, Garcia-Castrill Riego L. Emergency Medical Dispatch expert group at the Utstein Consensus Symposium 2005. Recommended guidelines for reporting on emergency medical dispatch when conducting research in emergency medicine: the Utstein style. *Resuscitation.* 2008;79:193–7.
9. Kuisma M, Boyd J, Väyrynen T, Repo J, Nousila-Wiik M, Holmström P. Emergency call processing and survival from out-of-hospital ventricular fibrillation. *Resuscitation.* 2005;67:89–93.
10. Berdowski J, Beekhuis F, Zwinderman AH, Tijssen JGP, Koster RW. Importance of the first link: description and recognition of an out-of-hospital cardiac arrest in an emergency call. *Circulation.* 2009;119:2096–102.
11. Palma E, Antonaci D, Coli A, Cicolini G. Analysis of emergency medical services triage and dispatch errors by registered nurses in Italy. *J Emerg Nurs.* 2014;40:476–83.
12. Andersen MS, Johnsen SP, Hansen AE, Skjaereth E, Hansen CM, Sørensen JN, Jepsen SB, Hansen JB, Christensen EF. Preventable deaths following emergency medical dispatch—an audit study. *Scand J Trauma Resusc Emerg Med.* 2014;22:74.
13. Ek B, Edström P, Toutin A, Svedlund M. Reliability of a Swedish pre-hospital dispatch system in prioritizing patients. *Int Emerg Nurs.* 2013;21:143–9.
14. Dami F, Golay C, Pasquier M, Fuchs V, Carron P-N, Hugli O. Prehospital triage accuracy in a criteria based dispatch centre. *BMC Emerg Med.* 2015;15:32.
15. Rørtveit S, Meland E, Hunskaar S. Changes of triage by GPs during the course of prehospital emergency situations in a Norwegian rural community. *Scand J Trauma Resusc Emerg Med.* 2013;21:89.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



