

Polyphosphoinositides in the nucleus: Roadmap of their effectors and mechanisms of interaction



Rhian G. Jacobsen^a, Fatemeh Mazloumi Gavvani^{a,1}, Amanda J. Edson^a,
Marianne Goris^b, Altanchimeg Altankhuyag^{c,d}, Aurélia E. Lewis^{a,*}

^a Department of Biological Sciences, University of Bergen, Bergen, 5008, Norway

^b NORCE Norwegian Research Center, Bergen, 5008, Norway

^c Department of Biomedicine, University of Bergen, Bergen, 5021, Norway

^d Department of Clinical Science, University of Bergen, Bergen, 5021, Norway

ARTICLE INFO

Keywords:

Polyphosphoinositides
Phosphoinositides
Nucleus
Electrostatic interactions
Polybasic motif
K/R motif

ABSTRACT

Biomolecular interactions between proteins and polyphosphoinositides (PPIIn) are essential in the regulation of the vast majority of cellular processes. Consequently, alteration of these interactions is implicated in the development of many diseases. PPIIn are phosphorylated derivatives of phosphatidylinositol and consist of seven species with different phosphate combinations. PPIIn signal by recruiting proteins via canonical domains or short polybasic motifs. Although their actions are predominantly documented on cytoplasmic membranes, six of the seven PPIIn are present within the nucleus together with the PPIIn kinases, phosphatases and phospholipases that regulate their turnover. Importantly, the contribution of nuclear PPIIn in the regulation of nuclear processes has led to an increased recognition of their importance compared to their more accepted cytoplasmic roles. This review summarises our knowledge on the identification and functional characterisation of nuclear PPIIn-effector proteins as well as their mode of interactions, which tend to favour polybasic motifs.

1. Introduction

Polyphosphoinositides (PPIIn) (nomenclature as detailed in (Michell *et al.*, 2006)) are phosphorylated derivatives of the glycerophospholipid phosphatidylinositol (PtdIns), composed of two hydrophobic fatty acyl tails esterified to a glycerol backbone, which is itself linked to the D-myoinositol head group (Fig. 1A). Phosphorylation/dephosphorylation of the hydroxyl groups present on positions 3, 4 and 5 of the inositol head group by PPIIn kinases and phosphatases can produce seven distinct PPIIn, *i.e.* three monophosphorylated PPIIn (PtdIns3P, PtdIns4P and PtdIns5P), three diphosphorylated (PtdIns(3,4)P₂, PtdIns(3,5)P₂ and PtdIns(4,5)P₂) and one triphosphorylated (PtdIns(3,4,5)P₃) (Balla, 2013). The presence of acyl chains allows PPIIn to be embedded in membranes. On the other hand, the headgroup is exposed to the solvent where it can interact with proteins containing well-defined structured PPIIn-binding domains or short polybasic motifs (Hammond and Balla, 2015; Pemberton and Balla, 2018). In both cases, interaction occurs through electrostatic interactions between basic residues and the phosphates displayed on the headgroup. Depending on the level and localisation of PPIIn species, different proteins can be recruited to specific membrane locations in a timely manner, hence providing the cell with a PPIIn-dependent signalling code.

* Corresponding author.

E-mail address: aurelia.lewis@uib.no (A.E. Lewis).

¹ present address: The Sars International Centre for Marine Molecular Biology, Bergen 5006, Norway.

<https://doi.org/10.1016/j.jbior.2019.04.001>

Received 20 March 2019; Received in revised form 2 April 2019; Accepted 3 April 2019

Available online 06 April 2019

2212-4926/ © 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abbreviations	
ABP	actin-binding protein
BASP1	brain acid soluble protein 1
BAZ2B	BROMO domain adjacent to zinc finger 2B
CKI α	casein kinase I α
DAG	diacylglycerol
EBP1	ErbB3 binding protein-1
FYVE	Fab1p, YOTB, Vac1 and EEA1
GAP43	growth associated protein 43
HDAC1	histone deacetylase 1
ING2	inhibitor of growth protein 2
IP ₃	inositol 1,4,5-trisphosphate
IPMK	inositol polyphosphate multikinase
K/R	lysine/arginine motif following the motif K/R-(X ₃₋₇)-K-X-K/R-K/R
LRH-1	liver receptor homolog-1
MARCKS	myristoylated alanine-rich C-kinase substrate
NLS	nuclear localisation signal
NoLS	nucleolar localisation signal
N-WASP	neuronal Wiskott-Aldrich syndrome protein
OGT	O-linked beta-N-acetylglucosamine transferase
PBR	polybasic region
Pf1	PHD factor 1
PH	pleckstrin homology
PHD	plant homeodomain
PHF8	PHD finger protein 8
PI3K	phosphoinositide 3-kinase
PI4K	phosphatidylinositol 4-kinase
PIKE-L	PI3K enhancer-L
PIP ₃ BP	PtdIns(3,4,5)P ₃ binding protein
PIP2K	phosphatidylinositol phosphate kinase
PLC	phospholipase C
Pol I/II	RNA polymerase I/II
PPIn	polyphosphoinositide
PtdIns	phosphatidylinositol
PTEN	phosphatase and tensin homolog deleted on chromosome 10
RNP	ribonucleoprotein
SAP30	sin3A-associated protein 30
SAP30L	human SAP30-like protein
SF-1	steroidogenic factor-1
SHIP	src homology 2 (SH2) domain containing inositol phosphatase
TAF3	transcription initiation factor TFIID subunit 3
Topo II α	DNA topoisomerase II α
UBF	upstream binding factor
UHRF1	ubiquitin-like PHD and RING finger domain-containing protein 1

Since the 1950s when the PPIn cycle was discovered, research efforts have focused on the biochemical and functional characterisation of PPIn located at the plasma membrane and further on organelle membranes. It took at least another decade before the first studies demonstrated the existence of phospholipids (Manzoli *et al.*, 1977; Rose and Frenster, 1965) and then of PPIn within the confines of the nucleus, *i.e.* separate from the nuclear envelope (Cocco *et al.*, 1987; Payrastra *et al.*, 1992). Indeed, not only were PPIn detected in nuclei stripped of their nuclear envelope, but also their levels changed upon the addition of different cell stimuli (Clarke *et al.*, 2001; Cocco *et al.*, 1989; Divecha *et al.*, 1991). Since then, evidence has accumulated to detail the localisation of PPIn and the identification of effector proteins in the nucleus. This led to the recognition of PPIn being instrumental in a number of different nuclear processes such as splicing and chromatin remodelling (see recent reviews (Barlow *et al.*, 2010; Davis *et al.*, 2015; Fiume *et al.*, 2012; Hamann and Blind, 2018; Martelli *et al.*, 2011; Shah *et al.*, 2013)). Identifying which effector proteins these PPIn interact with and understanding their mode of interaction are key to deciphering the functions of these lipids in the nucleus. This review focuses on these two aspects.

2. Nuclear polyphosphoinositides and their metabolic enzymes

All PPIn, except PtdIns(3,5)P₂, have been detected in the nucleus by using different approaches (Table 1 and reviewed in (Fiume *et al.*, 2012; Shah *et al.*, 2013)). PtdIns3P has been detected in nuclei by the use of a specific probe consisting of two FYVE (Fab1p, YOTB, Vac1p and EEA1) domains and electron microscopy (Gillooly *et al.*, 2000). Mass levels of nuclear PtdIns4P, PtdIns5P and PtdIns(4,5)P₂ have been measured by radiolabelling (Clarke *et al.*, 2001). PtdIns(4,5)P₂ has also been identified in the nucleus using a PtdIns(4,5)P₂ specific probe, the pleckstrin homology (PH) domain of phospholipase C (PLC) δ 1 combined with electron microscopy (Watt *et al.*, 2002). In addition, using immunostaining, PtdIns(4,5)P₂ was detected in nuclear speckles along with mRNA-processing components, in the newly described nuclear lipid islets as well as in nucleoli but as a minor detectable pool (Boronnikov *et al.*, 1998; Osborne *et al.*, 2001; Sobol *et al.*, 2013; Sobol *et al.*, 2018; Yildirim *et al.*, 2013). Using either a specific PtdIns(3,4,5)P₃ probe (the PH domain of general receptor of phosphoinositides 1) or antibodies, PtdIns(3,4,5)P₃ was shown to localise to the nucleoplasm and nucleolus (Karlsson *et al.*, 2016; Kumar *et al.*, 2010; Kwon *et al.*, 2010; Lindsay *et al.*, 2006).

Consistently, several PPIn metabolising enzymes are present in the nucleus, notably in discrete sub-nuclear areas common to the localisation of their respective PPIn substrate and/or product (Table 2 and Fig. 1B). In particular, strong evidence shows the presence of a PPIn metabolic cycle in nuclear speckles (alias inter chromatin granule clusters), hubs of mRNA processing in association with transcriptional events (Spector and Lamond, 2011) and the nucleolus, the ribosome factory (Pederson, 2011). The type III PtdIns4K α and β isoforms, generating PtdIns4P, have been detected in nucleoli and nuclear speckles respectively (Kakuk *et al.*, 2006, 2008; Szivak *et al.*, 2006). The β form, in particular, was found in nuclear speckles when phosphorylated on Serine 496 or Threonine 504, while the Serine 294 phosphorylated enzyme is present on the Golgi (Szivak *et al.*, 2006). The type I phosphatidylinositol phosphate kinase (PIPKI α), which phosphorylates PtdIns4P to generate PtdIns(4,5)P₂, has also been found in the nucleolus in G1/S phase and shown to affect rRNA gene silencing through its physical association with H3K9me3 and heterochromatin protein 1 α (Chakrabarti *et al.*, 2015). PIPKI α as well as PIPKI γ isoform 4, localise to nuclear speckles (Boronnikov *et al.*, 1998; Mellman *et al.*, 2008; Schill and




Fig. 1. Nuclear polyphosphoinositide metabolism. **A**) Phosphatidylinositol (PtdIns) structure PI(18:0/20:4 (5Z,8Z,11Z, 14Z)) downloaded from the LIPID MAPS structure database (LM ID: LMGP06010010) (Fahy *et al.*, 2009; Sud *et al.*, 2007). Hydroxyl groups that are sites of phosphorylation on the inositol head group are indicated with arrows and carbon atom numbers are shown in red. **B**) Nuclear polyphosphoinositide cycle. Polyphosphoinositide kinases and phosphatases as well as phospholipases known to reside in the nucleus are shown (see also Table 2). Abbreviations: 4-Ptase: 4-phosphatase, DAG: diacylglycerol, IP₃: inositol 1,4,5-trisphosphate, IPMK: inositol polyphosphate multikinase, P: phosphate group, PI3K: phosphoinositide 3-kinase, PI4K: phosphatidylinositol 4-kinase, PIPK: phosphatidylinositol phosphate kinase, PLC: phospholipase C, PTEN: phosphatase and tensin homolog deleted on chromosome 10, SHIP: src homology 2 (SH2) domain containing inositol phosphatase.

Table 1
Nuclear localisation of polyphosphoinositides.

PPIn	Sub-nuclear localisation (References)
PtdIns3P	Nucleolus (Gillooly <i>et al.</i> , 2000)
PtdIns4P	Nucleus (Clarke <i>et al.</i> , 2001; Kalasova <i>et al.</i> , 2016; Vann <i>et al.</i> , 1997) Nucleolus (Kalasova <i>et al.</i> , 2016)
PtdIns5P	Nucleus and chromatin enriched fraction, detected upon stress induction (Clarke <i>et al.</i> , 2001; Jones <i>et al.</i> , 2006)
PtdIns(3,4)P ₂	Nuclear membrane (Watt <i>et al.</i> , 2004; Yokogawa <i>et al.</i> , 2000)
PtdIns(4,5)P ₂	Interchromatin granules/Nuclear speckles (Boronenkov <i>et al.</i> , 1998; Osborne <i>et al.</i> , 2001) Nucleolus (Kalasova <i>et al.</i> , 2016; Osborne <i>et al.</i> , 2001; Sobol <i>et al.</i> , 2013; Yildirim <i>et al.</i> , 2013)
PtdIns(3,4,5)P ₃	Nuclear lipid islets (Sobol <i>et al.</i> , 2018) Nuclear matrix (Kumar <i>et al.</i> , 2010; Kwon <i>et al.</i> , 2010; Lindsay <i>et al.</i> , 2006) Nucleolus (Karlsson <i>et al.</i> , 2016)

Anderson, 2009). These studies are consistent with the possible detection of PtdIns4P in nucleoli as well as nuclear speckles, as reported by Kalasova *et al.*, although no markers were used in the study (Kalasova *et al.*, 2016). In addition, the type II PIPK α and β , which phosphorylate PtdIns5P to generate PtdIns(4,5)P₂, are also present in nuclear speckles (Boronenkov *et al.*, 1998). PLC β 1, which metabolises PtdIns(4,5)P₂ to diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP₃), has also been found in the nucleus and in particular in nuclear speckles (Manzoli *et al.*, 1995; Martelli *et al.*, 1992; Tabellini *et al.*, 2003). Other isoforms are also known to be nuclear, such as PLC δ 1 and δ 4 (Kunrath-Lima *et al.*, 2018; Okada *et al.*, 2010). Finally, consistent with the presence of a PtdIns(3,4,5)

Table 2
Nuclear PPIn-metabolising enzymes, their substrates and nuclear localisation.

Nuclear PPIn kinases		
Enzyme	Substrate	Sub-nuclear localisation (References)
Type III α PI4K PI4K230	PtdIns	Nucleolus (Kakuk <i>et al.</i> , 2006, 2008)
Type III β PI4K PI4K92	PtdIns	Nucleus (de Graaf <i>et al.</i> , 2002) > Nuclear speckles (Szivak <i>et al.</i> , 2006)
Type Ia PIPK	PtdIns4P	Nuclear speckles (Boronenkov <i>et al.</i> , 1998; Mellman <i>et al.</i> , 2008) Nucleolus (Chakrabarti <i>et al.</i> , 2015)
Type I γ PIPK _{i4}	PtdIns4P	Nuclear speckles (Schill and Anderson, 2009)
Type II α PIPK	PtdIns5P	Nucleus (Bultsma <i>et al.</i> , 2010)
Type II β PIPK	PtdIns5P	Nuclear speckles (Boronenkov <i>et al.</i> , 1998)
Class I PI3K, p110 β	PtdIns(4,5)P ₂	Nucleus (Bultsma <i>et al.</i> , 2010; Ciruela <i>et al.</i> , 2000) Nucleoplasm, chromatin, double stranded break foci (Kumar <i>et al.</i> , 2010, 2011; Marques <i>et al.</i> , 2009) Nucleolus (Karlsson <i>et al.</i> , 2016)
IPMK	PtdIns(4,5)P ₂	Nucleus (Resnick <i>et al.</i> , 2005)
Class II α PI3K	PtdIns, PtdIns4P	Nuclear speckles (Didichenko and Thelen, 2001)
Class II β PI3K	PtdIns, PtdIns4P	Nuclear matrix (Sindic <i>et al.</i> , 2001) Nuclear lamina (Banfic <i>et al.</i> , 2009)
Nuclear PPIn phosphatases		
Enzyme	Substrate	Sub-nuclear localisation (References)
PTEN	PtdIns(3,4,5)P ₃	Nucleus (Deleris <i>et al.</i> , 2003; Shen <i>et al.</i> , 2007; Song <i>et al.</i> , 2008) Nucleolus (Li <i>et al.</i> , 2014), PTEN β (Liang <i>et al.</i> , 2017)
SHIP1	PtdIns(3,4,5)P ₃	Nucleolus (Ehm <i>et al.</i> , 2015)
SHIP2	PtdIns(3,4,5)P ₃ or PtdIns(4,5)P ₂ ?	Nuclear speckles (Deleris <i>et al.</i> , 2003; Elong Edimo <i>et al.</i> , 2011)
Type I PtdIns(4,5)P ₂ 4-phosphatase	PtdIns(4,5)P ₂	Translocates to the nucleus upon cell stress (Zou <i>et al.</i> , 2007)
Nuclear phospholipase C		
Enzyme	Substrate	Sub-nuclear localisation (References)
PLC β 1	PtdIns(4,5)P ₂	Nuclear speckles (Tabellini <i>et al.</i> , 2003)
PLC δ 1	PtdIns(4,5)P ₂	Nucleus (Okada <i>et al.</i> , 2010)
PLC δ 4	PtdIns(4,5)P ₂	Nucleus (Kunrath-Lima <i>et al.</i> , 2018)

Abbreviations: IPMK: inositol polyphosphate multikinase, PI3K: phosphoinositide 3-kinase, PI4K: phosphatidylinositol 4-kinase, PIPK: phosphatidylinositol phosphate kinase, PLC: phospholipase C, PPIn: polyphosphoinositide, PtdIns: phosphatidylinositol, PTEN: phosphatase and tensin homolog deleted on chromosome 10, SHIP: src homology 2 (SH2) domain containing inositol phosphatase.

P₃ pool in the nucleus and nucleolus, enzymes generating (class IA phosphoinositide 3-kinase (PI3K) p110 β) or metabolising PtdIns (3,4,5)P₃ (phosphatase and tensin homolog deleted on chromosome 10 (PTEN) and src homology 2 (SH2) domain containing inositol phosphatase (SHIP) 1/2) have been detected in the nucleus. In particular, the PI3K catalytic subunit p110 β harbours a nuclear




Fig. 2. Modes of protein interaction with polyphosphoinositides in the nucleus. Schematic representation of different modes of PPIn interaction in the following nuclear proteins in **A**) ligand binding domain (LBD) of steroidogenic factor-1 (SF-1) shielding the acyl chains of a representative PPIn (Blind *et al.*, 2014), **B**) ErbB3 binding protein-1 (EBP1) showing the human sequence of the C-terminal unstructured region including the K/R motif (Karlsson *et al.*, 2016) and **C**) Inhibitor of growth 2 (ING2), showing the human sequence of the plant homeodomain (PHD) with the polybasic region (PBR) (Gozani *et al.*, 2003; Huang *et al.*, 2007). Hydrophobic regions/interactions are indicated in orange. Basic residues are shown in blue. Basic residues which have been mutated and implicated in binding are indicated with +. Illustrations are not drawn to scale.

localisation signal (NLS) and was shown to be important for DNA replication, DNA repair and cell survival (Kumar et al., 2010, 2011; Marques et al., 2009). However, some of the reported nuclear functions of p110 β have been shown to be kinase-independent (Kumar et al., 2010; Marques et al., 2009). Regarding a possible nucleolar role for p110 β , one study showed that p110 (likely β) interacts, phosphorylates and activates the ribosomal DNA transcription factor, upstream binding factor (UBF), in the nucleus upon insulin growth factor-1 stimulation (Drakas et al., 2004). This study did not however evaluate the lipid kinase activity of p110(β). Alternatively, inositol polyphosphate multikinase (IPMK), found in the nucleus, has PI3K activity (Maag et al., 2011; Resnick et al., 2005) and was shown to phosphorylate PtdIns(4,5) P_2 -bound to steroidogenic factor-1 (SF-1) to PtdIns(3,4,5) P_3 (Blind et al., 2012). The phosphatases PTEN and SHIP1/2, which dephosphorylate PtdIns(3,4,5) P_3 to produce PtdIns(4,5) P_2 and PtdIns(3,4) P_2 respectively, are also present in the nucleus (Deleris et al., 2003; Ehm et al., 2015; Elong Edimo et al., 2011; Shen et al., 2007). SHIP2 has been found in nuclear speckles when phosphorylated on Serine 132 and suggested to use PtdIns(4,5) P_2 as a substrate rather than PtdIns(3,4,5) P_3 *in vivo* (Deleris et al., 2003; Elong Edimo et al., 2011). The nuclear roles of PTEN are thought to be either dependent (Liu et al., 2005) or independent of its lipid phosphatase activity (Lindsay et al., 2006; Shen et al., 2007; Song et al., 2011). PTEN has also been found in promyelocytic leukaemia protein (PML) bodies, at the centromere and in nucleoli (Li et al., 2014; Shen et al., 2007; Song et al., 2008). The class II PI3Ks α and β , which can produce PtdIns3P or PtdIns(3,4) P_2 , are found in nuclear speckles or in the nuclear matrix in association with lamin A/C, respectively (Banfic et al., 2009; Didichenko and Thelen, 2001; Sindic et al., 2001).

One fundamental question about the presence of PPIIn within the nucleus is how their hydrophobic tails are shielded from the aqueous environment. Although our understanding of the biophysical nature of these PPIIn in the nucleus remains incomplete, several possibilities have been proposed. Recent elegant structural studies demonstrated the acyl chains to be buried in the hydrophobic ligand binding pocket of the nuclear receptors (NR) SF-1/NR5A1 ((Blind et al., 2012, 2014), as represented in Fig. 2A) and liver receptor homolog-1 (LRH-1/NR5A2) (Sablin et al., 2015), allowing the exposure of the inositol head group to the action of PPIIn kinases and phosphatases (Blind et al., 2012). Alternatively, the formation of micelles, perhaps with the abundant nuclear phospholipid phosphatidylcholine (Hunt et al., 2001), has been suggested but not demonstrated so far (Barlow et al., 2010; Irvine, 2006). Recently, a new type of nuclear structure referred to as nuclear lipid islets (NLIs) has been described as PtdIns(4,5) P_2 nuclear aggregates, possibly in the form of micelles with acyl chains facing inwards (Sobol et al., 2018; Sztacho et al., 2019). These structures are suggested to serve as platforms for RNA polymerase II (pol II) as an integral part for mRNA pol II transcription. Lipid droplets have also been documented in the nucleus (Layerenza et al., 2013) as well as in expansions of the inner nuclear membrane (Romanauska and Kohler, 2018) and may act as reservoir of PPIIn in their monolayer. In whatever physico-chemical structure PPIIn may be found in the nucleus, another issue is still unanswered: how is PtdIns transferred from its site of synthesis in the endoplasmic reticulum to the nucleus for the local action of PPIIn kinases and phosphatases? Answering this question will in no doubt give the field of nuclear PPIIn greater recognition.

3. Nuclear polyphosphoinositide functions

Elucidating interaction partners for nuclear PPIIn is fundamental to understanding the role they play in the nucleus. Several studies have begun to shed light on nuclear PPIIn effector proteins and the influence PPIIn have on diverse key nuclear processes.

3.1. Nuclear PPIIn in chromatin remodelling and gene transcription

PPIIn have been linked to several aspects of chromatin remodelling and regulation of gene transcription. PPIIn modulating the interaction of PPIIn-binding proteins with chromatin is one potential mechanism by which PPIIn influence these processes (Viiri et al., 2012). Some of these interactions are regulated via PtdIns5P, which although normally found at low levels in cells, increases in the nucleus in response to different types of cellular stress (Clarke et al., 2001; Jones et al., 2006; Zou et al., 2007). PtdIns5P binds to several nuclear proteins involved in chromatin remodelling harbouring a plant homeodomain (PHD)-type zinc finger. PHD zinc fingers can bind to methylated lysines in histone tails while other parts of the protein can recruit chromatin-remodelling complexes (Sanchez and Zhou, 2011). One of these proteins is the histone code reader, inhibitor of growth protein 2 (ING2), a component of the transcriptional co-repressor Sin3a-histone deacetylase 1 (HDAC1) complex (Bua et al., 2013; Gozani et al., 2003). PtdIns5P contributes to the localisation of ING2 in the nucleus where it associates with the chromatin (Gozani et al., 2003; Jones et al., 2006). This is indeed counteracted by the overexpression of the type II PtdIns5P 4-kinase β , which decreases the levels of PtdIns5P. Overexpression of ING2 induces p53 acetylation and p53-dependent induction of apoptosis and this activity was shown to be dependent upon an intact PPIIn-binding motif (Gozani et al., 2003; Zou et al., 2007). In response to etoposide-induced DNA damage in particular, PtdIns5P was demonstrated to be required for the association of ING2 with target gene promoters leading to their transcriptional repression (Bua et al., 2013). Interaction of all three mono-phosphorylated PPIIn with the co-repressor sin3A-associated protein 30-like (SAP30L) was reported to decrease its association with chromatin *in vitro* and to reduce its transcriptional repression activity (Viiri et al., 2009). Association of the basal transcription initiation factor TFIID subunit 3 (TAF3) with PPIIn, including PtdIns5P, was shown to be required for TAF3-regulated gene transcription necessary for muscle differentiation (Stijf-Bultsma et al., 2015). In plants, the extended PHD finger of *Arabidopsis* homolog of trithorax ATX1 binds selectively to PtdIns5P, which leads to its translocation from the nucleus to the cytoplasm, and a decrease in the transcriptional activity of ATX1 (Alvarez-Venegas et al., 2006; Ndamukong et al., 2010). Furthermore, interaction of Ubiquitin-like PHD and RING finger domain-containing protein 1 (UHRF1) with PtdIns5P leads to a conformational change allowing binding of UHRF1 to H3K9me3 (Gelato et al., 2014).

Other nuclear factors are influenced by PtdIns(4,5) P_2 . The role of this PPIIn is associated with chromatin remodelling as it allows the retention of BRG1, a component of the SWI/SNF-like BRG/BRM-associated factors (BAF) chromatin-remodelling complex in the

nucleus and in particular on the chromatin (Zhao *et al.*, 1998). PtdIns(4,5) P_2 binds to a basic motif in the C-terminal part of BRG1 and enhances actin filament binding (Rando *et al.*, 2002). The functional consequence of this interaction is, however, not clear. What is known is that actin plays roles in transcription (Visa and Percipalle, 2010) and may hence influence the action of the BAF complex, and this, in turn, may be regulated by PtdIns(4,5) P_2 .

Association of the myristoylated corepressor brain acid soluble protein 1 (BASP1) with PtdIns(4,5) P_2 is a requisite for HDAC1 binding at gene promoters regulated by Wilms' tumour 1 protein, and the subsequent repression of these genes (Toska *et al.*, 2012). The nuclear receptors SF-1 (Blind *et al.*, 2014) and LRH-1 (Sablin *et al.*, 2015) interact with PPIIn via their hydrophobic ligand-binding pockets while leaving the inositol head group exposed to possible modification (Fig. 2A). Interestingly, the phosphorylation status of the PPIIn associated with SF-1 has been shown to be critical for transcriptional activity, with phosphorylation of PtdIns(4,5) P_2 to PtdIns(3,4,5) P_3 by IPMK associated with higher transcriptional activity and dephosphorylation by PTEN to PtdIns(4,5) P_2 with reduced activity (Blind *et al.*, 2012). Alternatively, histones H1 and H3 themselves have also been identified as PtdIns(4,5) P_2 -binding proteins and PtdIns(4,5) P_2 was identified to interact with the C-terminal of H1 (Yu *et al.*, 1998). The authors further demonstrate that PtdIns(4,5) P_2 can alleviate the inhibition of mRNA pol II transcription imposed by H1 *in vitro*. Most recently, Choi *et al.* have shown that both nuclear PtdIns(4,5) P_2 and PIPKI α interact with the tumour suppressor p53, which increases its stability during cellular stress (Choi *et al.*, 2019).

3.2. PPIIn in nucleolar processes

Several PPIIn species have been detected in the nucleolus, including PtdIns3P, PtdIns(4,5) P_2 , PtdIns(3,4,5) P_3 and perhaps also PtdIns4P (Gillooly *et al.*, 2000; Kalasova *et al.*, 2016; Karlsson *et al.*, 2016; Osborne *et al.*, 2001; Sobol *et al.*, 2013; Yildirim *et al.*, 2013). PtdIns(4,5) P_2 was shown to bind to RNA pol I and to be necessary for transcription at the rDNA promoter (Yildirim *et al.*, 2013). In addition, PtdIns(4,5) P_2 binds both to UBF and fibrillarin, which resulted in a reduction of binding of UBF to rDNA and altered mobility of the fibrillarin/snRNA complex, suggesting a possible competitive mode of action. Furthermore, PtdIns(4,5) P_2 engages with Pol I subunits and UBF during mitosis and co-localisation is maintained even when Pol I or II transcription is inhibited, suggesting that PtdIns(4,5) P_2 is part of the nucleolar organising region (NOR) complex of which the organisation is cell cycle dependent (Sobol *et al.*, 2013). Recently, PtdIns(4,5) P_2 has been shown to regulate the levels of H3K9me2 at the rDNA promoter by binding to and inhibiting the activity of the histone lysine demethylase PHD finger protein 8 (PHF8), likely resulting in a reduction in rRNA gene transcription (Ulicna *et al.*, 2018). Consistent with these studies showing a role for PtdIns(4,5) P_2 in nucleolar processes, PIPKI α , one of the PPIIn kinases generating PtdIns(4,5) P_2 using PtdIns4P, was found in the nucleolus in the G1/S phase of the cell cycle (Chakrabarti *et al.*, 2015). PIPKI α associates with H3K9me3 and plays roles in rDNA gene silencing. Whether or not the action of PIPKI α is due to its lipid kinase activity was not investigated, but in light of the study of Ulicna *et al.*, this mechanism would be worth exploring.

3.3. Nuclear PPIIn and cell survival

Nuclear PI3K activity and PtdIns(3,4,5) P_3 have been shown to mediate the anti-apoptotic response to nerve growth factor (NGF) together with the effector protein, nucleophosmin (alias B23) (Ahn *et al.*, 2004, 2005). Upon the addition of NGF, PtdIns(3,4,5) P_3 regulates the interaction of nucleophosmin with caspase-activated DNase (CAD), thereby preventing its nuclease activity and DNA fragmentation (Ahn *et al.*, 2005). The interaction between PtdIns(3,4,5) P_3 and nucleophosmin in particular was required for the prevention of cell death. Additionally, nucleophosmin associates with active Akt in the nucleus following NGF stimulation, which increases its stability and enhances cell survival (Lee *et al.*, 2008). The GTPase phosphatidylinositol 3 (PI3)-kinase enhancer (PIKE)-L (now known as Arf-GAP with GTPase, ANK repeat and PH domain-containing protein 2), is another nuclear protein which binds to PtdIns(3,4,5) P_3 , and contributes to cell survival through activation of PI3K and Akt (Hu *et al.*, 2005).

3.4. PPIIn in mRNA processing and export

Nuclear speckles (also known as interchromatin granule clusters) are nuclear hubs in which pre-mRNA splicing factors congregate, but also contain a variety of other proteins (Galganski *et al.*, 2017; Spector and Lamond, 2011). Localisation of PtdIns(4,5) P_2 to nuclear speckles was evident with its co-detection with speckle markers such as the splicing factor SC-35 or Sm proteins (Boronenkov *et al.*, 1998; Osborne *et al.*, 2001). Nuclear PtdIns(4,5) P_2 interacts with components of small nuclear ribonucleoproteins (RNP), the hyperphosphorylated form of the large subunit of RNA pol II and U1-U6 snRNAs (Osborne *et al.*, 2001). The same study also showed that its immunodepletion inhibited splicing of pre-mRNAs *in vitro*. Other nuclear speckle-associated proteins are also known to interact with PPIIn. Aly (alias THO complex subunit 4), for example, can associate with PtdIns(4,5) P_2 and PtdIns(3,4,5) P_3 and mutation of key basic residues prevented PtdIns(3,4,5) P_3 -binding and reduced Aly-mediated mRNA export activities and cell proliferation (Okada *et al.*, 2008). Moreover, PtdIns(3,4,5) P_3 , produced by IPMK, was shown to regulate Aly-mediated recognition of specific mRNAs for nuclear export, particularly for those involved in homologous recombination DNA repair (Wickramasinghe *et al.*, 2013). Furthermore, a study searching for nuclear proteins interacting with the nuclear PIPKI α , identified the non-canonical poly(A) polymerase called nuclear speckle targeted PIPKI α regulated-poly(A) polymerase (Star-PAP), and showed that PtdIns(4,5) P_2 directly stimulates the polyadenylation activity of target mRNAs (Mellman *et al.*, 2008). Further studies from the same group demonstrated that casein kinase I α (CKI α), which is also PtdIns(4,5) P_2 -sensitive, is part of the Star-PAP complex together with PIPKI α and can regulate Star-PAP activity targeting the 3'-end mRNA processing of select mRNAs (Gonzales *et al.*, 2008).

Collectively, these studies clearly demonstrate the far-reaching influence that PPIIn have on nuclear events via their interaction with an array of different effector proteins.

4. Mapping polyphosphoinositide-protein interactomes to understand their function

4.1. PPIIn interactomic studies

To globally capture PPIIn binding proteins, multiple studies have taken a proteomic approach ((Best, 2014; D'Santos and Lewis, 2012) and listed in Table 3). Affinity matrices containing synthetic PPIIn followed by mass spectrometric fingerprinting have been used to isolate PtdIns(3,4,5) P_3 -binding proteins in pig leukocytes (Krugmann et al., 2002). A chemical proteomics method using cleavable lipid baits has also been used in macrophage whole cell extracts to identify mostly PtdIns(3,4) P_2 interacting proteins (Pasquali et al., 2007). Other studies using PPIIn pull downs to identify interacting proteins included bifunctional activity-based probes. In this approach, target proteins were labelled using PPIIn head groups containing both a photo affinity tag to bind the proteins and a secondary tag for analysis (Rowland et al., 2011). With the advancement of mass spectrometry, more quantitative methods have been employed. Bead immobilized or liposomes incorporated PPIIn have been used to identify PtdIns3P, PtdIns(3,5) P_2 , PtdIns(4,5) P_2 and PtdIns(3,4,5) P_3 target proteins in colorectal carcinoma cytosolic extracts (Catimel et al., 2008, 2009, 2013). To discriminate specific PPIIn binders from background proteins, quantitative approaches such as stable isotope labelling by amino acids in cell culture (SILAC) have been employed (Dixon et al., 2011; Jungmichel et al., 2014; Lewis et al., 2011). Alternative strategies include the analysis of yeast cDNA libraries combined with FACS-dependent identification of fluorescent PPIIn binding clones for the identification of PtdIns(4,5) P_2 and PtdIns(3,4,5) P_3 binding proteins (Bidlingmaier et al., 2011).

4.2. Nuclear PtdIns(4,5) P_2 interactome

To map the nuclear PPIIn-protein interaction networks, we established an approach allowing the enrichment of nuclear PPIIn-interacting proteins by incubating nuclei with the polyamine neomycin (Lewis et al., 2011). Due to the high affinity of neomycin for PPIIn via electrostatic interactions (Gabev et al., 1989; Schacht, 1976, 1978, 1979), neomycin was predicted to compete with proteins for binding to PPIIn. By incubating isolated nuclei with neomycin, PPIIn-binding proteins were displaced, allowing their collection and identification by quantitative mass spectrometry. Using this approach, 349 proteins specifically displaced by neomycin from intact nuclei were identified with enriched functions in DNA topological change, nucleosome positioning/assembly, chromatin assembly, ribosome biogenesis, rRNA processing, mRNA splicing and processing (Lewis et al., 2011). Amongst them, a few known PPIIn-effector proteins were identified such as Aly (Okada et al., 2008), histone H1 (Yu et al., 1998) and nucleophosmin (Ahn et al., 2005) as well as several speckle-associated proteins with RNA splicing functions (Osborne et al., 2001). However, many of the neomycin-displaced proteins had no previous history as nuclear PPIIn-effector proteins and presented potentially novel roles for nuclear PPIIn. PPIIn

Table 3

PPIIn interactomics studies.

PPIIn interactome analysed	Method	Cell type/subcellular compartment	Reference
PtdIns(3,4,5) P_3	PPIIn conjugated beads and MS	Pig leukocyte cytosolic extract	(Krugmann et al., 2002)
Mostly PtdIns(3,4) P_2	PPIIn conjugated to cleavable S-S bond biotin + streptavidin beads and MS	Primary macrophage cytosolic extract	(Pasquali et al., 2007)
PtdIns(4,5) P_2	Biotinylated PPIIn, streptavidine conjugated beads and MS	Secretory granules from bovine adrenal chromaffin cells	(Osborne et al., 2007)
PtdIns(3,5) P_2 & PtdIns(4,5) P_2	PPIIn conjugated beads or liposomes and MS	LIM1215 colon cancer cell cytosolic extract	(Catimel et al., 2008)
PtdIns(3,4,5) P_3	PPIIn conjugated beads or liposomes and MS	LIM1215 colon cancer cell cytosolic extract	(Catimel et al., 2009)
PtdIns(4,5) P_2	PPIIn conjugated beads and quantitative MS	Neomycin extracted nuclear proteins isolated from murine erythroleukemia cells	(Lewis et al., 2011)
PtdIns(3,4) P_2	Stimulation of class I PI3K \pm wortmannin, biotinylated PPIIn coupled to streptavidin beads and SILAC-based quantitative MS	1321N1 astrocytoma membrane fractions	(Dixon et al., 2011)
PtdIns(4,5) P_2 and PtdIns(3,4,5) P_3	yeast surface-displayed human protein fragment libraries + fluorescent PPIIn	N/A	(Bidlingmaier et al., 2011)
PtdIns(3,4,5) P_3	PPIIn activity probe for click chemistry and MS	MDA-MB-435 cancer cell cytosolic extract	(Rowland et al., 2011)
PtdIns3P	PPIIn conjugated beads or liposomes and MS	LIM1215 colon cancer cell cytosolic, membrane and nuclear extracts	(Catimel et al., 2013)
All PPIIn	Triple SILAC-based quantitative MS	HeLa S3 whole cell extract	(Jungmichel et al., 2014)

Table updated from (D'Santos and Lewis, 2012). SILAC: stable isotope labeling by amino acids in cell culture; MS: mass spectrometry; N/A: not applicable.

binding of DNA topoisomerase II α (Topo II α), for example, was validated and an inhibitory effect of PPI on kDNA decatenation activity was demonstrated, suggesting a role for PPI in the regulation of DNA topological changes (Lewis et al., 2011). Other neomycin-extracted proteins were subsequently verified to interact with PPI and include ErbB3 binding protein-1 (EBP1, alias proliferation-associated protein 2G4, PA2G4) (Karlsson et al., 2016), PHF8 (Ulicna et al., 2018) and UHRF1 (Gelato et al., 2014). Additionally, neomycin extracts were subjected to quantitative interactomics following PtdIns(4,5)P₂ lipid-pull down and this led to the identification of 34 PtdIns(4,5)P₂-binding proteins, 28 of which were annotated to the nucleus, and known to be involved in mRNA transcription regulation, mRNA splicing and protein folding/complex assembly (Lewis et al., 2011). Two of these proteins were subsequently found common to the PtdIns(4,5)P₂-binding proteins identified by the PPI interactomics study by Jungmichel et al. (2014), i.e. Elongation factor 1-delta and TBC1 domain family member 24.

5. Nuclear polyphosphoinositide-protein modes of interaction – polybasic regions lead the way

5.1. Diverse modes of interactions for PPI-binding proteins

Proteins interact with PPI via domains of diverse structure and PPI binding specificities and affinities. They include the

Table 4
Polybasic motifs involved in polyphosphoinositide-protein interaction.

Actin-binding and membrane-binding proteins			
Sequence	Protein	Motif type	Reference
135-KSGLKYKK 161-KLFQVKGRR	Gelsolin	K/R	(Yu et al., 1992)
139-KLYQVKGKK	gCap39 ^a	K/R	(Yu et al., 1990, 1992)
112-KQGLVIRK 138-RLLVKGGKR 816-KQQLKKEK	Villin	K/R	(Bazari et al., 1988; Kumar et al., 2004)
13-KVFNDMKVRK	Cofilin	K/R	(Matsuzaki et al., 1988; Yu et al., 1992)
127-KCYEMASHLRR	Profilin I	K/R-like	(Lassing and Lindberg, 1985; Skare and Karlsson, 2002; Lambrechts et al., 2002)
151-KKKKKRFSFKKSFKLSGFSFKKNNK	MARCKS	K/R-like	(Wang et al., 2001)
30-KAHKAAIKIQASFRGHITRKKLKGEEK	GAP43 ^b	K/R-like	(Laux et al., 2000)
183-KEKKKGAKKKRLTK	N-WASP	PBR	(Papayannopoulos et al., 2005)
Nuclear PPI-binding proteins			
Sequence	Protein	Motif type	Reference
105-RRKKREQKK	Pf1	K/R, PBR	(Kaadige and Ayer, 2006)
123-RNRRKRK	SAP30	PBR	(Viiri et al., 2009)
85-RNKRKRK	SAP30L	PBR	(Viiri et al., 2009)
264-KTMDKSTKTKKDRRSR	ING2	PBR	(Gozani et al., 2003)
1-60 region including a polybasic motif ^c	BASP1 (alias NAP-22)	PBR	(Terashita et al., 2002)
641-RTGKGWKRKSAGGGPS	UHRF1	PBR	(Gelato et al., 2014)
918-IKDKKHKHKKRHRAH	TAF3	PBR	(Stijf-Bultsma et al., 2015)
27-RGRGR and 79-RPK	Aly	RG motif	(Okada et al., 2008)
65-KKEKEMKK 364-RKTQKKKKKK	EBP1 ^d EBP1 ^d	Reverse K/R K/R	(Karlsson et al., 2016)
PHD-BRD linker ^e	BAZ2B	K/R	(Kostrhon et al., 2017)
827-KSRPKKKK	PHF8	K/R, PBR	(Ulicna et al., 2018)
K194, K200 and K228, K234	NPM	PBR?	(Ahn et al., 2005)
981-KKVRGKVVWQRISPLFNTK	OGT	Reverse K/R	(Yang et al., 2008)
1402-EEVRQKKSSRKRKRDS	BRG1	K/R, PBR	(Rando et al., 2002)
370-KSKKGQSTSRHKK	p53	PBR	(Choi et al., 2019)

Basic residues in bold indicate residues shown to be required for PPI interaction. BASP1: brain acid soluble protein 1, BAZ2B: BROMO domain adjacent to zinc finger 2B, EBP1: Erb 3 binding protein-1, GAP43: Growth Associated Protein 43, ING2: inhibitor of growth protein 2, NPM: nucleophosmin, OGT: O-linked beta-N-acetylglucosamine transferase, Pf1: PHD factor 1, PHF8: PHD finger protein 8, N-WASP: neuronal Wiskott-Aldrich syndrome protein, SAP30: Sin3A-associated protein 30, SAP30L: human SAP30-like protein, TAF3: Transcription initiation factor TFIID subunit 3, UHRF1: E3 ubiquitin-protein ligase UHRF1. K/R: lysine/arginine motif following the motif K/R-(X_{3,7})-K-X-K/R-K/R, PBR-polybasic region.

^a Recommended protein name: macrophage-capping protein.

^b Recommended protein name: neuromodulin.

^c The basic motif lies between aa 4–10 (1-MGGKLSKKKKGYNVN) of the first 60 aa shown to bind to PPI.

^d Isoform 1 also known as p48.

^e PHD-BRD linker:1982-ASGQTLKIKLHVKGKKTNESKGGKVVTLTGDTEDSASTSSSLKRGKDKLKKRMEENTSINLSKQESFT.

following domains: PH, FYVE, PX (Phox homology), ENTH/ANTH (epsin amino-terminal homology), FERM (band 4.1, ezrin, radixin and moesin), PROPPIN and TRAF (Choy *et al.*, 2017; Hammond and Balla, 2015; Kutateladze, 2010; Pemberton and Balla, 2018). Alternatively, short polybasic PPIIn-binding motifs or regions (PBR) have also been reported to interact with PPIIn (Janmey *et al.*, 2018; McLaughlin and Murray, 2005; Yin and Janmey, 2003). These motifs consist of short stretches of basic amino acids, often denoted as lysine/arginine-rich patches/clusters or KR-motifs (K/R-(X₃₋₇)-K-X-K/R-K/R), first identified in the actin-binding protein (ABP) gelsolin (Martin, 1998; Yu *et al.*, 1992). Gelsolin is a multifunctional protein involved in cytoskeletal remodelling by severing and capping actin filaments. This activity is regulated by PtdIns(4,5)P₂ interacting at the same site as the actin-binding site (Feng *et al.*, 2001; Yu *et al.*, 1992). KR-motifs have been identified in many PPIIn-binding cytoskeletal proteins of the gelsolin family, such as gCap39, villin, cofilin and profilin and the Wiskott Aldrich syndrome protein (WASP) ((Janmey *et al.*, 2018) and Table 4). These motifs rely on electrostatic interactions between their basic residues and the negatively charged phosphate groups. In addition, hydrophobic residues can also contribute to the interaction to the acyl chains of PPIIn, as shown for gelsolin and cofilin (Feng *et al.*, 2001; Gorbatyuk *et al.*, 2006; Janmey and Stossel, 1987). Some of these ABPs known to engage in PtdIns(4,5)P₂ interaction in the cytoplasm can also be found in the nucleus (reviewed in (Barlow *et al.*, 2010)). For example, profilin I, which binds to PtdIns(4,5)P₂ via two motifs, one of them overlapping with the actin binding site, is also present in nuclear speckles as well as cajal bodies, and has been implicated in mRNA splicing (Lambrechts *et al.*, 2002; Skare and Karlsson, 2002; Skare *et al.*, 2003). Other ABPs, such as N-WASP (Wu *et al.*, 2006), gCAP39 (Onoda and Yin, 1993; Onoda *et al.*, 1993), as well as cofilins (Munsie *et al.*, 2012), have also been detected in the nucleus in addition to the cytoplasm. To the best of our knowledge, the functional role(s) of the interaction with PPIIn have however not been explored in the regulation of nuclear processes thought to be mediated by these proteins.

Another well characterized example of PPIIn-binding motif can be found in the myristoylated alanine-rich C-kinase substrate (MARCKS) and growth-associated protein 43 (GAP43) proteins ((Laux *et al.*, 2000; Wang *et al.*, 2002) see Table 4). MARCKS and GAP43 anchor into the bilayer through an N-terminal myristate and two saturated acyl chains respectively, and bind electrostatically to a cluster of PtdIns(4,5)P₂ via a conserved effector domain, consisting of a K/R like PBR. The membrane interaction of MARCKS is regulated by phosphorylation of serine residues located within the effector domain by protein kinase C, resulting in a reduction of the electrostatic attraction between the protein and PPIIn and causing MARCKS to dissociate from the plasma membrane and to translocate to the cytoplasm (McLaughlin and Murray, 2005; Ohmori *et al.*, 2000). Interestingly, MARCKS has also been detected in the nucleus due to its effector domain acting as an NLS (Rohrbach *et al.*, 2015).

Many small GTPases in the Ras and Rho families harbour C-terminal PBR consisting of four or more lysine or arginine residues, responsible for their association with the plasma membrane via PtdIns(4,5)P₂ and PtdIns(3,4,5)P₃ (Heo *et al.*, 2006). In the case of the small GTPase Rac1, the polybasic motif was demonstrated to also act as a functional NLS (Lanning *et al.*, 2003). The presence of canonical NLS sequences (K-K/R-x-K/R) in the polybasic motif of several Ras and Rho family members indicates a dual regulatory role in membrane binding and nucleo-cytoplasmic shuttling of these proteins (Williams, 2003).

5.2. Polybasic motifs as a mechanism for nuclear protein-PPIIn interaction

In the case of *true* nuclear proteins, *i.e.* residing dominantly in the nucleus, only a few are known to interact with PPIIn. via a structured domain. For example, basic residues in the PH domain of Tfb1 (p62), a subunit of the general transcription factor II H (TFIIH) interacts preferably with PtdIns3P and PtdIns5P, and interaction with PtdIns5P could potentially regulate its binding with the activation domain of VP16 due to an overlapping interaction site (Di Lello *et al.*, 2005). PtdIns(3,4,5)P₃ binding protein (PIP₃BP) binds to PtdIns(3,4,5)P₃ via two individual PH domains (Tanaka *et al.*, 1997, 1999). Consistently, in our study, only seven of the 349 proteins identified following neomycin displacement contained structured PPIIn-binding domains, whereas 165 contained at least one basic PPIIn-binding K/R motif (Lewis *et al.*, 2011). Furthermore, 19 of the 28 nuclear PtdIns(4,5)P₂ specific-binding proteins contained these motifs. Using mutational studies, we have subsequently verified the involvement of PBR or K/R motifs in PPIIn binding in three proteins identified, namely EBP1 (Karlsson *et al.*, 2016), heterogeneous nuclear ribonucleoprotein U (hnRNP U) and Topo II α (unpublished results). EBP1 harbours two K/R motifs, one localised in the unstructured C-terminus and a reversely orientated K/R motif in the N-terminus on a protruding loop (Table 4 (Karlsson *et al.*, 2016)). Both of these K/R motifs contributed to PPIIn binding as site-directed mutations of lysine residues to alanines in both motifs were required to prevent PPIIn binding of the full length protein (Karlsson *et al.*, 2016). The contribution of K/R motifs or PBRs in PPIIn binding has also been demonstrated in a number of other nuclear proteins, hence underscoring the relevance of electrostatic interactions for PPIIn-binding in the nucleus. These include ING2 (Bua *et al.*, 2013; Gozani *et al.*, 2003), PHD factor 1 (Pfl) (Kaadige and Ayer, 2006), SAP30L (Viiri *et al.*, 2009), UHRF1 (Gelato *et al.*, 2014), TAF3 (Stijf-Bultsma *et al.*, 2015), BAZ2B (Kostrhon *et al.*, 2017), PHF8 (Ulicna *et al.*, 2018) as well as p53 (Choi *et al.*, 2019) and all require basic residues for the interaction (see Table 4). For example, PHF8 contains two K/R motifs and one PHD, but only the most C-terminal K/R motif was found to bind to PtdIns(4,5)P₂. This binding was abolished by mutating all the lysine residues to alanine or glycine within the motif (Ulicna *et al.*, 2018). In the case of Pfl, a PBR located C-terminal of a PHD zinc finger was shown to be essential for PPIIn binding while the PHD showed little contribution (Kaadige and Ayer, 2006). Basic residues have also been shown to be important for PPIIn-binding in other nuclear proteins such as nucleophosmin, which binds via two lysine pairs within its C-terminus but in the absence of defined PPIIn-binding motif (Ahn *et al.*, 2005). Another example is Aly, which binds via basic residues in a glycine-arginine-rich domain in its N-terminal (Okada *et al.*, 2008). O-linked beta-N-acetylglucosamine transferase (OGT), on the other hand, involves two lysines within a reverse K/R motif (Yang *et al.*, 2008).

In addition to electrostatic properties of the PBR, other features or structures can influence PPIIn interaction with nuclear proteins. The contribution of hydrophobic regions is one of these features as shown in a number of nuclear PtdIns(3,4,5)P₃-binding proteins, as this permits interaction with acyl chains (Bidlingmaier *et al.*, 2011). NMR analysis of the C-terminal PBR of EBP1 showed that

hydrophobic interactions between non-polar residues upstream of the motif, particularly ³⁵⁶ALL³⁵⁸, and acyl chains of 16 carbons contributed to binding to PtdIns(3,4,5)P₃ in addition to electrostatic interactions ((Karlsson et al., 2016), as illustrated in Fig. 2B). Consistently, a hydrophobic region upstream of the PBR is required for the ability of SAP30L to bind monophosphorylated PPIIn, as the PBR alone was unable to bind PPIIn unless this hydrophobic region was also present (Viiri et al., 2009). In one instance, binding of the repressor BASP1 to PtdIns(4,5)P₂ requires the covalent binding of a myristoyl group within the region required for PPIIn interaction (Terashita et al., 2002; Toska et al., 2012). Alternatively, PBR-mediated PPIIn interaction can be influenced by the presence of a zinc finger of the PHD type, located N-terminal to the PBR. This is the case for ING2 as shown in Fig. 2C, as well as in ING1, ACF1 (alias BAZ1A), and TAF3, implying a common mechanism (Gozani et al., 2003; Stijf-Bultsma et al., 2015). Molecular modelling of the PHD-PBR of ING2 with PtdIns5P showed that, in at least one of the models, the PBR forms a PPIIn binding pocket which encases the inositol ring while one of the acyl chains is oriented towards the PHD ((Huang et al., 2007), as illustrated in Fig. 2C). The basic linker region between the PHD domain and the bromodomain of BROMO domain adjacent to zinc finger 2B (BAZ2B) interacts with PPIIn via different K/R motifs and via the contribution of the proximal part of the PHD (Kostrhon et al., 2017). Similarly, SAP30L, display an atypical zinc-finger followed by a PBR, which has also been recognised as a PPIIn-interacting module, particularly to monophosphorylated PPIIn (Viiri et al., 2009). Interestingly, the zinc finger was shown to support PPIIn interaction mediated by the PBR while the PBR was required for PPIIn specificity. Again, the PBR of Pf1 and ING2 play a major role in direct PPIIn binding particularly in determining PPIIn specificity (Kaadige and Ayer, 2006). These studies emphasise that PBRs are key to PPIIn binding and the environment and features around these regions can sometime influence binding, perhaps by shielding acyl chains.

As shown in Table 4, it is interesting to notice that many reported nuclear PPIIn-binding proteins harbour K/R motifs that were initially shown as a binding site for both PtdIns(4,5)P₂ and actin in ABP. As discussed earlier, BRG1 is an example of a nuclear protein which harbours such a motif showing the same binding activity to PtdIns(4,5)P₂ and actin (Rando et al., 2002). Considering the accumulating roles of both PPIIn and actin in nuclear processes, it may be worth evaluating both their contribution when characterising the functions of PPIIn-effector proteins in the nucleus.

5.3. Influence of PPIIn-mediated interaction on nuclear localisation

A few different studies have indicated that PPIIn can influence the nuclear localisation of some PPIIn-interacting proteins due to the presence of basic residues in structured domains or short polybasic motifs. In particular, K/R motifs and PBRs bear resemblance to NLS, i.e. the bipartite NLS (KR-(X₉₋₂₉)-KKK and monopartite K-K/R-X-K/R, and nucleolar localisation signals (NoLS), such as R/K-R/K-X-R/K (Emmott and Hiscox, 2009; McLane and Corbett, 2009). For example, the C-terminal K/R motif of EBP1 overlaps with a NoLS and was shown to be required for both nucleolar targeting and PPIIn binding via four lysines (Karlsson et al., 2016). The PBR in SAP30L and BASP1 serves both as a NLS and PPIIn binding site (Terashita et al., 2002; Toska et al., 2012; Viiri et al., 2009). The PPIIn binding site within the PH domain of the nuclear protein PIKE-L is also a putative NLS (Hu et al., 2005). Incidentally, disruption of PPIIn binding can translocate nuclear effector proteins from the nucleus to the cytoplasm. Mutation of the PPIIn binding site in the PH domain of PIKE-L causes its translocation to the cytoplasm (Hu et al., 2005). Mutation of the PBR which binds PtdIns(4,5)P₂ in the C-terminal of p53 also caused an increased distribution of p53 to the cytoplasm (Choi et al., 2019). As mentioned earlier, decreased levels of PtdIns5P releases ING2 from the chromatin leading to its translocation to the cytoplasm, suggesting that the localisation of ING2 is dependent upon PtdIns5P (Gozani et al., 2003; Jones et al., 2006). This was further substantiated by the effect of synthetic PtdIns5P, as well as other PPIIn, on the sub-nuclear targeting of fluorescently labelled NLS-PHD finger of ING2, which was altered in the wild type but not in the PPIIn binding mutant (Gozani et al., 2005). Similarly, PIP₃BP, which harbours two PH domains, both necessary for PtdIns(3,4,5)P₃ binding, is translocated from the nucleus to the cytoplasm upon the introduction of constitutively active PI3K, which is prevented by loss of the PH domain binding to PPIIn (Tanaka et al., 1997, 1999). Another protein, OGT translocates from the nucleus to the plasma membrane upon serum stimulation in a PtdIns(3,4,5)P₃-dependent manner (Yang et al., 2008).

Disruption of PPIIn binding can also affect sub-nuclear localisation. For example, alteration of the basic motif of Aly prevents its localisation to nuclear speckles, redistributing it to the nucleoplasm (Okada et al., 2008). Insulin receptor substrate 3 (IRS3) can localise both to the plasma membrane and to nuclear speckles and contains a PH domain which binds mostly to PtdIns3P, in contrast to the PH domains of IRS1 and 2 which demonstrate different PPIIn specificities (Maffucci et al., 2003; Razzini et al., 2000). A point mutation in the PH domain, which abolishes PtdIns3P binding, decreases the localisation of full-length IRS3 to both the plasma membrane and the nuclear speckles (Maffucci et al., 2003). PDZ (postsynaptic density protein-95, discs large, zonula occludens) domains, which can bind to PPIIn, including PtdIns(4,5)P₂, have been suggested to localise zonula occludens 2 (ZO-2) to nuclear speckles (Meerschaert et al., 2009) and syntenin-2 to nuclear speckles but most likely to nucleoli (Geeraerts et al., 2013; Mortier et al., 2005). Finally, the pericentric heterochromatin localisation of UHRF1 is dependent upon an intact PPIIn-binding PBR, at least due to PtdIns5P (Gelato et al., 2014).

These studies may implicate a role for nuclear PPIIn to compete with interacting partners or induce conformation changes of effector proteins to allow their redistribution within the nucleus or even as far as to the cytoplasm and on the plasma membrane. The fact that PPIIn can target the same protein on membranes as well as in membrane-less sub nuclear sites is, however, a paradox but appears to require similar binding strategies, as found in peripheral proteins and ABPs.

6. Conclusions

Mounting evidence have clearly shown that, just as cytoplasmic PPIIn, nuclear PPIIn can provide a signalling code allowing the regulation of diverse nuclear processes. Still, many questions are unanswered regarding their functional characterisation. Continuous

research efforts focusing on mapping the effector proteins of all known nuclear PPIs as well as deciphering their mode of interaction are needed to increase our understanding of PPI signaling in the nucleus.

Conflicts of interest

The authors declare to have no conflict of interest with the publication of this manuscript.

Acknowledgements

Research in our laboratory is funded by the University of Bergen, the Meltzer Research Fund, the Nansen Fund and the Norwegian Cancer Society (grant number 2183087). We thank Vandana Ardawatia, Diana C. Turcu and Andrea Papdiné Morovicz for their critical reading of the manuscript and helpful discussions.

References

- Ahn, J.Y., Liu, X., Cheng, D., Peng, J., Chan, P.K., Wade, P.A., Ye, K., 2005. Nucleophosmin/B23, a nuclear PI(3,4,5)P(3) receptor, mediates the antiapoptotic actions of NGF by inhibiting CAD. *Mol. Cell.* 18 (4), 435–445.
- Ahn, J.Y., Rong, R., Liu, X., Ye, K., 2004. PIKE/nuclear PI 3-kinase signaling mediates the antiapoptotic actions of NGF in the nucleus. *EMBO J.* 23 (20), 3995–4006.
- Alvarez-Venegas, R., Sadder, M., Hlavacka, A., Baluska, F., Xia, Y., Lu, G., Firsov, A., Sarath, G., Moriyama, H., Dubrovsky, J.G., Avramova, Z., 2006. The Arabidopsis homolog of trithorax, ATX1, binds phosphatidylinositol 5-phosphate, and the two regulate a common set of target genes. *Proc. Natl. Acad. Sci. U. S. A.* 103 (15), 6049–6054.
- Balla, T., 2013. Phosphoinositides: tiny lipids with giant impact on cell regulation. *Physiol. Rev.* 93 (3), 1019–1137.
- Banfic, H., Visnjic, D., Mise, N., Balakrishnan, S., Deplano, S., Korchev, Y.E., Domin, J., 2009. Epidermal growth factor stimulates translocation of the class II phosphoinositide 3-kinase PI3K-C2beta to the nucleus. *Biochem. J.* 422 (1), 53–60.
- Barlow, C.A., Laishram, R.S., Anderson, R.A., 2010. Nuclear phosphoinositides: a signaling enigma wrapped in a compartmental conundrum. *Trends Cell Biol.* 20 (1), 25–35.
- Bazari, W.L., Matsudaira, P., Wallek, M., Smeal, T., Jakes, R., Ahmed, Y., 1988. Villin sequence and peptide map identify six homologous domains. *Proc. Natl. Acad. Sci. U. S. A.* 85 (14), 4986–4990.
- Best, M.D., 2014. Global approaches for the elucidation of phosphoinositide-binding proteins. *Chem. Phys. Lipids* 182, 19–28.
- Bidlingmaier, S., Wang, Y., Liu, Y., Zhang, N., Liu, B., 2011. Comprehensive analysis of yeast surface displayed cDNA library selection outputs by exon microarray to identify novel protein-ligand interactions. *Mol. Cell. Proteomics* 10 (3) M110 005116.
- Blind, R.D., Sablin, E.P., Kuchenbecker, K.M., Chiu, H.J., Deacon, A.M., Das, D., Fletterick, R.J., Ingraham, H.A., 2014. The signaling phospholipid PIP3 creates a new interaction surface on the nuclear receptor SF-1. *Proc. Natl. Acad. Sci. U. S. A.* 111 (42), 15054–15059.
- Blind, R.D., Suzawa, M., Ingraham, H.A., 2012. Direct modification and activation of a nuclear receptor-PIP(2) complex by the inositol lipid kinase IPMK. *Sci. Signal.* 5 (229), ra44.
- Boronenkov, I.V., Lojans, J.C., Umeda, M., Anderson, R.A., 1998. Phosphoinositide signaling pathways in nuclei are associated with nuclear speckles containing pre-mRNA processing factors. *Mol. Biol. Cell* 9 (12), 3547–3560.
- Bua, D.J., Martin, G.M., Binda, O., Gozani, O., 2013. Nuclear phosphatidylinositol-5-phosphate regulates ING2 stability at discrete chromatin targets in response to DNA damage. *Sci. Rep.* 3, 2137.
- Bultsma, Y., Keune, W.J., Divecha, N., 2010. PIP4Kbeta interacts with and modulates nuclear localization of the high-activity PtdIns5P-4-kinase isoform PIP4Kalpha. *Biochem. J.* 430 (2), 223–235.
- Catimel, B., Kapp, E., Yin, M.X., Gregory, M., Wong, L.S., Condrón, M., Church, N., Kershaw, N., Holmes, A.B., Burgess, A.W., 2013. The PI(3)P interactome from a colon cancer cell. *J. Proteomics* 82, 35–51.
- Catimel, B., Schieber, C., Condrón, M., Patsiouras, H., Connolly, L., Catimel, J., Nice, E.C., Burgess, A.W., Holmes, A.B., 2008. The PI(3,5)P2 and PI(4,5)P2 interactomes. *J. Proteome Res.* 7 (12), 5295–5313.
- Catimel, B., Yin, M.X., Schieber, C., Condrón, M., Patsiouras, H., Catimel, J., Robinson, D.E., Wong, L.S., Nice, E.C., Holmes, A.B., Burgess, A.W., 2009. PI(3,4,5)P3 interactome. *J. Proteome Res.* 8 (7), 3712–3726.
- Chakrabarti, R., Sanyal, S., Ghosh, A., Bhar, K., Das, C., Siddhanta, A., 2015. Phosphatidylinositol-4-phosphate 5-kinase 1 alpha modulates ribosomal RNA gene silencing through its interaction with histone H3 lysine 9 trimethylation and heterochromatin protein HP1-alpha. *J. Biol. Chem.* 290 (34), 20893–20903.
- Choi, S., Chen, M., Cryns, V.L., Anderson, R.A., 2019. A nuclear phosphoinositide kinase complex regulates p53. *Nat. Cell Biol.* 21, 462–475.
- Choy, C.H., Han, B.K., Botelho, R.J., 2017. Phosphoinositide diversity, distribution, and effector function: Stepping Out of the Box. *Bioessays* 39 (12).
- Ciruela, A., Hinchliffe, K.A., Divecha, N., Irvine, R.F., 2000. Nuclear targeting of the beta isoform of type II phosphatidylinositol phosphate kinase (phosphatidylinositol 5-phosphate 4-kinase) by its alpha-helix 7. *Biochem. J.* 346 Pt 3, 587–591.
- Clarke, J.H., Letcher, A.J., D'Santos, C.S., Halstead, J.R., Irvine, R.F., Divecha, N., 2001. Inositol lipids are regulated during cell cycle progression in the nuclei of murine erythroleukaemia cells. *Biochem. J.* 357 (Pt 3), 905–910.
- Cocco, L., Gilmour, R.S., Ognibene, A., Letcher, A.J., Manzoli, F.A., Irvine, R.F., 1987. Synthesis of polyphosphoinositides in nuclei of Friend cells. Evidence for polyphosphoinositide metabolism inside the nucleus which changes with cell differentiation. *Biochem. J.* 248 (3), 765–770.
- Cocco, L., Martelli, A.M., Gilmour, R.S., Ognibene, A., Manzoli, F.A., Irvine, R.F., 1989. Changes in nuclear inositol phospholipids induced in intact cells by insulin-like growth factor I. *Biochem. Biophys. Res. Commun.* 159 (2), 720–725.
- D'Santos, C.S., Lewis, A.E., 2012. Functional proteomics: mapping lipid-protein interactomes. In: Leung, H.-C.E. (Ed.), *Integrative Proteomics*. InTech.
- Davis, W.J., Lehmann, P.Z., Li, W., 2015. Nuclear PI3K signaling in cell growth and tumorigenesis. *Front Cell Dev Biol* 3, 24.
- de Graaf, P., Klapisz, E.E., Schulz, T.K., Cremers, A.F., Verkleij, A.J., van Bergen en Henegouwen, P.M., 2002. Nuclear localization of phosphatidylinositol 4-kinase beta. *J. Cell Sci.* 115 (Pt 8), 1769–1775.
- Deleris, P., Bacqueville, D., Gayral, S., Carrez, L., Salles, J.P., Perret, B., Breton-Douillon, M., 2003. SHIP-2 and PTEN are expressed and active in vascular smooth muscle cell nuclei, but only SHIP-2 is associated with nuclear speckles. *J. Biol. Chem.* 278 (40), 38884–38891.
- Di Lello, P., Nguyen, B.D., Jones, T.N., Potempa, K., Kobor, M.S., Legault, P., Omichinski, J.G., 2005. NMR structure of the amino-terminal domain from the Tfb1 subunit of TFIIB and characterization of its phosphoinositide and VP16 binding sites. *Biochemistry* 44 (21), 7678–7686.
- Didichenko, S.A., Thelen, M., 2001. Phosphatidylinositol 3-kinase c2alpha contains a nuclear localization sequence and associates with nuclear speckles. *J. Biol. Chem.* 276 (51), 48135–48142.
- Divecha, N., Banfic, H., Irvine, R.F., 1991. The polyphosphoinositide cycle exists in the nuclei of Swiss 3T3 cells under the control of a receptor (for IGF-I) in the plasma membrane, and stimulation of the cycle increases nuclear diacylglycerol and apparently induces translocation of protein kinase C to the nucleus. *EMBO J.* 10 (11), 3207–3214.
- Dixon, M.J., Gray, A., Boisvert, F.M., Agacan, M., Morrice, N.A., Gourlay, R., Leslie, N.R., Downes, C.P., Batty, I.H., 2011. A screen for novel phosphoinositide 3-kinase effector proteins. *Mol. Cell. Proteomics* 10 (4) M110 003178.
- Drakas, R., Tu, X., Baserga, R., 2004. Control of cell size through phosphorylation of upstream binding factor 1 by nuclear phosphatidylinositol 3-kinase. *Proc. Natl.*

- Acad. Sci. U. S. A. 101 (25), 9272–9276.
- Ehm, P., Nalaskowski, M.M., Wundenberg, T., Jucker, M., 2015. The tumor suppressor SHIP1 colocalizes in nucleolar cavities with p53 and components of PML nuclear bodies. *Nucleus* 6 (2), 154–164.
- Elong Edimo, W., Derua, R., Janssens, V., Nakamura, T., Vanderwinden, J.M., Waelkens, E., Erneux, C., 2011. Evidence of SHIP2 Ser 132 phosphorylation, its nuclear localization and stability. *Biochem. J.* 439 (3), 391–401.
- Emmott, E., Hiscox, J.A., 2009. Nucleolar targeting: the hub of the matter. *EMBO Rep.* 10 (3), 231–238.
- Fahy, E., Subramaniam, S., Murphy, R.C., Nishijima, M., Raetz, C.R., Shimizu, T., Spener, F., van Meer, G., Wakelam, M.J., Dennis, E.A., 2009. Update of the LIPID MAPS comprehensive classification system for lipids. *J. Lipid Res.* 50 (Suppl. 1), S9–S14.
- Feng, L., Mejillano, M., Yin, H.L., Chen, J., Prestwich, G.D., 2001. Full-contact domain labeling: identification of a novel phosphoinositide binding site on gelsolin that requires the complete protein. *Biochemistry* 40 (4), 904–913.
- Fiume, R., Keune, W.J., Faenza, I., Bultsma, Y., Ramazzotti, G., Jones, D.R., Martelli, A.M., Somner, L., Follo, M.Y., Divecha, N., Cocco, L., 2012. Nuclear phosphoinositides: location, regulation and function. *Subcell. Biochem.* 59, 335–361.
- Gabev, E., Kasianowicz, J., Abbott, T., McLaughlin, S., 1989. Binding of neomycin to phosphatidylinositol 4,5-bisphosphate (PIP₂). *Biochim. Biophys. Acta* 979 (1), 105–112.
- Galganski, L., Urbanek, M.O., Krzyzosiak, W.J., 2017. Nuclear speckles: molecular organization, biological function and role in disease. *Nucleic Acids Res.* 45 (18), 10350–10368.
- Geeraerts, A., Hsiu-Fang, F., Zimmermann, P., Engelborghs, Y., 2013. The characterization of the nuclear dynamics of syntenin-2, a PIP₂ binding PDZ protein. *Cytometry* 83 (9), 866–875.
- Gelato, K.A., Tauber, M., Ong, M.S., Winter, S., Hiragami-Hamada, K., Sindlinger, J., Lemak, A., Bultsma, Y., Houliston, S., Schwarzer, D., Divecha, N., Arrowsmith, C.H., Fischle, W., 2014. Accessibility of different histone H3-binding domains of UHRF1 is allosterically regulated by phosphatidylinositol 5-phosphate. *Mol. Cell.* 54 (6), 905–919.
- Gillooly, D.J., Morrow, I.C., Lindsay, M., Gould, R., Bryant, N.J., Gaullier, J.M., Parton, R.G., Stenmark, H., 2000. Localization of phosphatidylinositol 3-phosphate in yeast and mammalian cells. *EMBO J.* 19 (17), 4577–4588.
- Gonzales, M.L., Mellman, D.L., Anderson, R.A., 2008. CK1 α is associated with and phosphorylates star-PAP and is also required for expression of select star-PAP target messenger RNAs. *J. Biol. Chem.* 283 (18), 12665–12673.
- Gorbatyuk, V.Y., Nosworthy, N.J., Robson, S.A., Bains, N.P., Maciejewski, M.W., Dos Remedios, C.G., King, G.F., 2006. Mapping the phosphoinositide-binding site on chick cofilin explains how PIP₂ regulates the cofilin-actin interaction. *Mol. Cell.* 24 (4), 511–522.
- Gozani, O., Field, S.J., Ferguson, C.G., Ewalt, M., Mahlke, C., Cantley, L.C., Prestwich, G.D., Yuan, J., 2005. Modification of protein sub-nuclear localization by synthetic phosphoinositides: evidence for nuclear phosphoinositide signaling mechanisms. *Adv. Enzym. Regul.* 45, 171–185.
- Gozani, O., Karuman, P., Jones, D.R., Ivanov, D., Cha, J., Lugovskoy, A.A., Baird, C.L., Zhu, H., Field, S.J., Lessnick, S.L., Villasenor, J., Mehrotra, B., Chen, J., Rao, V.R., Brugge, J.S., Ferguson, C.G., Payrastra, B., Myszkla, D.G., Cantley, L.C., Wagner, G., Divecha, N., Prestwich, G.D., Yuan, J.Y., 2003. The PHD finger of the chromatin-associated protein ING2 functions as a nuclear phosphoinositide receptor. *Cell* 114 (1), 99–111.
- Hamann, B.L., Blind, R.D., 2018. Nuclear phosphoinositide regulation of chromatin. *J. Cell. Physiol.* 233 (1), 107–123.
- Hammond, G.R., Balla, T., 2015. Polyphosphoinositide binding domains: key to inositol lipid biology. *Biochim. Biophys. Acta* 1851 (6), 746–758.
- Heo, W.D., Inoue, T., Park, W.S., Kim, M.L., Park, B.O., Wandless, T.J., Meyer, T., 2006. PI(3,4,5)P₃ and PI(4,5)P₂ lipids target proteins with polybasic clusters to the plasma membrane. *Science* 314 (5804), 1458–1461.
- Hu, Y., Liu, Z., Ye, K., 2005. Phosphoinositide lipids bind to phosphatidylinositol 3 (PI3)-kinase enhancer GTPase and mediate its stimulatory effect on PI3-kinase and Akt signalings. *Proc. Natl. Acad. Sci. U. S. A.* 102 (46), 16853–16858.
- Huang, W., Zhang, H., Davrazou, F., Kutateladze, T.G., Shi, X., Gozani, O., Prestwich, G.D., 2007. Stabilized phosphatidylinositol-5-phosphate analogues as ligands for the nuclear protein ING2: chemistry, biology, and molecular modeling. *J. Am. Chem. Soc.* 129 (20), 6498–6506.
- Hunt, A.N., Clark, G.T., Attard, G.S., Postle, A.D., 2001. Highly saturated endonuclear phosphatidylcholine is synthesized in situ and colocalized with CDP-choline pathway enzymes. *J. Biol. Chem.* 276 (11), 8492–8499.
- Irvine, R.F., 2006. Nuclear inositol signalling – expansion, structures and clarification. *Biochim. Biophys. Acta* 1761 (5–6), 505–508.
- Janmey, P.A., Bucki, R., Radhakrishnan, R., 2018. Regulation of actin assembly by PI(4,5)P₂ and other inositol phospholipids: an update on possible mechanisms. *Biochem. Biophys. Res. Commun.* 506 (2), 307–314.
- Janmey, P.A., Stossel, T.P., 1987. Modulation of gelsolin function by phosphatidylinositol 4,5-bisphosphate. *Nature* 325 (6102), 362–364.
- Jones, D.R., Bultsma, Y., Keune, W.J., Halstead, J.R., Elouarrat, D., Mohammed, S., Heck, A.J., D'Santos, C.S., Divecha, N., 2006. Nuclear PtdIns5P as a transducer of stress signaling: an in vivo role for PIP4K β . *Mol. Cell* 23 (5), 685–695.
- Jungmichel, S., Sylvestersen, K.B., Choudhary, C., Nguyen, S., Mann, M., Nielsen, M.L., 2014. Specificity and commonality of the phosphoinositide-binding proteome analyzed by quantitative mass spectrometry. *Cell Rep.* 6 (3), 578–591.
- Kaadige, M.R., Ayer, D.E., 2006. The polybasic region that follows the plant homeodomain zinc finger 1 of Pfl is necessary and sufficient for specific phosphoinositide binding. *J. Biol. Chem.* 281 (39), 28831–28836.
- Kakuk, A., Friedlander, E., Vereb Jr., G., Kasa, A., Balla, A., Balla, T., Heilmeyer Jr., L.M., Gergely, P., Vereb, G., 2006. Nucleolar localization of phosphatidylinositol 4-kinase PI4K230 in various mammalian cells. *Cytometry* 69 (12), 1174–1183.
- Kakuk, A., Friedlander, E., Vereb Jr., G., Lisboa, D., Bagossi, P., Toth, G., Gergely, P., Vereb, G., 2008. Nuclear and nucleolar localization signals and their targeting function in phosphatidylinositol 4-kinase PI4K230. *Exp. Cell Res.* 314 (13), 2376–2388.
- Kalassova, I., Faberova, V., Kalendova, A., Yildirim, S., Ulicna, L., Venit, T., Hozak, P., 2016. Tools for visualization of phosphoinositides in the cell nucleus. *Histochem. Cell Biol.* 145 (4), 485–496.
- Karlsson, T., Altankhyug, A., Dobrovol'ska, O., Turcu, D.C., Lewis, A.E., 2016. A polybasic motif in ErbB3-binding protein 1 (EBP1) has key functions in nucleolar localization and polyphosphoinositide interaction. *Biochem. J.* 473 (14), 2033–2047.
- Kostrhon, S., Kontaxis, G., Kaufmann, T., Schirghuber, E., Kubicek, S., Konrat, R., Slade, D., 2017. A histone-mimicking interdomain linker in a multidomain protein modulates multivalent histone binding. *J. Biol. Chem.* 292 (43), 17643–17657.
- Krugmann, S., Anderson, K.E., Ridley, S.H., Risso, N., McGregor, A., Coadwell, J., Davidson, K., Eguinoa, A., Ellson, C.D., Lipp, P., Manifava, M., Ktistakis, N., Painter, G., Thuring, J.W., Cooper, M.A., Lim, Z.Y., Holmes, A.B., Dove, S.K., Michell, R.H., Grewal, A., Nazarian, A., Erdjument-Bromage, H., Tempst, P., Stephens, L.R., Hawkins, P.T., 2002. Identification of ARAP3, a novel PI3K effector regulating both Arf and Rho GTPases, by selective capture on phosphoinositide affinity matrices. *Mol. Cell.* 9 (1), 95–108.
- Kumar, A., Fernandez-Capetillo, O., Carrera, A.C., 2010. Nuclear phosphoinositide 3-kinase beta controls double-strand break DNA repair. *Proc. Natl. Acad. Sci. U. S. A.* 107 (16), 7491–7496.
- Kumar, A., Redondo-Munoz, J., Perez-Garcia, V., Cortes, I., Chagoyen, M., Carrera, A.C., 2011. Nuclear but not cytosolic phosphoinositide 3-kinase beta has an essential function in cell survival. *Mol. Cell Biol.* 31 (10), 2122–2133.
- Kumar, N., Zhao, P., Tomar, A., Galea, C.A., Khurana, S., 2004. Association of villin with phosphatidylinositol 4,5-bisphosphate regulates the actin cytoskeleton. *J. Biol. Chem.* 279 (4), 3096–3110.
- Kunrath-Lima, M., de Miranda, M.C., Ferreira, A.D.F., Faraco, C.C.F., de Melo, M.I.A., Goes, A.M., Rodrigues, M.A., Faria, J., Gomes, D.A., 2018. Phospholipase C delta 4 (PLCdelta4) is a nuclear protein involved in cell proliferation and senescence in mesenchymal stromal stem cells. *Cell. Signal.* 49, 59–67.
- Kutateladze, T.G., 2010. Translation of the phosphoinositide code by PI effectors. *Nat. Chem. Biol.* 6 (7), 507–513.
- Kwon, I.S., Lee, K.H., Choi, J.W., Ahn, J.Y., 2010. PI(3,4,5)P₃ regulates the interaction between Akt and B23 in the nucleus. *BMB Rep* 43 (2), 127–132.
- Lambrechts, A., Jonckheere, V., Dewitte, D., Vandekerckhove, J., Ampe, C., 2002. Mutational analysis of human profilin I reveals a second PI(4,5)P₂ binding site neighbouring the poly(L-proline) binding site. *BMC Biochem.* 3, 12.
- Lanning, C.C., Ruiz-Velasco, R., Williams, C.L., 2003. Novel mechanism of the co-regulation of nuclear transport of SmgGDS and Rac1. *J. Biol. Chem.* 278 (14),

- 12495–12506.
- Lassing, I., Lindberg, U., 1985. Specific interaction between phosphatidylinositol 4,5-bisphosphate and profilactin. *Nature* 314 (6010), 472–474.
- Laux, T., Fukami, K., Thelen, M., Golub, T., Frey, D., Caroni, P., 2000. GAP43, MARCKS, and CAP23 modulate PI(4,5)P(2) at plasmalemmal rafts, and regulate cell cortex actin dynamics through a common mechanism. *J. Cell Biol.* 149 (7), 1455–1472.
- Layzerenza, J.P., Gonzalez, P., Garcia de Bravo, M.M., Polo, M.P., Sisti, M.S., Ves-Losada, A., 2013. Nuclear lipid droplets: a novel nuclear domain. *Biochim. Biophys. Acta* 1831 (2), 327–340.
- Lee, S.B., Xuan Nguyen, T.L., Choi, J.W., Lee, K.H., Cho, S.W., Liu, Z., Ye, K., Bae, S.S., Ahn, J.Y., 2008. Nuclear Akt interacts with B23/NPM and protects it from proteolytic cleavage, enhancing cell survival. *Proc. Natl. Acad. Sci. U. S. A.* 105 (43), 16584–16589.
- Lewis, A.E., Sommer, L., Arntzen, M.O., Strahm, Y., Morrice, N.A., Divecha, N., D'Santos, C.S., 2011. Identification of nuclear phosphatidylinositol 4,5-bisphosphate-interacting proteins by neomycin extraction. *Mol. Cell. Proteomics* 10 (2) M110 003376.
- Li, P., Wang, D., Li, H., Yu, Z., Chen, X., Fang, J., 2014. Identification of nucleolus-localized PTEN and its function in regulating ribosome biogenesis. *Mol. Biol. Rep.* 41 (10), 6383–6390.
- Liang, H., Chen, X., Yin, Q., Ruan, D., Zhao, X., Zhang, C., McNutt, M.A., Yin, Y., 2017. PTENbeta is an alternatively translated isoform of PTEN that regulates rDNA transcription. *Nat. Commun.* 8, 14771.
- Lindsay, Y., McCoull, D., Davidson, L., Leslie, N.R., Fairservice, A., Gray, A., Lucocq, J., Downes, C.P., 2006. Localization of agonist-sensitive PtdIns(3,4,5)P3 reveals a nuclear pool that is insensitive to PTEN expression. *J. Cell Sci.* 119 (Pt 24), 5160–5168.
- Liu, J.L., Sheng, X., Hortobagyi, Z.K., Mao, Z., Gallick, G.E., Yung, W.K., 2005. Nuclear PTEN-mediated growth suppression is independent of Akt down-regulation. *Mol. Cell Biol.* 25 (14), 6211–6224.
- Maag, D., Maxwell, M.J., Hardesty, D.A., Boucher, K.L., Choudhari, N., Hanno, A.G., Ma, J.F., Snowman, A.S., Pietropaoli, J.W., Xu, R., Storm, P.B., Saiardi, A., Snyder, S.H., Resnick, A.C., 2011. Inositol polyphosphate multikinase is a physiologic PI3-kinase that activates Akt/PKB. *Proc. Natl. Acad. Sci. U. S. A.* 108 (4), 1391–1396.
- Maffucci, T., Razzini, G., Ingrosso, A., Chen, H., Iacobelli, S., Sciacchitano, S., Quon, M.J., Falasca, M., 2003. Role of pleckstrin homology domain in regulating membrane targeting and metabolic function of insulin receptor substrate 3. *Mol. Endocrinol.* 17 (8), 1568–1579.
- Manzoli, F.A., Maraldi, N.M., Cocco, L., Capitani, S., Facchini, A., 1977. Chromatin phospholipids in normal and chronic lymphocytic leukemia lymphocytes. *Cancer Res.* 37 (3), 843–849.
- Manzoli, L., Billi, A.M., Gilmour, R.S., Martelli, A.M., Matteucci, A., Rubbini, S., Weber, G., Cocco, L., 1995. Phosphoinositide signaling in nuclei of Friend cells: tiiazofurin down-regulates phospholipase C beta 1. *Cancer Res.* 55 (14), 2978–2980.
- Marques, M., Kumar, A., Poveda, A.M., Zuluaga, S., Hernandez, C., Jackson, S., Pasero, P., Carrera, A.C., 2009. Specific function of phosphoinositide 3-kinase beta in the control of DNA replication. *Proc. Natl. Acad. Sci. U. S. A.* 106 (18), 7525–7530.
- Martelli, A.M., Gilmour, R.S., Bertagnolo, V., Neri, L.M., Manzoli, L., Cocco, L., 1992. Nuclear localization and signalling activity of phosphoinositidase C beta in Swiss 3T3 cells. *Nature* 358 (6383), 242–245.
- Martelli, A.M., Ognibene, A., Buontempo, F., Fini, M., Bressanin, D., Goto, K., McCubrey, J.A., Cocco, L., Evangelisti, C., 2011. Nuclear phosphoinositides and their roles in cell biology and disease. *Crit. Rev. Biochem. Mol. Biol.* 46 (5), 436–457.
- Martin, T.F., 1998. Phosphoinositide lipids as signaling molecules: common themes for signal transduction, cytoskeletal regulation, and membrane trafficking. *Annu. Rev. Cell Dev. Biol.* 14, 231–264.
- Matsuzaki, F., Matsumoto, S., Yahara, I., Yonezawa, N., Nishida, E., Sakai, H., 1988. Cloning and characterization of porcine brain cofilin cDNA. Cofilin contains the nuclear transport signal sequence. *J. Biol. Chem.* 263 (23), 11564–11568.
- McLane, L.M., Corbett, A.H., 2009. Nuclear localization signals and human disease. *IUBMB Life* 61 (7), 697–706.
- McLaughlin, S., Murray, D., 2005. Plasma membrane phosphoinositide organization by protein electrostatics. *Nature* 438 (7068), 605–611.
- Meerschaert, K., Tun, M.P., Remue, E., De Ganck, A., Boucherie, C., Vanloo, B., Degeest, G., Vandekerckhove, J., Zimmermann, P., Bhardwaj, N., Lu, H., Cho, W., Gettemans, J., 2009. The PDZ2 domain of zonula occludens-1 and -2 is a phosphoinositide binding domain. *Cell. Mol. Life Sci.* 66 (24), 3951–3966.
- Mellman, D.L., Gonzales, M.L., Song, C., Barlow, C.A., Wang, P., Kendzioriski, C., Anderson, R.A., 2008. A PtdIns4,5P2-regulated nuclear poly(A) polymerase controls expression of select mRNAs. *Nature* 451 (7181), 1013–1017.
- Michell, R.H., Heath, V.L., Lemmon, M.A., Dove, S.K., 2006. Phosphatidylinositol 3,5-bisphosphate: metabolism and cellular functions. *Trends Biochem. Sci.* 31 (1), 52–63.
- Mortier, E., Wuytens, G., Leenaerts, I., Hannes, F., Heung, M.Y., Degeest, G., David, G., Zimmermann, P., 2005. Nuclear speckles and nucleoli targeting by PIP2-PDZ domain interactions. *EMBO J.* 24 (14), 2556–2565.
- Munsie, L.N., Desmond, C.R., Truant, R., 2012. Cofilin nuclear-cytoplasmic shuttling affects cofilin-actin rod formation during stress. *J. Cell Sci.* 125 (Pt 17), 3977–3988.
- Ndamukong, I., Jones, D.R., Lapko, H., Divecha, N., Avramova, Z., 2010. Phosphatidylinositol 5-phosphate links dehydration stress to the activity of ARABIDOPSIS TRITHORAX-LIKE factor ATX1. *PLoS One* 5 (10), e13396.
- Ohmori, S., Sakai, N., Shirai, Y., Yamamoto, H., Miyamoto, E., Shimizu, N., Saito, N., 2000. Importance of protein kinase C targeting for the phosphorylation of its substrate, myristoylated alanine-rich C-kinase substrate. *J. Biol. Chem.* 275 (34), 26449–26457.
- Okada, M., Jang, S.W., Ye, K., 2008. Akt phosphorylation and nuclear phosphoinositide association mediate mRNA export and cell proliferation activities by ALY. *Proc. Natl. Acad. Sci. U. S. A.* 105 (25), 8649–8654.
- Okada, M., Taguchi, K., Maekawa, S., Fukami, K., Yagisawa, H., 2010. Calcium fluxes cause nuclear shrinkage and the translocation of phospholipase C-delta1 into the nucleus. *Neurosci. Lett.* 472 (3), 188–193.
- Onoda, K., Yin, H.L., 1993. gCap39 is phosphorylated. Stimulation by okadaic acid and preferential association with nuclei. *J. Biol. Chem.* 268 (6), 4106–4112.
- Onoda, K., Yu, F.X., Yin, H.L., 1993. gCap39 is a nuclear and cytoplasmic protein. *Cell Motil. Cytoskelet.* 26 (3), 227–238.
- Osborne, S.L., Thomas, C.L., Gschmeissner, S., Schiavo, G., 2001. Nuclear PtdIns(4,5)P2 assemblies in a mitotically regulated particle involved in pre-mRNA splicing. *J. Cell Sci.* 114 (Pt 13), 2501–2511.
- Osborne, S.L., Wallis, T.P., Jimenez, J.L., Gorman, J.J., Meunier, F.A., 2007. Identification of secretory granule phosphatidylinositol 4,5-bisphosphate-interacting proteins using an affinity pulldown strategy. *Mol. Cell. Proteomics* 6 (7), 1158–1169.
- Papayannopoulos, V., Co, C., Prehoda, K.E., Snapper, S., Taunton, J., Lim, W.A., 2005. A polybasic motif allows N-WASP to act as a sensor of PIP2 density. *Mol. Cell.* 17 (2), 181–191.
- Pasquali, C., Bertschy-Meier, D., Chabert, C., Curchod, M.L., Arod, C., Booth, R., Mechtler, K., Vilbois, F., Xenarios, I., Ferguson, C.G., Prestwich, G.D., Camps, M., Rommel, C., 2007. A chemical proteomics approach to phosphatidylinositol 3-kinase signaling in macrophages. *Mol. Cell. Proteomics* 6 (11), 1829–1841.
- Payrastre, B., Nievers, M., Boonstra, J., Breton, M., Verkleij, A.J., Van Bergen en Henegouwen, P.M., 1992. A differential location of phosphoinositide kinases, diacylglycerol kinase, and phospholipase C in the nuclear matrix. *J. Biol. Chem.* 267 (8), 5078–5084.
- Pederson, T., 2011. The nucleolus. *Cold Spring Harb Perspect Biol* 3 (3).
- Pemberton, J.G., Balla, T., 2018. Polyphosphoinositide-binding domains: Insights from peripheral membrane and lipid-transfer proteins. *Adv. Exp. Med. Biol.* https://doi.org/10.1007/5584_2018_288.
- Rando, O.J., Zhao, K., Janmey, P., Crabtree, G.R., 2002. Phosphatidylinositol-dependent actin filament binding by the SWI/SNF-like BAF chromatin remodeling complex. *Proc. Natl. Acad. Sci. U. S. A.* 99 (5), 2824–2829.
- Razzini, G., Ingrosso, A., Branaccio, A., Sciacchitano, S., Esposito, D.L., Falasca, M., 2000. Different subcellular localization and phosphoinositides binding of insulin receptor substrate protein pleckstrin homology domains. *Mol. Endocrinol.* 14 (6), 823–836.
- Resnick, A.C., Snowman, A.M., Kang, B.N., Hurt, K.J., Snyder, S.H., Saiardi, A., 2005. Inositol polyphosphate multikinase is a nuclear PI3-kinase with transcriptional regulatory activity. *Proc. Natl. Acad. Sci. U. S. A.* 102 (36), 12783–12788.
- Rohrbach, T.D., Shah, N., Jackson, W.P., Feeney, E.V., Scanlon, S., Gish, R., Khodadadi, R., Hyde, S.O., Hicks, P.H., Anderson, J.C., Jarboe, J.S., Willey, C.D., 2015. The effector domain of MARCKS is a nuclear localization signal that regulates cellular PIP2 levels and nuclear PIP2 localization. *PLoS One* 10 (10), e0140870.

- Romanouska, A., Kohler, A., 2018. The inner nuclear membrane is a metabolically active territory that generates nuclear lipid droplets. *Cell* 174 (3), 700–715 e718.
- Rose, H.G., Frenster, J.H., 1965. Composition and metabolism of lipids within repressed and active chromatin of interphase lymphocytes. *Biochim. Biophys. Acta Lipids Lipid Metab.* 106 (3), 577–591.
- Rowland, M.M., Bostic, H.E., Gong, D., Speers, A.E., Lucas, N., Cho, W., Cravatt, B.F., Best, M.D., 2011. Phosphatidylinositol 3,4,5-trisphosphate activity probes for the labeling and proteomic characterization of protein binding partners. *Biochemistry* 50 (51), 11143–11161.
- Sablin, E.P., Blind, R.D., Uthayaruban, R., Chiu, H.J., Deacon, A.M., Das, D., Ingraham, H.A., Fletterick, R.J., 2015. Structure of Liver Receptor Homolog-1 (NR5A2) with PIP3 hormone bound in the ligand binding pocket. *J. Struct. Biol.* 192 (3), 342–348.
- Sanchez, R., Zhou, M.M., 2011. The PHD finger: a versatile epigenome reader. *Trends Biochem. Sci.* 36 (7), 364–372.
- Schacht, J., 1976. Inhibition by neomycin of polyphosphoinositide turnover in subcellular fractions of Guinea-pig cerebral cortex in vitro. *J. Neurochem.* 27 (5), 1119–1124.
- Schacht, J., 1978. Purification of polyphosphoinositides by chromatography on immobilized neomycin. *J. Lipid Res.* 19 (8), 1063–1067.
- Schacht, J., 1979. Isolation of an aminoglycoside receptor from Guinea pig inner ear tissues and kidney. *Arch. Oto-Rhino-Laryngol.* 224 (1–2), 129–134.
- Schill, N.J., Anderson, R.A., 2009. Two novel phosphatidylinositol-4-phosphate 5-kinase type Igamma splice variants expressed in human cells display distinctive cellular targeting. *Biochem. J.* 422 (3), 473–482.
- Shah, Z.H., Jones, D.R., Sommer, L., Foulger, R., Bultsma, Y., D'Santos, C., Divecha, N., 2013. Nuclear phosphoinositides and their impact on nuclear functions. *FEBS J.* 280 (24), 6295–6310.
- Shen, W.H., Balajee, A.S., Wang, J., Wu, H., Eng, C., Pandolfi, P.P., Yin, Y., 2007. Essential role for nuclear PTEN in maintaining chromosomal integrity. *Cell* 128 (1), 157–170.
- Sindic, A., Aleksandrova, A., Fields, A.P., Volinia, S., Banfic, H., 2001. Presence and activation of nuclear phosphoinositide 3-kinase C2beta during compensatory liver growth. *J. Biol. Chem.* 276 (21), 17754–17761.
- Skare, P., Karlsson, R., 2002. Evidence for two interaction regions for phosphatidylinositol(4,5)-bisphosphate on mammalian profilin I. *FEBS Lett.* 522 (1–3), 119–124.
- Skare, P., Kreivi, J.P., Bergstrom, A., Karlsson, R., 2003. Profilin I colocalizes with speckles and Cajal bodies: a possible role in pre-mRNA splicing. *Exp. Cell Res.* 286 (1), 12–21.
- Sobol, M., Krausova, A., Yildirim, S., Kalasova, I., Faberova, V., Vrkoslav, V., Philimonenko, V., Marasek, P., Pastorek, L., Capek, M., Lubovska, Z., Ulicna, L., Tsuji, T., Lisa, M., Cvacka, J., Fujimoto, T., Hozak, P., 2018. Nuclear phosphatidylinositol 4,5-bisphosphate islets contribute to efficient RNA polymerase II-dependent transcription. *J. Cell Sci.* 131 (8).
- Sobol, M., Yildirim, S., Philimonenko, V.V., Marasek, P., Castano, E., Hozak, P., 2013. UBF complexes with phosphatidylinositol 4,5-bisphosphate in nucleolar organizer regions regardless of ongoing RNA polymerase I activity. *Nucleus* 4 (6), 478–486.
- Song, M.S., Carracedo, A., Salmena, L., Song, S.J., Egia, A., Malumbres, M., Pandolfi, P.P., 2011. Nuclear PTEN regulates the APC-CDH1 tumor-suppressive complex in a phosphatase-independent manner. *Cell* 144 (2), 187–199.
- Song, M.S., Salmena, L., Carracedo, A., Egia, A., Lo-Coco, F., Teruya-Feldstein, J., Pandolfi, P.P., 2008. The deubiquitinylation and localization of PTEN are regulated by a HAUSP-PML network. *Nature* 455 (7214), 813–817.
- Spector, D.L., Lamond, A.I., 2011. Nuclear speckles. *Cold Spring Harb Perspect Biol* 3 (2).
- Stijf-Bultsma, Y., Sommer, L., Tauber, M., Baalbaki, M., Giardoglou, P., Jones, D.R., Gelato, K.A., van Pelt, J., Shah, Z., Rahnamoun, H., Toma, C., Anderson, K.E., Hawkins, P., Lauberth, S.M., Haramis, A.P., Hart, D., Fischle, W., Divecha, N., 2015. The basal transcription complex component TAF3 transduces changes in nuclear phosphoinositides into transcriptional output. *Mol. Cell.* 58 (3), 453–467.
- Sud, M., Fahy, E., Cotter, D., Brown, A., Dennis, E.A., Glass, C.K., Merrill Jr., A.H., Murphy, R.C., Raetz, C.R., Russell, D.W., Subramaniam, S., 2007. LMSD: LIPID MAPS structure database. *Nucleic Acids Res.* 35, D527–D532 Database issue.
- Szivaik, I., Lamb, N., Heilmeyer, L.M., 2006. Subcellular localization and structural function of endogenous phosphorylated phosphatidylinositol 4-kinase (PI4K92). *J. Biol. Chem.* 281 (24), 16740–16749.
- Sztacho, M., Sobol, M., Balaban, C., Escudeiro Lopes, S.E., Hozak, P., 2019. Nuclear phosphoinositides and phase separation: important players in nuclear compartmentalization. *Adv Biol Regul* 71, 111–117.
- Tabellini, G., Bortul, R., Santi, S., Riccio, M., Baldini, G., Cappellini, A., Billi, A.M., Berezney, R., Ruggeri, A., Cocco, L., Martelli, A.M., 2003. Diacylglycerol kinase-theta is localized in the speckle domains of the nucleus. *Exp. Cell Res.* 287 (1), 143–154.
- Tanaka, K., Horiguchi, K., Yoshida, T., Takeda, M., Fujisawa, H., Takeuchi, K., Umeda, M., Kato, S., Ihara, S., Nagata, S., Fukui, Y., 1999. Evidence that a phosphatidylinositol 3,4,5-trisphosphate-binding protein can function in nucleus. *J. Biol. Chem.* 274 (7), 3919–3922.
- Tanaka, K., Imajoh-Ohmi, S., Sawada, T., Shirai, R., Hashimoto, Y., Iwasaki, S., Kaibuchi, K., Kanaho, Y., Shirai, T., Terada, Y., Kimura, K., Nagata, S., Fukui, Y., 1997. A target of phosphatidylinositol 3,4,5-trisphosphate with a zinc finger motif similar to that of the ADP-ribosylation-factor GTPase-activating protein and two pleckstrin homology domains. *Eur. J. Biochem.* 245 (2), 512–519.
- Terashita, A., Funatsu, N., Umeda, M., Shimada, Y., Ohno-Iwashita, Y., Epan, R.M., Maekawa, S., 2002. Lipid binding activity of a neuron-specific protein NAP-22 studied in vivo and in vitro. *J. Neurosci. Res.* 70 (2), 172–179.
- Toska, E., Campbell, H.A., Shandilya, J., Goodfellow, S.J., Shore, P., Medler, K.F., Roberts, S.G., 2012. Repression of transcription by WT1-BASP1 requires the myristoylation of BASP1 and the PIP2-dependent recruitment of histone deacetylase. *Cell Rep.* 2 (3), 462–469.
- Ulicna, L., Kalendova, A., Kalasova, I., Vacik, T., Hozak, P., 2018. PIP2 epigenetically represses rRNA genes transcription interacting with PHF8. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1863 (3), 266–275.
- Vann, L.R., Wooding, F.B., Irvine, R.F., Divecha, N., 1997. Metabolism and possible compartmentalization of inositol lipids in isolated rat-liver nuclei. *Biochem. J.* 327 (Pt 2), 569–576.
- Viiiri, K., Maki, M., Lohi, O., 2012. Phosphoinositides as regulators of protein-chromatin interactions. *Sci. Signal.* 5 (222), pe19.
- Viiiri, K.M., Janis, J., Siggers, T., Heinonen, T.Y., Valjakka, J., Bulyk, M.L., Maki, M., Lohi, O., 2009. DNA-binding and -bending activities of SAP30L and SAP30 are mediated by a zinc-dependent module and monophosphoinositides. *Mol. Cell Biol.* 29 (2), 342–356.
- Visa, N., Percipalle, P., 2010. Nuclear functions of actin. *Cold Spring Harb Perspect Biol* 2 (4), a000620.
- Wang, J., Arbuзова, A., Hangyas-Mihalayne, G., McLaughlin, S., 2001. The effector domain of myristoylated alanine-rich C kinase substrate binds strongly to phosphatidylinositol 4,5-bisphosphate. *J. Biol. Chem.* 276 (7), 5012–5019.
- Wang, J., Gambhir, A., Hangyas-Mihalayne, G., Murray, D., Golebiewska, U., McLaughlin, S., 2002. Lateral sequestration of phosphatidylinositol 4,5-bisphosphate by the basic effector domain of myristoylated alanine-rich C kinase substrate is due to nonspecific electrostatic interactions. *J. Biol. Chem.* 277 (37), 34401–34412.
- Watt, S.A., Kimber, W.A., Fleming, I.N., Leslie, N.R., Downes, C.P., Lucocq, J.M., 2004. Detection of novel intracellular agonist responsive pools of phosphatidylinositol 3,4-bisphosphate using the TAPP1 pleckstrin homology domain in immunoelectron microscopy. *Biochem. J.* 377 (Pt 3), 653–663.
- Watt, S.A., Kular, G., Fleming, I.N., Downes, C.P., Lucocq, J.M., 2002. Subcellular localization of phosphatidylinositol 4,5-bisphosphate using the pleckstrin homology domain of phospholipase C delta1. *Biochem. J.* 363 (Pt 3), 657–666.
- Wickramasinghe, V.O., Savill, J.M., Chavali, S., Jonsdottir, A.B., Rajendra, E., Gruner, T., Laskey, R.A., Babu, M.M., Venkitaraman, A.R., 2013. Human inositol polyphosphate multikinase regulates transcript-selective nuclear mRNA export to preserve genome integrity. *Mol. Cell.* 51 (6), 737–750.
- Williams, C.L., 2003. The polybasic region of Ras and Rho family small GTPases: a regulator of protein interactions and membrane association and a site of nuclear localization signal sequences. *Cell. Signal.* 15 (12), 1071–1080.
- Wu, X., Yoo, Y., Okuhama, N.N., Tucker, P.W., Liu, G., Guan, J.L., 2006. Regulation of RNA-polymerase-II-dependent transcription by N-WASP and its nuclear-binding partners. *Nat. Cell Biol.* 8 (7), 756–763.
- Yang, X., Ongusaha, P.P., Miles, P.D., Havstad, J.C., Zhang, F., So, W.V., Kudlow, J.E., Michell, R.H., Olefsky, J.M., Field, S.J., Evans, R.M., 2008. Phosphoinositide signalling links O-GlcNAc transferase to insulin resistance. *Nature* 451 (7181), 964–969.
- Yildirim, S., Castano, E., Sobol, M., Philimonenko, V.V., Dzijaik, R., Venit, T., Hozak, P., 2013. Involvement of phosphatidylinositol 4,5-bisphosphate in RNA polymerase I transcription. *J. Cell Sci.* 126 (Pt 12), 2730–2739.

- Yin, H.L., Janmey, P.A., 2003. Phosphoinositide regulation of the actin cytoskeleton. *Annu. Rev. Physiol.* 65, 761–789.
- Yokogawa, T., Nagata, S., Nishio, Y., Tsutsumi, T., Ihara, S., Shirai, R., Morita, K., Umeda, M., Shirai, Y., Saitoh, N., Fukui, Y., 2000. Evidence that 3'-phosphorylated polyphosphoinositides are generated at the nuclear surface: use of immunostaining technique with monoclonal antibodies specific for PI 3,4-P(2). *FEBS Lett.* 473 (2), 222–226.
- Yu, F.X., Johnston, P.A., Sudhof, T.C., Yin, H.L., 1990. gCap39, a calcium ion- and polyphosphoinositide-regulated actin capping protein. *Science* 250 (4986), 1413–1415.
- Yu, F.X., Sun, H.Q., Janmey, P.A., Yin, H.L., 1992. Identification of a polyphosphoinositide-binding sequence in an actin monomer-binding domain of gelsolin. *J. Biol. Chem.* 267 (21), 14616–14621.
- Yu, H., Fukami, K., Watanabe, Y., Ozaki, C., Takenawa, T., 1998. Phosphatidylinositol 4,5-bisphosphate reverses the inhibition of RNA transcription caused by histone H1. *Eur. J. Biochem.* 251 (1–2), 281–287.
- Zhao, K., Wang, W., Rando, O.J., Xue, Y., Swiderek, K., Kuo, A., Crabtree, G.R., 1998. Rapid and phosphoinositol-dependent binding of the SWI/SNF-like BAF complex to chromatin after T lymphocyte receptor signaling. *Cell* 95 (5), 625–636.
- Zou, J., Marjanovic, J., Kisseleva, M.V., Wilson, M., Majerus, P.W., 2007. Type I phosphatidylinositol-4,5-bisphosphate 4-phosphatase regulates stress-induced apoptosis. *Proc. Natl. Acad. Sci. U. S. A* 104 (43), 16834–16839.