

Article

Diversity and Distribution of Mites (Acari: Ixodida, Mesostigmata, Trombidiformes, Sarcoptiformes) in the Svalbard Archipelago

Anna Seniczak ^{1,*}, Stanisław Seniczak ², Marla D. Schwarzfeld ³ and Stephen J. Coulson ^{4,5} and Dariusz J. Gwiazdowicz ⁶

¹ Department of Natural History, University Museum of Bergen, University of Bergen, Postboks 7800, 5020 Bergen, Norway

² Department Evolutionary Biology, Faculty of Biological Sciences, Kazimierz Wielki University, J.K. Chodkiewicza 30, 85-064 Bydgoszcz, Poland; stseni@ukw.edu.pl

³ Canadian National Collection of Insects, Arachnids and Nematodes, Agriculture and Agri-food Canada, 960 Carling Avenue, Ottawa, ON K1A 0C6, Canada; marla.schwarzfeld@canada.ca

⁴ Swedish Species Information Centre, Swedish University of Agricultural Sciences, SLU Artdatabanken, Box 7007, 75007 Uppsala, Sweden; stephen.coulson@slu.se

⁵ Department of Arctic Biology, University Centre in Svalbard, P.O. Box 156, 9171 Longyearbyen, Svalbard, Norway

⁶ Faculty of Forestry, Poznań University of Life Sciences, Wojska Polskiego 71c, 60-625 Poznań, Poland; dariusz.gwiazdowicz@up.poznan.pl

* Correspondence: Anna.Seniczak@uib.no

Received: 21 July 2020; Accepted: 19 August 2020; Published: 25 August 2020

Abstract: Svalbard is a singular region to study biodiversity. Located at a high latitude and geographically isolated, the archipelago possesses widely varying environmental conditions and unique flora and fauna communities. It is also here where particularly rapid environmental changes are occurring, having amongst the fastest increases in mean air temperature in the Arctic. One of the most common and species-rich invertebrate groups in Svalbard is the mites (Acari). We here describe the characteristics of the Svalbard acarofauna, and, as a baseline, an updated inventory of 178 species (one Ixodida, 36 Mesostigmata, 43 Trombidiformes, and 98 Sarcoptiformes) along with their occurrences. In contrast to the Trombidiformes and Sarcoptiformes, which are dominated in Svalbard by species with wide geographical distributions, the Mesostigmata include many Arctic species (39%); it would thus be an interesting future study to determine if mesostigmatid communities are more affected by global warming than other mite groups. A large number of new species (42 spp.) have been described from Svalbard, including 15 that have so far been found exclusively there. It is yet uncertain if any of these latter species are endemic: six are recent findings, the others are old records and, in most cases, impossible to verify. That the Arctic is still insufficiently sampled also limits conclusions concerning endemism.

Keywords: checklist; Astigmata; Endeostigmata; Oribatida; Prostigmata; climate change; Arctic; Svalbard

1. Introduction

The invertebrate fauna of the Svalbard archipelago is amongst the best known for any region in the Arctic [1]. However, the inventory has been collated from studies over a period of 150 years and has never been subjected to a critical review and taxonomic revision. Without an improved

understanding of the current fauna it is not possible to understand the current terrestrial ecosystem or determine its response to on-going environmental change.

The dramatic environmental changes that we see today may be most discernable in polar regions, which possess both a unique flora and fauna and are undergoing the greatest rates of climate change. A particularly interesting area to study this is the Svalbard archipelago that is located at 78° N, isolated, and with a distinctive and diverse flora and fauna that are poorly studied [2]. Svalbard is currently experiencing one of the fastest temperature increases and one of the highest rates of sea ice loss in the Arctic [3]. During the past 50 years, the mean air temperature in Svalbard has increased by 4 °C, the winter mean air temperature has increased by 7 °C, the snow season duration has decreased by 20 days, and precipitation has increased by 65% [4]. For example, at Svalbard airport in 2018, the mean annual temperature was -1.8 °C, winter temperature was -5.3 °C, and annual precipitation was 252.5 mm [5]. These factors, as well as others that are tightly connected to the climate (e.g., faster melting of glaciers, changes in permafrost, landslides, avalanches, and flooding), affect the ecosystems in different manners. Even though a few species benefit from a warming climate, for example vertebrates, most Arctic species in Svalbard are experiencing negative effects in the warming environment [6].

Just south of Svalbard, two sea currents meet, a northern branch of the warm North Atlantic Drift and the southerly flowing cold Arctic Current, with consequences for the local climates on the east and west coasts of the archipelago [7] and potentially for species immigration histories, thus affecting species communities. A good example is the distinctive acarofauna of Edgeøya in the east of the archipelago compared to that observed on the west coast [8] and which includes an oribatid species new to science and not so far recorded elsewhere [9].

As a consequence of the increasing temperatures and changes in precipitation patterns, many glaciers are receding and revealing new land surfaces for primary succession. Therefore, regions such as Svalbard offer a unique opportunity to identify the pioneer species that first colonize barren post-glacial habitats and to analyze long-term processes of change, e.g., in invertebrate community assembly [10].

The diversity and richness of the environment is much more than simply lists of species occurring in a given area; it is also the regional specificity of these species, for example, those species that are considered to be typically High Arctic. Changing climatic conditions that determine and stimulate changes in habitats also affect the ranges of some mite species. Several studies have considered changes in mite communities in Svalbard in space and time; their dispersal abilities [11–17], as well as their emergence in new areas or retreat from previously occupied locations [10,18–20]. The uniqueness of polar areas, with their characteristic climates and isolated island distribution that often restricts dispersal processes, limits the range of some species. In this context, knowledge of zoogeography (ranges of occurrence of species) is extremely valuable because this suggests not only the migration routes, but also the ability of selected species to colonize new areas. Nonetheless, the climate of Svalbard is changing fast. These changes will undoubtedly provide opportunities for new species to colonise the region with difficult to foresee outcomes. The consequences of human introductions of invasive and alien species (IAS) have been described in the Antarctic [21] and there are examples of similar introductions in Svalbard [18].

Understanding mite ecology is also important since mites are excellent bioindicators of environmental changes and their presence, or changes in their communities, can be used in biological monitoring of naturally occurring processes as well as the consequences of human impacts, for example, those related to industrial activities in Svalbard's natural environment [20,22].

We here present a review of the mite fauna of Svalbard. These results can be the basis for further analyzes of the acarofauna (e.g., zoogeographic, ecological, taxonomic, and parasitological).

2. Material and Methods

The data were extracted from 104 papers published between 1871 and 2020. The localities presented in Figure 1 are based on Table 1. Distribution of *Ixodes uriae* White, 1852, the only representative of the order Ixodida in Svalbard, follows [23]. The nomenclature and arrangement of

the Mesostigmata families follows [24,25], while the species nomenclature follows [26] with a few modifications [27,28]. Distribution of the families follows [27–30].

The arrangement of the Trombidiformes families follow [31] with the exception of the Eriophyidae that is included in the Endeostigmata, as suggested by [32,33]. Names and distributions of the Bdellidae follow [34]; Cunaxidae—[35]; Cocceupodidae and Eupodidae—[36,37] for nomenclature, and [38,39] for distributions; Penthaleidae—[40]; Penthalodidae—[41]; Rhagidiidae—[42–44]; Halacaridae—[45]; Triophtydeiidae—[46] for taxonomy, and [47,48] for distribution; Tydeidae—[49,50]; Iolinidae—[51,52] for taxonomy, and [47,53] for distributions; Sperchontidae—[48,54,55]; Trombidiidae—[56]; Syringophilidae—[57]; Stigmeidae—[58]; Tetranychidae—[59,60]; and Pygmephoridae—[61].

The nomenclature and arrangement of the Sarcoptiformes families follow [62,63]. For the Endeostigmata, the nomenclature and arrangement of families follow [63,64]. The distribution of Nanorchestidae follow [65]; Alicorhagiidae—[66]; and Eriophyidae—[67]. The species names of the Oribatida (without Astigmata) follow [68] with a few exceptions [69,70] and their distribution is given after [9,68,71,72]. The Astigmata species (families Acaridae, Alloptidae, and Avenzoariidae) follow [73,74] and their distribution is according to [75].

Full names of species are provided in Table 1, while in other tables and figures abbreviations are used. The species within families are organized alphabetically. The names of localities follow in alphabetical order (Table 1).

3. Results

3.1. Ixodida


Only one species, the seabird tick (*Ixodes uriae*), is known from Svalbard. This species has a wide geographical distribution (Table 1) but in Svalbard has been found only recently on Bjørnøya and Spitsbergen [76–78].

3.2. Mesostigmata

Thus far, 36 species of Mesostigmata from 13 families have been found in Svalbard (Table 1). Amongst these families, the richest in species is the Ascidae (12 spp.), followed by the Zerconidae (6 spp.); these two families contribute almost 50% of mesostigmatic alpha diversity in Svalbard (Figure 2).

The peculiarity of the mesostigmatic mite communities of Svalbard compared to other mite groups is manifested by the description of one new genus (*Arctoseius*) and seven species [*Halolaelaps coulsoni* Gwiazdowicz & Teodorowicz, 2017; *H. gerlachi* Hirschmann, 1966; *Antennoseius (Vitzthumia) oudemansi* (Thor, 1930); *Arctoseius laterincisus* Thor, 1930; *Proctolaelaps parvanalis* (Thor, 1930); *Neoseiulus grumantensis* Kolodochka & Gwiazdowicz, 2014; and *N. magnanalis* (Thor, 1930)], all first described from the largest island of the archipelago—Spitsbergen. Moreover, based on recently collected samples from this island, redescriptions of several species have been published: *Zercon solenites* Haarløv, 1942; *Antennoseius (Vitzthumia) oudemansi*; *Arctoseius haarlovi* Lindquist, 1963; *Proctolaelaps parvanalis*; *Neoseiulus ellesmerei* (Chant & Hansell, 1971); and *N. magnanalis* [27–29,79,80].

Svalbard



Map Key

(54) Advent City	(74) Carmichaelpynten	(88) Großer Trichter	(108) Kinnvika
(51) Adventdalen	(38) Colesbukta	(49) Grumantbyen	(87) Knudsenheia
(41) Adventfjorden	(114) Dirksbukta	(111) Gyldénayane	(12) Koefoedodden
(36) Aldegondabreen	(18) Diskobukta	(58) Hanaskogdalen	(81) Kolhaugen
(7) Alfredjellet	(8) Ellasjøen	(6) Hellevatnet	(23) Kong Karls Land
(47) Arctowskifjellet	(39) Endalen	(52) Helvetiadalen	(82) Kongsfjorden
(5) Austervåg	(1) Engelskelva	(105) Hinlopenstretet	(65) Krykkjefjellet
(34) Barentsburg	(31) Engelskhytta	(59) Hiorthfjellet	(115) Lågøya
(22) Barentsøya	(102) Eremitten	(48) Hiorthhamn	(107) Lakssjøen
(27) Bellsund	(56) Fivelflyene	(14) Hopen	(95) Liefdefjorden
(62) Billefjorden	(72) Fjortende Julibukta	(13) Hopen radio	(104) Lomfjorden
(11) Birds colony	(94) Flathukken	(25) Hornsund	(55) Longyearbyen
(98) Biskayerhuken	(109) Florabukta	(53) Isdammen	(67) MacKenziedale
(40) Bjørndalen	(110) Fosterøyane	(44) Isfjord radio	(77) Magdalenefjorden
(20) Blåbukta	(68) Fuglehukken fyr	(42) Isfjorden	(33) Mälardalen
(71) Blomstrandhalvøya	(2) Fugleodden	(21) Kapp Heuglin	(69) Midtre Lovénbreen
(99) Bockfjorden	(61) Gipsdalen	(30) Kapp Laila	(4) Mosevatnet
(43) Bohemanneset	(35) Grønfjordbreen	(19) Kapp Lee	(90) Mosselbukta
(63) Brucebyen	(37) Grønfjorden	(70) Kiærstranda	(89) Mushamna
(15) Negerdalen	(106) Sofiaøya		
(76) Nissenfjella	(78) Sørgattet		
(29) Nordenskiöld Land	(112) Sørfjorden		
(101) Nordenskiöldøya	(24) Sørkapp		
(80) Ny-Ålesund	(91) Station Islands at Liefdefjorden		
(100) Ny-Friesland	(9) Stevatnet		
(84) Observasjonsholmen	(86) Storholmen		
(83) Ossian Sarsfjellet	(85) Stuphallet		
(64) Petuniabukta	(92) Sven Olssonodden		
(46) Pyramiden	(10) Teltvika		
(26) Recherchefjorden	(60) Tempelfjorden		
(96) Reinsdyrflya	(16) Tjuvfjorden		
(73) Richardlaguna	(45) Trygghamna		
(113) Ringhorndalen	(3) Tunheim		
(17) Russebukta	(28) Vårsolbukta		
(57) Sassendalen	(50) Vestpynten		
(32) Sassenfjorden	(79) Virgohamna		
(75) Signehamna	(103) Von Otterøya		
(66) Silene Hill	(93) Worsleyneset		



Figure 1. Localities in Svalbard where mites were found.

Table 1. List of mites of the Svalbard archipelago.

No.	TAXON	DISTRIBUTION	LOCALITIES
IXODIDA			
Ixodidae			
1.	<i>Ixodes uriae</i> White, 1852	Afrotropical, Australasian, Neotropical, Holarctic	Bjørnøya [77]; Spitsbergen: Ossian Sarsfjellet [76–78]
MESOSTIGMATA			
Dinychidae			
2	<i>Uroseiulus acuminatus</i> (C.L. Koch, 1847)	Palearctic	Spitsbergen [81,82]
Zerconidae			
3.	<i>Zercon andrei</i> Sellnick, 1958	Palearctic	Bjørnøya: Alfredfjellet, “Birds colony”, Teltvika [29,30,83]; Spitsbergen: Hornsund [84]
4.	<i>Z. curiosus</i> Trägårdh, 1910	Palearctic	Spitsbergen: Longyearbyen [85]
5.	<i>Z. forsslundi</i> Sellnick, 1958	Palearctic	Edgeøya: Diskobukta, Kapp Heuglin, Kapp Lee, Negerdalen, Russebukta [8]; Hopen: Koefoedodden, Norwegian Meteorological Institute Station [86]; Nordaustlandet: Kinnvika [30,87]; Spitsbergen: Adventdalen, Billefjord, Bjørndalen, Endalen, Hornsund, Ny-Ålesund, Ossian Sarsfjellet, Petuniabukta, Storholmen, Trygghamna [10,20,30,85,88–90] Edgeøya: Diskobukta, Kapp Lee [8]; Hopen: Koefoedodden, Norwegian Meteorological Institute Station [86]; Nordaustlandet: Kinnvika [29,87]; Spitsbergen: Adventdalen, Bjørndalen, Blomstrandhalvøya, Endalen, Fjortende Julibukta, Grønfjorden, Magdalenefjorden, Petuniabukta, Storholmen, Vårsolbukta [20,27,30,89,90]
6.	<i>Z. solenites</i> Haarløw, 1942 **	Arctic	Spitsbergen: Adventdalen, Barentsburg, Hanaskogdalen, Hjorthammmen [91]
7.	<i>Z. triangularis</i> C.L. Koch, 1836	Palearctic	
8.	<i>Z. zelawaiensis</i> Sellnick, 1944	Palearctic	Bjørnøya: Alfredfjellet [88]; Spitsbergen: Hornsund [83]
Parasitidae			
<i>Paragamasus</i>			
9.	<i>(Aclerogamasus) insertus</i> (Micherdzinski, 1969)	Palearctic	Spitsbergen: Barentsburg [18,19]
10.	<i>Vulgarogamasus</i> <i>immanis</i> (Berlese, 1904)	Palearctic	Spitsbergen: Barentsburg [92]
11.	<i>V. remberti</i> (Oudemans 1912)	Palearctic	Spitsbergen: Barentsburg [18,19]
Digamasellidae			
12.	<i>Dendrolaelaps foveolatus</i> (Leitner, 1949)	Palearctic	Spitsbergen: Pyramiden [20]
Halolaelapidae			
<i>Halolaelaps coulsoni</i>			
13.	Gwiazdowicz & Teodorowicz, 2017 *†	Arctic	Spitsbergen: Pyramiden [93]
14.	<i>H. gerlachi</i> Hirschmann, 1966 *†	Arctic	Spitsbergen: Ny-Ålesund [94]
15.	<i>Saproscenans baloghi</i> Karg, 1964	Palearctic	Edgeøya: Diskobukta [8]
Ologamasidae			
16.	<i>Gamasellus borealis</i> (C.L. Koch, 1879)	Palearctic	Spitsbergen: Dirksbukta [7]
Eviphidiidae			
17.	<i>Thinoseius spinosus</i> (Willmann, 1939)	Palearctic	Spitsbergen: Longyearbyen, Storholmen [90,95]
Macrochelidae			
<i>Macrocheles</i>			
18.	<i>muscaedomesticae</i> (Scopoli, 1772)	Palearctic	Spitsbergen: Barentsburg [84]
Ascidae			
19.	<i>Antennoseius</i>	Arctic	Bjørnøya: “Birds colony” [83]; Edgeøya: Diskobukta, Kapp Lee,

	<i>(Vitzthumia) oudemansi</i> (Thor, 1930) **		Russebukta [8]; Hopen: Koefoedodden, Norwegian Meteorological Institute Station [86]; Nordaustlandet: Kinnvika [30,79,87]; Spitsbergen: Adventdalen, Barentsburg, Bjørndalen, Endalen, Florabukta, Hiorthamn, Magdalenefjorden, Petuniabukta, Pyramiden, Storholmen, Vestpynten [19,20,22,30,79,85,88–91]
20.	<i>Arctoseius babenkoi</i> Makarova, 1999	Palearctic	Spitsbergen: Hornsund, Pyramiden [20,84,96]
21.	<i>A. cetratus</i> (Sellnick, 1940)	Holarctic	Spitsbergen [96]
22.	<i>A. haarlovi</i> Lindquist, 1963 **	Arctic	Bjørnøya: “Birds colony” [83]; Edgeøya: Diskobukta, Kapp Lee, Negerdalen, Russebukta [8]; Hopen: Koefoedodden, Norwegian Meteorological Institute Station [86]; Spitsbergen: Adventdalen, Barentsburg, Bjørndalen, Endalen, Hornsund, Longyearbyen, Magdalenefjorden, Mushamna, Ny-Ålesund, Petuniabukta, Pyramiden, Storholmen, Vestpynten [10,19,20,22,30,80,89,90,97]
23.	<i>A. laterincisus</i> Thor, 1930 *†	Arctic	Spitsbergen: Adventdalen, Barentsburg, Hanaskogdalen, Hjorthamn [91] Edgeøya: Blåbukta, Diskobukta, Kapp Heuglin, Kapp Lee, Negerdalen, Russebukta [8]; Hopen: Koefoedodden, Norwegian Meteorological Institute Station [86]; Nordaustlandet: Florabukta, Kinnvika [30,87]; Spitsbergen: Adventdalen, Barentsburg, Billefjorden, Endalen, Grønfjorden, Hornsund, Magdalenefjorden, Mushamna, Ny-Ålesund, Petuniabukta, Pyramiden, Storholmen, Sørkapp, Vårsolbukta [10,19,20,30,88–90,96]
24.	<i>A. multidentatus</i> Evans, 1955	Arctic	Spitsbergen [96]
25.	<i>A. ornatus</i> Lindquist, 1961	Holarctic	Bjørnøya: Alfredfjellet [83]; Spitsbergen: Hornsund [88,96]
26.	<i>A. tajmyricus</i> Petrova & Makarova, 1991	Palearctic	Edgeøya: Blåbukta, Kapp Heuglin, Negerdalen [8]; Nordaustlandet: Kinnvika [30,87]; Spitsbergen: Adventdalen, Endalen, Hornsund [30,89]
27.	<i>A. tschernovi</i> Makarova, 2000	Arctic	Bjørnøya: Alfredfjellet, „Birds colony” [83]; Edgeøya: Kapp Heuglin, Kapp Lee, Negerdalen, Russebukta [8]; Spitsbergen: Adventdalen, Bjørndalen, Endalen, Ossian Sarsfjellet, Petuniabukta, Vestpynten, Vårsolbukta [20,22,30,88,89,96]
28.	<i>A. weberi</i> Evans, 1955	Arctic	Bjørnøya: Alfredfjellet, “Birds colony” [83]; Spitsbergen [84]
29.	<i>Zerconopsis labradorensis</i> Evans & Till, 1960	Arctic	Bjørnøya: Teltvika [29,30]
30.	<i>Z. muestairi</i> (Schweizer, 1949)	Palearctic	Melicharidae
31.	<i>Proctolaelaps parvanalis</i> (Thor, 1930) ***	Arctic	Bjørnøya: Alfredfjellet, “Birds colony” [83]; Edgeøya: Blåbukta, Diskobukta, Kapp Lee, Russebukta [8]; Spitsbergen: Adventdalen, Bjørndalen, Endalen, Hiorthamn, Longyearbyen, Mälardalen, Ny-Ålesund, Petuniabukta [10,20,22,29,30,85,89,91]
32.	Phytoseiidae <i>Neoseiulus ellesmerei</i> (Chant & Hansell, 1971) **	Arctic	Edgeøya: Russebukta [8,28]; Spitsbergen: Adventdalen, Endalen, Longyearbyen [28]
33.	<i>N. grumantensis</i> Kolodochka & Gwiazdowicz, 2014 *†	Arctic	Spitsbergen: Longyearbyen, Petuniabukta [28]
34.	<i>N. magnanalans</i> (Thor, 1930) *†	Arctic	Spitsbergen: Adventdalen, Hiorthamn, Mälardalen, Ny-Ålesund, Petuniabukta [10,20,22,28,30,85,89,91]
35.	Dermyssidae <i>Dermyssus</i> <i>hirundinis</i> (Hermann, 1804)	Holarctic	Edgeøya: Blåbukta [8]; Spitsbergen: Longyearbyen, Ny-Ålesund [90,98]
36.	Laelapidae <i>Haemogamasus ambulans</i> (Thorel, 1872)	Holarctic	Bjørnøya [99]; Spitsbergen: Adventdalen, Endalen, Grumantbyen, Fjortende Julibukta, Ossian Sarsfjellet, Storholmen [90,91,100]
37.	<i>Laelaps hilaris</i> C.L. Koch, 1836	Holarctic	Spitsbergen: Grumantbyen [100]
	TROMBIDIFORMES		
	Prostigmata		
	Bdellidae		

38.	<i>Bdella longicornis</i> (Linnaeus, 1758)	Holarctic, Neotropic, Indomalayan	Bjørnøya: south [99,101]; Prins Karls Forland: MacKenziedale, Richardlaguna almost to Carmichaelpynten, Silene Hill [99,101]; Spitsbergen: Adventfjorden, Bellsund, Billefjorden, Bohemanneset, Brucebyen, Dirksbukta, Eremitten, Gipsdal, Hiorthhamn, Sorgfjorden, Station Islands at Liefdefjorden, Salmon Lake at Dirksbukta [7,91,99,101,102]
39.	<i>B. semiscutata</i> Thor, 1930 *	Palearctic	Spitsbergen: Engelskhytta, Hiorthfjellet, Longyearbyen [91,103]
40.	<i>Odontoscirrus lapidaria</i> (Kramer, 1881)	Cosmopolitan	Bjørnøya [104]; Spitsbergen: Barentsburg [91]
41.	<i>Cyta latirostris</i> (Hermann, 1804)	Cosmopolitan	Bjørnøya: Ellasjøen, south [99,101,105]; Prins Karls Forland: MacKenziedale, Silene Hill [99,101]; Spitsbergen: Adventfjorden, Barentsburg, Billefjorden, Brucebyen, Engelskhytta, "Großer Trichter", Grønfjorden, Longyearbyen, Magdalenefjorden, Salmon Lake at Dirksbukta, [7,91,99,101,103]; Svalbard [106]
42.	<i>Neomolgus capillatus</i> (Kramer, 1881)	Palearctic, Australasian	Spitsbergen: Hiorthhamn [91]
43.	<i>N. littoralis</i> (Linnaeus, 1758)	Holarctic	Bjørnøya: south [99,101,104,105]; Prins Karls Forland: Silene Hill [99,101]; Islands of Hinlopenstretet: Fosterøyane, Gyldenøyane, Vaigattøyane including Von Otterøya and Nordenskiöldøya [7]; Spitsbergen: Adventfjorden, Billefjorden, Bohemanneset, Brucebyen, Eremitten, Grønfjorden, between Hiorthhamn and Advent City, Isfjorden, Liefdefjorden, Ny-Friesland, Salmon Lake at Dirksbukta, , Sassenfjorden, Sorgfjorden, south side of Reinsdyrflya [7,91,99,101,102,105]
44.	<i>N. pallipes</i> (C.L. Koch, 1879)	Holarctic	Spitsbergen: Adventfjorden, Billefjorden, Brucebyen, south side of Reinsdyrflya [7,99,101]
	Cunaxidae		
45.	<i>Cunaxoides croceus</i> (C.L. Koch, 1838)	Holarctic	Spitsbergen: "Großer Trichter" [103]
	Cocceupodidae		
46.	<i>Cocceupodes mollicellus</i> C.L. Koch 1838	Holarctic	Spitsbergen: "Großer Trichter", Hiorthhamn, Longyearbyen, Magdalenefjorden [91,103,106]
	Eupodidae		
47.	<i>Eupodes variegatus</i> C.L. Koch, 1838	Holarctic, Afrotropical	Bjørnøya: Engelskelva, Fugleodden [91]; Spitsbergen: Hanaskogdalen, Hiorthhamn, Longyearbyen? [91,103,106]
48.	<i>Neoprotoreuneses borneri</i> (Thor, 1934) * ¹	Holarctic	Spitsbergen: "Großer Trichter", Magdalenefjorden, Tempelfjorden [66,103]
	Penthaleidae		
49.	<i>Penthaleus major</i> (Dugès, 1834)	Cosmopolitan	Bjørnøya [102]; Spitsbergen: Hiorthhamn, Longyearbyen, Magdalenefjorden, Reinsdyrflya-Liefdefjorden, Station Islands at Liefdefjorden [7,91,103]
	Penthalodidae		
50.	<i>Penthalodes ovalis</i> (Dugès, 1834)	Holarctic	Bjørnøya [105]; Spitsbergen: Barentsburg, Hiorthhamn, Liefdefjorden, Longyearbyen, south side of Reinsdyrflya [7,91,103]
	Rhagidiidae		
51.	<i>Coccorhagidia clavifrons</i> (Canestrini, 1886)	Cosmopolitan	Spitsbergen: "Großer Trichter", Hiorthhamn, Tempelfjorden [91,103]
52.	<i>Rhagidia gelida</i> Thorell, 1872 *	Holarctic	Bjørnøya [101,102,104,105], south [99]; Spitsbergen: Adventfjorden, Bellsund, Billefjorden, Brucebyen, Dirksbukta, Eremitten, "Großer Trichter", Hiorthhamn, Lomfjorden, Reinsdyrflya-Liefdefjorden, Salmon Lake at Dirksbukta, south side of Reinsdyrflya [7,91,99,101–103,105]
	Halacaridae		
53.	<i>Bradyagaue alberti</i> (Trouessart, 1902) *	Arctic	Spitsbergen: north [107]; Hopen [107]
54.	<i>Copidognathus poucheti</i> (Trouessart, 1893) *	Atlantic Ocean, Arctic	Spitsbergen: western coast [81]; expedition to Spitsbergen: Bellsund, Isfjorden, and Prins Karls Forland [108]
55.	<i>C. reticulatus</i> (Trouessart, 1893) *	Atlantic Ocean, Arctic	Spitsbergen: western coast [81]; expedition to Spitsbergen: Bellsund, Isfjorden, and Prins Karls Forland [108], Hinlopenstretet [107]
56.	<i>C. richardi</i> (Trouessart, 1902) *	Atlantic Ocean, Arctic	Spitsbergen: north [107]; Hopen [107]
57.	<i>Halacarellus subterraneus</i> Schulz,	Atlantic Ocean, Arctic,	Spitsbergen [45]

	1933	Palearctic	
58.	<i>H. subcrispus</i> Bartsch, 1978	Atlantic Ocean, Arctic	Spitsbergen [45]
59.	<i>Halacarus borealis</i> Trouessart, 1893 *	Atlantic Ocean, Arctic	Spitsbergen: western coast [81]; expedition to Spitsbergen: Bellsund, Isfjorden, and Prins Karls Forland [108]
60.	<i>Isobactrus levius</i> (Viets, 1927)	Atlantic Ocean, Arctic	Spitsbergen: Grønfjordbreen [109]
61.	<i>Rhombognathides</i> <i>spinipes</i> (Viets, 1933)	Atlantic Ocean, Arctic	Spitsbergen: Grønfjordbreen [109]
62.	<i>Rhombognathus subtilis</i> Bartsch, 1975	Atlantic Ocean, Arctic	Spitsbergen [45]
63.	<i>Thalassarachna coeca</i> (Trouessart, 1902) *	Arctic	Hopen [107]
64.	<i>T. princeps</i> (Trouessart, 1902) *	Arctic	Hopen [107]
	<hr/>		
65.	<i>Triophtydeidae</i> <i>Triophtydeus piniculus</i> (Oudemans, 1929) ²	Holarctic, Australasian	Spitsbergen: Magdalenefjorden [103]
	<hr/>		
66.	<i>Tydeidae</i> <i>Tydeus langei</i> Thor, 1934 ^{3,*,+†}	Arctic	Spitsbergen: "Großer Trichter", Magdalenefjorden, Tempelfjorden [103]
67.	<i>T. svalbardensis</i> Thor, 1932 ^{3,*,+†}	Arctic	Bjørnøya: Engelskelva, Fugleodden [91,110]; Spitsbergen: "Großer Trichter", Hanaskogdalen, Hiorthhamn, Longyearbyen, Magdalenefjorden, Tempelfjorden [91,103]
	<hr/>		
68.	<i>Iolinidae</i> <i>Tydaeolus globifer</i> (Thor, 1931)	Arctic	Spitsbergen: "Großer Trichter", Magdalenefjorden, Tempelfjorden [103]; Svalbard [47]
69.	<i>T. tenuiclaviger</i> (Thor, 1931)	Palearctic	Spitsbergen: Tempelfjorden [103]; Svalbard [47]
70.	<i>Microtydeus constans</i> Thor, 1931	Arctic	Spitsbergen: "Großer Trichter", Magdalenefjorden [103]; Svalbard [47]
	<hr/>		
71.	<i>Sperchontidae</i> <i>Sperchon brevirostris</i> (Koenike, 1895)	Holarctic	Bjørnøya: Ellasjøen, Engelskelva, Hellevatnet, Mosevatnet, Stevatnet [91,99,111]
	<hr/>		
72.	<i>Trombidiidae</i> <i>Podothrombium bicolor</i> (Hermann, 1804)	Palearctic	Spitsbergen: Hiorthhamn [91]
73.	<i>P. curtipalpe</i> (Thor, 1900) *	Holarctic	Spitsbergen: Hiorthhamn [91]
74.	<i>P. svalbarensis</i> Oudemans, 1930 ^{*,+†}	Arctic	Spitsbergen: Hiorthhamn [91]
	<hr/>		
75.	<i>Syringophilidae</i> <i>Chenophila nansenii</i> Skoracki & Zawierucha 2016 ^{*,+†}	Arctic	Spitsbergen: Nissenfjella [57]
	<hr/>		
76.	<i>Stigmaeidae</i> <i>Eustigmaeus oudemani</i> (Thor, 1930) ^{*,+†}	Arctic	Spitsbergen: Hiorthhamn [91]
77.	<i>E. pulchellus</i> (Thor, 1930) ^{*,+†}	Arctic	Spitsbergen: Hiorthhamn [91]
	<hr/>		
78.	<i>Tetranychidae</i> <i>Bryobia borealis</i> Oudemans, 1930 *	Holarctic	Spitsbergen: "Großer Trichter", Tempelfjorden [103]; Svalbard [106,112]
79.	<i>B. praetiosa</i> C.L. Koch 1836	Cosmopolitan	Bjørnøya [105]; Spitsbergen: Salmon Lake at Dirksbukta, south side of Reindsdyrflya [7]
	<hr/>		
80.	<i>Pygmephoridae</i> <i>Kerdabania arctica</i> (Thor, 1934) ^{4,*,+†}	Arctic	Spitsbergen: Templefjorden [103]
	<hr/>		
	SARCOPTIFORMES		
	<hr/>		
81.	<i>Endoostigmata</i> <i>Nanorchestidae</i> <i>Nanorchestes arboriger</i> (Berlese, 1904) ⁵	Palearctic	Spitsbergen: "Großer Trichter", Hiorthamn, Magdalenefjorden [91,103]
	<hr/>		
	<i>Alicorhagiidae</i>		

82.	<i>Alicorhagia clavipilus</i> (Thor, 1931)*	Arctic	Spitsbergen: "Großer Trichter", Hiorthhamn [103,113]
83.	<i>A. plumipilis</i> (Thor, 1931)*	Palearctic	Spitsbergen: "Großer Trichter" [103]
<hr/>			
84.	Eriophyidae <i>Aceria saxifragae</i> (Rostrup, 1900)	Holarctic	Spitsbergen: Hornsund [67]
85.	<i>Cecidophyes siedleckii</i> Kiedrowicz, Szydlo & Skoracka, 2016 *†	Arctic	Spitsbergen: Hornsund [67]
<hr/>			
Oribatida			
86.	Brachychthoniidae <i>Brachychthonius</i> <i>laetepictus</i> Berlese, 1910	Palearctic	Spitsbergen: Magdalenefjorden, Pyramiden [20,103]
87.	<i>Eobrachychthonius</i> <i>borealis</i> Forsslund, 1942	Holarctic	Bjørnøya [83]; Spitsbergen: Barentsburg, Longyearbyen, Recherchefjorden [14,114] Danskøya: Virgohamna [115]; Spitsbergen: Bockfjorden, Fivelflyene, Flathuken, "Großer Trichter", Helvetiadalen, Hiorthhamn, Isdammen, Longyearbyen, Magdalenefjorden, Reinsdyrflya, Sassendalen, Sven Olssonodden, Sørgattet [91,103,115,116]
88.	<i>E. latior</i> (Berlese, 1910)	Holarctic	Edgeøya: Kapp Lee, Russebukta [8] Spitsbergen: Longyearbyen, Pyramiden, Recherchefjorden [20,114,116]
89.	<i>E. oudemansi</i> Hammen, 1952	Holarctic, Neotropical, Indomalayan, Antarctic	Spitsbergen: Hornsund [88]
90.	<i>Liochthonius alpestris</i> (Forsslund, 1958)	Palearctic	Spitsbergen: Adventdalen, Barentsburg, Fivelflyene, Flathuken, Hiorthhamn, Isdammen, Reinsdyrflya, Worsleysenet [19,91,115,116]
91.	<i>L. brevis</i> (Michael, 1888)	Holarctic	Edgeøya: Blåbukta, Kapp Heuglin [8]
92.	<i>L. clavatus</i> (Forsslund, 1942)	Palearctic	Bjørnøya [83]; Spitsbergen: Barentsburg, Hornsund, Longyearbyen, Midtre Lovénbreen, Ny-Ålesund, Recherchefjorden [10,14,15,88,114,117,118]
93.	<i>L. lapponicus</i> (Trägårdh, 1910)	Holarctic	Spitsbergen: Barentsburg, Hornsund [19,88]
94.	<i>L. muscorum</i> Forsslund, 1964	Palearctic	Spitsbergen: Kongsfjorden [119]; based on Hodkinson's material stored in Liverpool [120]
95.	<i>L. neglectus</i> Moritz, 1976	Palearctic	Bjørnøya [83]; Hopen: Koefoedodden, Hopen radio [86]; Spitsbergen: Barentsburg, "Großer Trichter", Hiorthhamn, Longyearbyen, Magdalenefjorden, Petuniabukta, Pyramiden, Recherchefjorden, Vestpynten [14,19,20,22,91,103,105,114]
96.	<i>L. sellnicki</i> (Thor, 1930) *	Holarctic	Edgeøya: Blåbukta, Kapp Heuglin, Kapp Lee, Negerdalen, Russabukta [8]
97.	<i>L. strenzkei</i> Forsslund, 1963	Holarctic	Spitsbergen: Midtre Lovénbreen [10]
98.	<i>L. tuxeni</i> (Forsslund, 1957) <i>Neoliochthonius</i> <i>piluliferus</i> (Forsslund, 1942)	Palearctic	Edgeøya: Kapp Heuglin [8]
99.	<hr/>		
100.	Eniochthoniidae <i>Eniochthonius</i> <i>minutissimus</i> (Berlese, 1903)	Cosmopolitan	Spitsbergen: Petuniabukta [22]
<hr/>			
101.	Phthiracaridae <i>Atropacarus striculus</i> (C.L. Koch, 1835)	Holarctic, Oriental, Neotropical, Australian	Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen, Midtre Lovénbreen [10,13,14,16]
<hr/>			
102.	Crotoniidae <i>Camisia anomia</i> Colloff, 1993	Arctic	Bjørnøya [83]; Spitsbergen: Endalen, Kolhaugen, Kongsfjorden [121–129]
103.	<i>C. biverrucata</i> (C.L. Koch, 1839)	Holarctic	Edgeøya: Blåbukta, Kapp Heuglin [8]; Spitsbergen: Longyearbyen [103]
104.	<i>C. borealis</i> (Thorell,	Holarctic	Danskøya: Virgohamna [115]; Spitsbergen: Barentsburg, Hornsund,

			1871) *	
105.	<i>C. dictyna</i> Colloff, 1993	Holarctic	Isfjorden, Longyearbyen, Recherchefjorden [14,16,102,114,115,130] Svalbard [71]; Spitsbergen: Midtre Lovénbreen [10]	
106.	<i>C. foveolata</i> Hammer, 1955	Holarctic	Bjørnøya [83]; Edgeøya: Blåbukta, Kapp Heuglin, Negerdalen, Russebukta [8]; Hopen: Hopen radio, Koefoedodden[86]; Spitsbergen: Aldegondabreen, Grønfjordenbreen, Hornsund, Kiærstranda, Knudsenheia, Midtre Lovénbreen [10,88,109,131]	
107.	<i>C. horrida</i> (Hermann, 1804)	Holarctic, Oriental, Ethiopian, Neotropical	Bjørnøya [91]; Edgeøya: Diskobukta, Kapp Lee, Russebukta [8]; Spitsbergen: Barentsburg, "Großer Trichter", Hiorthamn, Hiorthjellet, Longyearbyen, Magdalenefjorden, Kongsfjorden, Reinsdyrflya, Recherchefjorden, Vestpynten, Petuniabukta, Pyramiden [20,22,91,103,105,114,115,132]	
108.	<i>C. invenusta</i> (Michael, 1888)	Palaearctic	Svalbard [71]	
109.	<i>C. lapponica</i> (Trägårdh, 1910)	Holarctic	Spitsbergen: Barentsburg, Hornsund, Longyearbyen [14,16,88]	
110.	<i>C. spinifer</i> (C.L. Koch, 1836)	Holarctic, Oriental, Neotropical	Spitsbergen: Tempelfjorden [103]	
111.	<i>Capillonothrus capillatus</i> (Berlese, 1914)	Holarctic	Spitsbergen: Arctowskifjellet [115]	
112.	<i>Platynothrus peltifer</i> (C.L. Koch, 1839)	Holarctic, Oriental, Australian	Spitsbergen: Recherchefjorden, Sassendalen [114,115]	
113.	<i>P. punctatus</i> (L. Koch, 1879)	Palaearctic	Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen, Petuniabukta, Vestpynten [13,14,16,22]	
Hermanniiidae				
114.	<i>Hermannia reticulata</i> Thorell, 1871 *	Holarctic	Bjørnøya: Fugleodden, Tunheim [83,91,105,133]; Edgeøya: Blåbukta, Russebukta [134]; Prins Karls Forland: Silene Hill [99,101]; Spitsbergen: Barentsburg, Bell sund, Endalen, "Großer Trichter", Kongsfjorden, Longyearbyen, Magdalenefjorden, Midtre Lovénbreen, Ny-Ålesund, Observasjonsholmen, Recherchefjorden, Petuniabukta, Tempelfjorden, Vestpynten [10,12,15,22,91,102,103,114,121–123,125,129,132–134]	
115.	<i>H. scabra</i> (L. Koch, 1879)	Holarctic	Bjørnøya [83,105]; Edgeøya: Kapp Lee [135]; Prins Karls Forland: Fuglehuken fyr [135]; Spitsbergen: Kapp Laila, Petuniabukta [135]	
Malaconothridae				
116.	<i>Malaconothrus monodactylus</i> (Michael, 1888)	Palaearctic	Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen [13,14,16]	
Nanhermanniidae				
117.	<i>Nanhermannia sellnicki</i> Forsslund, 1958	Palearctic	Spitsbergen: Barentsburg, Longyearbyen [14]	
Nothriidae				
118.	<i>Nothrus palustris</i> C.L. Koch, 1839	Holarctic	Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen [13,16]	
Damaeidae				
119.	<i>Damaeus onustus</i> (C.L. Koch, 1844)	Palaearctic, Afrotropical	Spitsbergen: Barentsburg, Longyearbyen [16]	
120.	<i>Kunstidamaeus arcticus</i> Miko & Monson, 2013 *, [†]	Svalbard	Bjørnøya [83]; Spitsbergen [72]	
121.	<i>Subbelba montana</i> (Kulczynski, 1902)	Palearctic	Bjørnøya: Austervåg [91]	
Peloppiidae				
122.	<i>Ceratoppia bipilis</i> (Hermann, 1804)	Holarctic, Oriental, Neotropical	Prins Karls Forland: Silene Hill [99,101]; Spitsbergen: Kongsfjorden, Storholmen [7;12] Bjørnøya: Tunheim [91]; Spitsbergen: Barentsburg, Bockfjorden, "Großer Trichter", Grønfjordenen, Hiorthjellet, Hiorthamn, Isdammen, Kolhaugen, Kongsfjorden, Longyearbyen, Magdalenefjorden, Recherchefjorden, Worsleyneset [91,103,114,115,121–125,127]	
123.	<i>C. hoeli</i> Thor, 1930 *	Arctic	Bjørnøya [83]; Edgeøya: Kapp Lee, Russebukta [8]; Hopen: Hopen radio, Koefoedodden [86]; Spitsbergen: Barentsburg, Hornsund, Longyearbyen, Petuniabukta, Pyramiden, Vestpynten [14–	
124.	<i>C. sphaerica</i> (L. Koch, 1879)	Palaearctic		

16,20,22,130]				
Carabodidae				
125.	<i>Carabodes labyrinthicus</i> (Michael, 1879)	Holarctic	Spitsbergen: Barentsburg, Longyearbyen [14]	
126.	<i>C. marginatus</i> (Michael, 1884)	Palearctic	Spitsbergen: Barentsburg, Longyearbyen [16]	
Autognetidae				
127.	<i>Autogneta kaisilai</i> Karppinen, 1967 *	Arctic	Spitsbergen: Biskayerhuken [115]	
128	<i>Conchogneta dalecarlica</i> (Forsslund, 1947)	Palearctic	Spitsbergen: Midtre Lovénbreen [10]	
Oppiidae				
129.	<i>Dissorrhina ornata</i> (Oudemans, 1900)	Holarctic	Bjørnøya: Fugleodden, Mosevatnet [91]; Spitsbergen: Adventdalen, Barentsburg, Biskayerhuken, Fivelflyene, Hiorthhamn, Isdammen, Longyearbyen, Magdalenefjorden, Petuniabukta, Pyramiden [14,20,22,91,103,115]	
130.	<i>Lauroppia fallax</i> (Paoli, 1908)	Holarctic, IndoMalayan, Australian, Neotropical	Spitsbergen: Magdalenefjorden [103]	
131.	<i>Micropia minus</i> (Paoli, 1908)	Cosmopolitan	Spitsbergen: Hornsund [136]	
132.	<i>Moritzella microdentata</i> Gordeeva & Grishina, 1991	Palearctic	Spitsbergen: Hornsund [137]	
133.	<i>Moritzoppia splendens</i> (C.L. Koch, 1841)	Holarctic	Bjørnøya [83]; Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen [13,14,16]	
134.	<i>M. unicarinata</i> (Paoli, 1908)	Holarctic	Edgeøya: Negerdalen [8]; Spitsbergen: Barentsburg, Longyearbyen [14,16] Edgeøya: Kapp Heuglin [8]; Spitsbergen: Barentsburg, "Großer Trichter", Hanaskogdalen, Hiorthhamn, Kongsfjord, Longyearbyen, Magdalenefjorden, Midtre Lovénbreen, Vestpynten [10,14,19,22,91,103,125]	
135.	<i>Oppiella neerlandica</i> (Oudemans, 1900)	Holarctic	Spitsbergen: Adventdalen, Barentsburg, Grønfjordenbreen, Hornsund, Isdammen, Isfjorden, Kongsfjorden, Longyearbyen, Magdalenefjorden, Midtre Lovénbreen, Vestpynten [13,14,16,19,22,115,121,123,136]	
136.	<i>O. nova</i> (Oudemans, 1902)	Cosmopolitan	Edgeøya: Diskobukta, Kapp Lee, Russebukta [8]; Hopen: Hopen radio, Koefoedodden [84]; Spitsbergen: Adventdalen, Arctowskijfjellet, Barentsburg, Bockfjorden, Fivelflyene, Grønfjordenbreen, Hornsund, Isfjorden, Kolhaugen, Longyearbyen, Magdalenefjorden, Petuniabukta, Recherchefjorden, Reinsdyrflya, Stuphallet, Vestpynten, Worsleyneset [13,14,16,22,103,114,115,124,136,138]	
Suctobelbidae				
138.	<i>Suctobelba hammerae</i> Krivolutsky, 1965	Holarctic	Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen [13,14,16]	
139.	<i>Suctobelbella sarekensis</i> (Forsslund, 1941)	Holarctic	Spitsbergen: Adventdalen, Fivelflyene [115]	
140.	<i>S. subcornigera</i> (Forsslund, 1941)	Holarctic, Oriental, Australian	Spitsbergen: Barentsburg, Biskayerhuken, Grønfjordenbreen, Isfjorden, Longyearbyen, Reinsdyrflya, Worsleyneset [13,14,16,22,115]	
Tectocephidae				
141.	<i>Tectocephus alatus</i> Berlese, 1913	Palearctic	Spitsbergen: Recherchefjorden [114]	
142.	<i>T. knuellei</i> Vanek, 1960	Palearctic	Spitsbergen: Barentsburg, Longyearbyen [14,16]	
143.	<i>T. sarekensis</i> Trägårdh, 1910	Palearctic	Spitsbergen: Adventdalen, Blomstrandhalvøya, Bockfjorden, Isdammen, Kongsfjorden, Reinsdyrflya, Worsleyneset [115]	
144.	<i>T. velatus</i> (Michael, 1880)	Cosmopolitan	Bjørnøya [83]; Edgeøya: Diskobukta, Kapp Lee, Russebukta [8]; Spitsbergen: Barentsburg, Hiorthhamn, Kolhaugen, Kongsfjorden, Longyearbyen, Midtre Lovénbreen, Petuniabukta, Pyramiden, Vestpynten [10,14,16,19,20,22,91,103,105,124,125]	
Ameronothridae				
145.	<i>Ameronothrus lineatus</i> (Thorell, 1871) *	Palaeartic	Bjørnøya: Engelskelva, Fugleodden, Tunheim [83,88,99,101,103,105,132,133,139]; Lågøya: [7]; Prins Karls Forland:	

			MacKenizedale, Richardlaguna almost to Carmichaelpynten [99,101]; Spitsbergen: Adventdalen, Barentsburg, Bockfjorden, Colesbukta, Grønfjordenbreen, Kolhaugen, Liefdefjorden, Nordenskiöld Land, Ny-Ålesund, Observasjonsholmen, Reinsdyrflya, Smeerenburg, Worsleyneset [7,12,91,102,109,115,124,132,133,140–145]
146.	<i>A. nidicola</i> Sitnikova, 1975	Palearctic	Spitsbergen: Hornsund [88]
147.	<i>A. nigrofemoratus</i> (L. Koch, 1879)	Palearctic	Bjørnøya [83]
	Micreremidae		
148.	<i>Micreremus brevipes</i> (Michael, 1888)	Palearctic	Spitsbergen: Recherchefjorden [114]
	Scutoverticidae		
149.	<i>Scutovertex minutus</i> (C.L. Koch, 1835)	Palearctic	Spitsbergen: Bockfjorden, Recherchefjorden [114,115]
	Achipteriidae		
150.	<i>Achipteria punctata</i> (Nicolet, 1855)	Holarctic	Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen, Vestpynten [13,14,16,22]
	Tegoribatidae		
	<i>Scutozetes</i>		
151.	<i>clavatosensillus</i> Ermilov, Martens & Tolstikov, 2013	Palearctic	Spitsbergen: Mosselbukta [146]
	Oribatellidae		
152.	<i>Oribatella arctica</i> Thor, 1930 *	Holarctic	Edgeøya: Diskobukta, Kapp Lee [8,147]; Spitsbergen: Barentsburg, Hiorthamn, Longyearbyen, Mälardalen, Petuniabukta, Recherchefjorden [15,22,91,103,114]
	Oribatulidae		
153.	<i>Oribatula exilis</i> (Nicolet, 1855)	Holarctic	Bjørnøya: Fugleodden, Mosevatnet, Tunheim [91]; Prins Karls Forland: Silene Hill [99,101]; Spitsbergen: Adventdalen, Barentsburg, Engelskhytta, Hanaskogdalen, Hiorthamn, Isfjord radio, Longyearbyen [14,91]
154.	<i>O. tibialis</i> (Nicolet, 1855)	Holarctic	Bjørnøya [83]; Edgeøya: Diskobukta, Kapp Lee, Russebukta [8]; Spitsbergen: Barentsburg, Longyearbyen, Observasjonsholmen, Petuniabukta, Storholmen, Vestpynten [12,14,15,19,22]
155.	<i>O. venusta</i> Berlese, 1908	Holarctic	Bjørnøya [83]; Spitsbergen: Adventdalen, Fivelflyene, "Großer Trichter", Longyearbyen, Magdalenefjorden, Recherchefjorden, Reinsdyrflya, Worsleyneset [103,114,115]
156.	<i>Phauloplia lucorum</i> (C.L. Koch, 1841)	Palearctic	Spitsbergen: Bellsund, Isfjorden [81,133]
	Scheloribatidae		
157.	<i>Liebstadia similis</i> (Michael, 1888)	Holarctic, Indomalayan, Australian	Spitsbergen: Barentsburg, Grønfjordenbreen, Isfjorden, Longyearbyen [13,14,16]
	Ceratozetidae		
158.	<i>Ceratozetes spitsbergensis</i> Thor, 1934 *	Palearctic	Edgeøya: Blåbukta, Kapp Heuglin, Negerdalen [8]; Kong Karls Land [148]; Spitsbergen: Arctowskijfjellet, Barentsburg, "Großer Trichter", Kolhaugen, Longyearbyen, Magdalenefjorden, Pyramiden, Reinsdyrflya, Worsleyneset [14,20,103,115,124] Bjørnøya: Fugleodden, Tunheim [14–16,83,91,99,101,105,133]; Barentsøya [130]; Edgeøya: Blåbukta, Diskobukta, Kapp Heuglin, Kapp Lee, Negerdalen, Russebukta, Tjuvfjorden [8,146]; Hopen: Hopen radio, Koefoedodden [86]; Prins Karls Forland: Fuglehukken, MacKenizedale, Silene Hill [99,101,146]; Sofiaøya [146]; Spitsbergen: Adventfjorden, Barentsburg, Bellsund, Billefjorden, Blomstrandhalvøya, Bohemanneset, Dirksbukta, Endalen, Engelskhytta, "Großer Trichter", Grønfjorden, Hiorthfjellet, Hiorthamn, Hornsund, Isfjorden, Kolhaugen, Krykkjefjellet, Liefdefjorden, Longyearbyen, Magdalenefjorden, Midtre Lovénbreen, Ossian Sarsfjellet, Petuniabukta, Recherchefjorden, Reinsdyrflya, Sven Olssonodden, Storholmen, Vestpynten, Worsleyneset [7,10,12,14–17,19,22,91,99,101–103,114,115,121–125,129,130,132,133,146,148]
159.	<i>Diapterobates notatus</i> (Thorell, 1871) *	Holarctic	

160.	<i>Edwardzetes edwardsi</i> (Nicolet, 1855)	Holarctic	Spitsbergen: Tempelfjorden; possibly from the North Cape [103]
161.	<i>Fuscozetes coulsoni</i> A. & S. Seniczak, 2020 * [†]	Arctic	Edgeøya: Negerdalen [9]
162.	<i>Iugoribates gracilis</i> Sellnick, 1944	Holarctic	Spitsbergen: Kongsfjorden [119]; based on Hodkinson's material stored in Liverpool [120]
163.	<i>Oromurcia lucens</i> (L. Koch, 1879)	Holarctic	Barentsøya [130]; Spitsbergen: Vestpynten [22]
164.	<i>Svalbardia paludicola</i> Thor, 1930 *	Holarctic	Spitsbergen: Hanaskogdalen, Hiorthhamn, Hornsund, Reinsdyrflya, Ringhorndalen, Sørkapp [91,139,146]
165.	<i>Trichoribates berlesei</i> Jacot, 1929	Holarctic	Spitsbergen: Hornsund, Storholmen [12,88]
166.	<i>T. novus</i> (Sellnick, 1928)	Holarctic	Spitsbergen: Hornsund [88]
167.	<i>T. setiger</i> (Trägårdh, 1910)	Palaearctic	Bjørnøya [83]; Spitsbergen: Recherchefjorden [114]
<hr/>			
168.	Chamobatidae <i>Chamobates birulai</i> (Kulczynski, 1902) *	Palaearctic	Spitsbergen [133]
169.	<i>C. borealis</i> (Trägårdh, 1902)	Palaearctic	Spitsbergen: Hornsund [88]
<hr/>			
170.	Punctoribatidae <i>Mycobates bicornis</i> (Strenzke, 1954)	Palaearctic	Spitsbergen: Midtre Lovénbreen [10]
171.	<i>M. parmeliae</i> (Michael, 1884)	Holarctic	Spitsbergen: Barentsburg, Longyearbyen, Midtre Lovénbreen [10,14,16] Bjørnøya [91]; Edgeøya: Kapp Lee, Russebukta [8,149]; Spitsbergen: Adventdalen, Arctowskijellet, Bockfjorden, Engelskhytta, Fjortende Julibukta, "Großer Trichter", Hiorthfjellet, Hiorthhamn, Hornsund, Isdammen, Kongsfjorden, Longyearbyen, Magdalenefjorden, Mosselbukta, Recherchefjorden, Sassendalen, Signehamna, Storholmen, Sørkapp, Vestpynten [22,91,103,114,115,121,123,125,149]
172.	<i>M. sarekensis</i> (Trägårdh, 1910)	Holarctic	
173.	<i>M. tridactylus</i> Willmann, 1929	Holarctic	Spitsbergen: Barentsburg, Longyearbyen [14]
174.	<i>Punctoribates punctum</i> (C.L. Koch, 1839)	Holarctic, Oriental, Australian	Spitsbergen: Barentsburg, Longyearbyen, Petuniabukta [14,16,22]
<hr/>			
175.	Galumnidae <i>Pergalumna nervosa</i> (Berlese, 1914)	Holarctic	Spitsbergen: Hornsund [136]
<hr/>			
176.	Acaridae <i>Sancassania mycophagus</i> (Mégnin, 1874)	Cosmopolitan	Spitsbergen: Hornsund [91]
<hr/>			
177.	Alloptidae <i>Alloptes (Sternalloptes)</i> <i>stercorarii</i> Dubinin, 1952	Arctic	Spitsbergen: Ny-Ålesund [74]
<hr/>			
178.	Avenzoariidae <i>Zachvatkinia isolata</i> Mironov, 1989	Arctic, Neotropical, Australian, Afrotropical	Spitsbergen: Ny-Ålesund [74]

Note: *—new to science; **—redescription on the base of material from Svalbard; †—so far found only on Svalbard.¹ [150] proposed the genus *Neoproteunetes* for those species of "*Protoreunetes*" that remained in the Eupodidae after the type specimen of *Protoreunetes* (*P. agilis* Berlese, 1923) was transferred to *Ereynetes* [66]. It is implied that this new genus includes *Protoreunetes borneri* Thor, 1934, but the combination was not published.² Identity and generic affiliation of this species is questionable [46]; possible record from Greenland with inconclusive identification [48].³ Species not included in [50], *species inquirenda?*⁴ Inadequately described, not included in key to species [61].⁵ Although specimens from Svalbard were apparently examined, this species was not confirmed from Svalbard [65].




Figure 2. Diversity of the mite families in Svalbard: (A) Mesostigmata; (B) Trombidiformes; (C) Sarcoptiformes. The number of species occurring in Svalbard is presented on bars. No figure was made for the Ixodida, which is represented in Svalbard by one family only.

More than a half of the Mesostigmata species known from Svalbard have been recorded there only once (Table 2). The majority of these are recent findings, but four were reported only in the first half of the last century, including *Arctoseius laterincisus*, which has an Arctic distribution. Four other records come from the second half of the last century, including another species with an Arctic distribution—*Halolaelaps gerlachi*.

Table 2. Mite species found on Svalbard only once (or by the same author at a similar time).

Until 1950	1951–2000	After 2001	Reference
Mesostigmata			
<i>U. acuminatus</i>			[81]
	<i>Z. curiosus</i>		[85]
<i>Z. triangularis</i>			[91]
	<i>P. (A.) insertus</i>		[18]
	<i>V. immanis</i>		[92]
	<i>V. remberti</i>		[18]
	<i>D. foveolatus</i>		[20]
	<i>H. coulsoni</i> *		[93]
	<i>H. gerlachi</i> *		[94]
	<i>S. baloghi</i>		[8]
<i>G. borealis</i>			[7]
	<i>M. muscaedomesticae</i>		[84]
	<i>A. cetratus</i>		[96]
<i>A. laterincisus</i> *			[91]
	<i>A. ornatus</i>		[96]
		<i>N. grumantensis</i> *	[28]
	<i>L. hilaris</i>		[100]
Trombidiformes			
<i>B. semiscutata</i> *			[91,103]
<i>N. capillatus</i>			[91]
<i>C. croceus</i>			[103]
	<i>A. saxifragae</i>		[67]
<i>P. borneri</i> *			[103]
<i>P. maior</i>			[102]
<i>C. clavifrons</i>			[91,103]
<i>B. alberti</i> *			[107]
<i>C. poucheti</i> *			[81]
<i>C. reticulatus</i> *			[107]
<i>C. richardi</i> *			[107]
	<i>H. subterraneus</i>		[45]
	<i>H. subcrispus</i>		[45]
<i>H. borealis</i> *			[81]
	<i>I. levis</i>		[109]
	<i>R. spinipes</i>		[109]
	<i>R. subtilis</i>		[45]
<i>T. coeca</i> *			[107]
<i>T. princeps</i> *			[107]
<i>T. globifer</i> *			[103]
<i>T. tenuiclaviger</i> *			[103]
<i>M. constans</i> *			[103]
<i>T. langei</i> *			[103]
<i>P. bicolor</i>			[91]
<i>P. curtipalpe</i> *			[91]
<i>P. svalbardense</i> *			[91]
<i>E. oudemani</i> *			[91]
<i>E. pulchellus</i> *			[91]
	<i>C. nansenii</i> *		[57]
<i>K. arctica</i> *			[103]
Sarcophagiformes			
<i>A. clavigipilus</i>			[103,113]

<i>A. plumipilis</i> *		[103]
	<i>C. siedleckii</i> *	[67]
<i>L. alpestris</i>		[88]
	<i>L. clavatus</i>	[8]
	<i>L. neglectus</i>	[119]
	<i>L. tuxeni</i>	[10]
<i>C. spinifer</i>		[103]
	<i>C. capillatus</i>	[115]
	<i>N. sellnicki</i>	[14]
	<i>D. onustus</i>	[16]
<i>S. montana</i>		[91]
	<i>C. labyrinthicus</i>	[14]
	<i>C. marginatus</i>	[16]
<i>A. kaisilai</i> *		[115]
	<i>C. dalecarlica</i>	[10]
<i>L. fallax</i>		[103]
	<i>M. minus</i>	[136]
	<i>S. sarekensis</i>	[115]
	<i>T. alatus</i>	[114]
	<i>T. sarekensis</i>	[115]
	<i>A. nidicola</i>	[88]
	<i>A. nigrofemoratus</i>	[83]
	<i>S. clavatosensillus</i>	[146]
<i>E. edwardsi</i>		[103]
	<i>F. coulsoni</i> *	[9]
	<i>I. gracilis</i>	[119]
	<i>T. novus</i>	[88]
	<i>C. birulai</i> *	[133]
	<i>C. borealis</i>	[88]
	<i>M. bicornis</i>	[10]
<i>P. nervosa</i>		[136]
<i>S. mycophagus</i>		[91]
	<i>A. stercorellii</i>	[74]
	<i>Z. isolata</i>	[74]

Note: *—new to science.

Most of the Mesostigmata species known from Svalbard have a Palaearctic distribution (Figure 3). Arctic species are also abundant and comprise nearly 40% of Svalbards total mesostigmatic diversity, while Holarctic species are less numerous. Five species have so far been found only in Svalbard (Table 1), including recent records (*Halolaelaps coulsoni* and *Neoseiulus grumantensis*) and historic records (*Halolaelaps gerlachi*, *Arctoseius laterincisus* and *Neoseiulus magnanalis*). Mesostigmata have been recorded from five islands in the Svalbard archipelago (Figure 1). Although most of the species are known from Spitsbergen, *Saproscans baloghi* Karg, 1964, is known only from Edgeøya and *Zerconopsis muestairi* (Schweizer, 1949) has been found exclusively on Bjørnøya (Table 1).




Figure 3. Distribution of mite species occurring in Svalbard: (A) Mesostigmata; (B) Trombidiformes; (C) Sarcoptiformes. No figure was made for the Ixodida, which are represented in Svalbard by one species with a distribution in “at least two regions” (see Table 1 for details).

3.3. Trombidiformes

The order Trombidiformes is represented in Svalbard only by the suborder Prostigmata with 17 families and 43 species recorded. The most diverse are the marine Halacaridae (12 spp.), followed by the terrestrial Bdellidae (7 spp.); these two families together comprise 44% of all the Trombidiformes known from Svalbard (Figure 1).

A strikingly large number of new species (19) have been described from Svalbard; i.e., nearly half of all Trombidiformes known from this archipelago: *Bdella semiscutata* Thor, 1930; *Neoprottereunetes borneri* (Thor, 1934); *Rhagidia gelida* Thorell, 1872; *Bradyagae alberti* (Trouessart,

1902); *Copidognathus poucheti* (Trouessart, 1893); *C. reticulatus* (Trouessart, 1893); *C. richardi* (Trouessart, 1902); *Halacarus borealis* Trouessart, 1893; *Thalassarachna coeca* (Trouessart, 1902); *T. princeps* (Trouessart, 1902); *T. langei* Thor, 1934; *T. svalbardensis* Thor, 1932; *Podothrombium curtipalpe* (Thor, 1900); *P. svalbardense* Oudemans, 1930; *Chenophila nansenii* Skoracki & Zawierucha, 2016; *Eustigmaeus oudehami* (Thor, 1930); *E. pulchellus* (Thor, 1930); *Bryobia borealis* Oudemans, 1930; and *Kerdabania arctica* (Thor, 1934). Moreover, 85% of the new species described from Svalbard have been recorded only once and mostly from early studies of the mite fauna (Table 2).

Species with wide geographical distributions predominate; cosmopolitan and Holarctic species, and those present in at least two zoogeographic regions (except that the Holarctic is treated as one unit) together comprise 66% of all the Trombidiformes (Figure 2). Arctic species are also abundant representing 27% of the recorded species, while the fewest number of species have Palaearctic distributions. There are seven species which have only been recorded from Svalbard (Table 1), but these are mostly single old records of Sig Thor, including *Tydeus langei*, *T. svalbardensis*, *Podothrombium svalbardense*, *Eustigmaeus oudehami*, *E. pulchellus*, and *Kerdabania arctica*. The one exception is the recently described *Chenophila nansenii*.

Trombidiformes have been found on five islands/island groups, predominantly on Spitsbergen (40 species), with others collected from Bjørnøya (11), Prins Karls Forland (6), Hopen (4), and a single species on the small islands of the Hinlopenstretet. A large number of species (27) have been found exclusively on Spitsbergen, one species [*Sperchon brevirostris* (Koenike, 1895)] only on Bjørnøya and two others (*Thalassarachna coeca* and *T. princeps*) only on Hopen (Table 1).

3.4. Sarcoptiformes

In Svalbard, 98 species of Sarcoptiformes from two suborders (Endeostigmata with 5 species and Oribatida with 93 species) have been recorded (Table 1). They represent 33 families; the richest in species is the Brachychthoniidae (14 spp.), followed by the Crotoniidae (12 spp.), Ceratozetidae (10 spp.), Oppiidae (9 spp.), and Punctoribatidae (5 spp.). Together, these five families comprise 51% of the species diversity of Sarcoptiformes in Svalbard (Figure 1). Some families are represented by only 1–4 species, yet this constitutes a significant share of their known global diversity. For example, 22% of the species belonging to the endeostigmatid family Alicorhagiidae occur in Svalbard. The oribatid families Eniochthoniidae, Peloppiidae and Tectocepheidae are also represented by large proportions of their total known species diversity, with 17%, 27%, and 24%, respectively.

Sixteen new species of Sarcoptiformes have been described from Svalbard: *Alicorhagia clavipilus* (Thor, 1931); *A. plumipilis* (Thor, 1931); *Cecidophyes siedleckii* Kiedrowicz, Szydlo & Skoracka, 2016; *Liochthonius sellnicki* (Thor, 1930); *Camisia borealis* (Thorell, 1871); *Hermannia reticulata* Thorell, 1871; *Kunstidamaeus arcticus* Miko & Monson, 2013; *Ceratoppia hoeli* Thor, 1930; *Autogneta kaisilai* Karppinen, 1967; *Ameronothrus lineatus* (Thorell, 1871); *Oribatella arctica* Thor, 1930; *Ceratozetes spitsbergensis* Thor, 1934; *Diapterobates notatus* (Thorell, 1871); *Fuscozetes coulsoni* A. & S. Seniczak, 2020; *Svalbardia paludicola* Thor, 1930; and *Chamobates birulai* (Kulczynski, 1902).

Species with wide geographic ranges together form 58% (cosmopolitan, 5%; Holarctic, 42%; and those found in at least two regions, 11%) of the sarcoptiform species (Figure 2). Species with a Palaearctic distribution also make up a substantial proportion (33%), with the remaining species having an Arctic distribution. Two of these Arctic species have only been found in the north of Spitsbergen: *Autogneta kaisilai* in Biskayerhuken and *Scutozetes clavatosensillus* Ermilov, Martens & Tolstikov, 2013, in Mosselbukta; and *Fuscozetes coulsoni* exclusively on Edgeøya (Table 1, Figure 3). Three species have been recorded only from Svalbard and these are all recently described new species (Table 1).

Sarcoptiformes have been recorded from ten islands in the Svalbard archipelago, but not on Nordaustlandet. The most commonly collected species is *Diapterobates notatus* (Table 1) found on seven islands and at nearly all collecting localities. Five species [*Camisia foveolata* Hammer, 1955; *Hermannia reticulata*, *H. scabra* (L. Koch, 1879); *Ceratoppia sphaerica* (L. Koch, 1879); and *Ameronothrus lineatus* (Thorell, 1871)] have been found on four islands and another seven other species [*Camisia*

horrida (Hermann, 1804); *Mycobates sarekensis* (Trägårdh, 1910); *Ceratozetes spitsbergensis*; *Oppiella translamellata* (Willmann, 1923); *Tectocepheus velatus* (Michael, 1880); *Oribatula exilis* (Nicolet, 1855); and *O. tibialis* (Nicolet, 1855)] on three islands. All of these species also have a wider distribution on Spitsbergen (Table 1, Figure 3).

Some species have been found exclusively on one island (Table 1), including 57 found only on Spitsbergen, four [*Liochthonius clavatus* (Forsslund, 1942); *L. strenzkei* Forsslund, 1963; *Neoliochthonius piluliferus* (Forsslund, 1942); and *Fuscozetes coulsoni*] on Edgeøya, and two [*Subbelba montana* (Kulczynski, 1902) and *Ameronothrus nigrofemoratus* (L. Koch, 1879)] from Bjørnøya.

Approximately one third of the sarcoptiform species have been recorded in Svalbard only once (Table 2). Seven of them were found in the first half of the last century and, 12 others, in the second half of the last century, including the descriptions of four new species.

4. Discussion

The acarofauna of the Svalbard archipelago is diverse with 178 recorded species comprising one Ixodida, 36 Mesostigmata, 43 Trombidiformes, and 98 Sarcoptiformes. However, compared with other regions of the Arctic (Alaska, Greenland, Iceland, and Taimyr), the species diversity is lower [151]. This paucity is likely to be explained by a combination of the small area of Svalbard (60% covered by permanent snow or glaciers) [4], the greater geographic isolation, and in some cases, the more northerly locality and harsher climate. As an example, from Iceland there are 428 species of mites known, with eight Ixodida, 131 Mesostigmata, 101 Trombidiformes, and 188 Sarcoptiformes [151–153]. Although Iceland has an area only 1.5 times greater than Svalbard, the island lies at a lower latitude (between 63° and 66° north latitude), glaciers cover only 11% of its area, and it lies in the path of the North Atlantic Current, which results in a more ameliorated climate. The diversity of mites in Svalbard is also much poorer in comparison to mainland Norway: Ixodida comprise 8% here with Mesostigmata—15%, Trombidiformes—13%, and Sarcoptiformes—30% [154–157]. However, these differences are not unexpected considering that the Svalbard archipelago is situated 900 km from the northern border of mainland Norway and has an area of only one fifth of that of mainland Norway [158].

The geographical isolation of islands, as a rule, results in reduced biodiversity [159], but, on the other hand, nurtures unique endemic species [160]. Svalbard has a unique flora and fauna, including endemic invertebrates: three rotifers, four tardigrades, and two aphids [86]. Fifteen mite species have been found exclusively in Svalbard, six of them being recent observations. The remaining are single historic records (i.e., from the first half of the last century) and it is consequently uncertain if any of these are endemic to Svalbard or the result of taxonomic confusion. Finding and studying the types or neotypes in Svalbard, along with more extensive sampling in the Arctic and including molecular studies, could help resolve this question.

Nevertheless, the Svalbard acarofauna is unique due to its specific Arctic elements. Amongst the Svalbard Mesostigmata in particular there are many species with an Arctic distribution (which make up nearly 40% of this group) and are similar to that observed on Greenland [151]. By contrast, the Sarcoptiformes in Svalbard are predominated by species with wide geographical ranges extending beyond the Arctic, as also seen with the Sarcoptiformes species known from Greenland [151] and northern Russia [161]. This difference between the two orders may be explained by the younger geological age of the Mesostigmata and their faster evolution rate [162]. Similarly, the Trombidiformes, which are also an ancient group [163], are dominated in Svalbard by species with wide geographic distributions. Nonetheless, Arctic species seem to be abundant among the Trombidiformes of Svalbard, comprising 27% of the species recorded, but these data should be treated with caution because they are mostly based on historic records that need to be confirmed.

Another characteristic feature of the Svalbard acarofauna is the large number of species new to science described from this region—they form approximately 25% of all mite species known from Svalbard. Seven of these species belong to the Mesostigmata, 19 to Trombidiformes, and 16 to Sarcoptiformes. However, many of these new species were found only once and early in the study of the mite fauna of Svalbard. These species include one mesostigmatid species, 18 Trombidiformes

species (i.e., 90% of all new species from this group), and two Sarcoptiformes species. Confirming these identifications is complicated by outmoded descriptions, incomplete, or too general drawings, and by the unavailability of most type material. These species require special attention in future studies. They are possibly rare or have been wrongly identified but the possibility cannot be excluded that they have disappeared from Svalbard due to environmental changes or that they were introduced and their populations have not established in Svalbard.

Out of 36 mite species found in Svalbard only once by early workers, the majority of species (24) were collected by Sig Thor, including one new species of Mesostigmata, eight new species of Trombidiformes, and two new species of Sarcoptiformes (Table 2). Despite the widespread belief that the collection of Thor along with all his valuable type material was destroyed in accordance with Thor's wishes (see, e.g., remark 2 on page 1308 in [148]), this collection exists and is kept at the Natural History Museum in Oslo, Norway [164]. However, it is in a very poor condition and it is not certain whether the Svalbard material is in a suitable condition for re-examination.

Fortunately, some of Thor's species have been found by later sampling campaigns and redescribed, for instance, *Antennoseius (Vitzthumia) oudeansi*, *Neoseiulus magnanalis*, *Proctolaelaps parvanalis*, and the specimens obtained deposited in zoological collections as neotypes [28,79]. In addition, six new species of mites have been found recently in Svalbard, including two Mesostigmata—*Neoseiulus grumantensis* and *Halolaelaps coulsoni* [28,93], one Trombidiformes—*Chenophila nansenii* [57], and three Sarcoptiformes—*Cecidophyes siedleckii*, *Kunstidamaeus arcticus*, and *Fuscozetes coulsoni* [9,67,72].

Recently, the first species of Ixodida, the seabird tick (*Ixodes uriae*), was discovered in Svalbard [76–78]. This tick is a major parasite of seabirds breeding at high latitudes and has the most extensive geographical distribution of all tick species [23], being also common throughout mainland Norway [165]. Its increasing occurrence in Svalbard may be related to warming winters (the tick overwinters at the breeding sites of the seabirds) [78] as was similarly observed in Iceland with another tick species, *I. ricinus* Linnaeus, 1758, and which has become more common in recent years [153].

Studies conducted during the past 20 years have also added 16 species of Mesostigmata [8,18,20,27,28,30,85,92,93,95,98,166] and 36 species of Sarcoptiformes to the Svalbard fauna [8–10,67,71,72,74,146,167]. This indicates that despite the relatively long history of mite studies for a region in the Arctic, our knowledge remains surprisingly poor. It is striking, however, that the Trombidiformes have been much less studied in Svalbard than other mite groups (Table 1). The reference list presented here includes only 18 papers referring to the Trombidiformes (vs. 72 papers on Sarcoptiformes and 34 on Mesostigmata), most of which were published at the end of the 19th and first half of the 20th centuries. Only three come from the present century. Therefore, it is clear that in future studies this group in particular should receive more attention.

There are also some species in Svalbard that have extraordinary disjunct distribution patterns: they occupy mainly the Arctic and parts of the subarctic regions, but are also found in some distant localities, in harsh conditions. One example is an oribatid species, *Ceratozetes spitsbergensis*, which has been assumed to be an Arctic species, present in Svalbard, Alaska [148], northern Canada [168,169] and the Nordic Arctic of the Russian Far East [148,170], but was found also in the Altai Mountains in Mongolia, at an altitude of 2800–3200 m a.s.l. [171] and in the Alps in Austria, at an altitude of 3300 m a.s.l. [172]. Since it was not found at lower elevations and/or lower latitudes, its presence in the Alps was explained by one of these theories: relict distribution [a cold-adapted species that was widely distributed in Europe during the Last Glacial Maximum (LGM), but when conditions became warmer it retreated to very restricted areas at high altitudes], or the nunatak hypothesis (survival of species on ice-free refuges) [172]. Some studies suggest that parts of Svalbard, e.g., Amsterdamsøya (north-west of Svalbard), remained ice-free during the LGM [173]. It is thus possible that some invertebrates survived the LGM *in situ* in these glacial refugia, although, due to the harsh conditions prevailing over an extended period of time, it seems likely that most biota could probably not survive on nunataks [86]. Another oribatid mite with an interesting distribution is *Scutozetes clavatosensillus*, which has been found only in Svalbard and in Nepal

(Dhaulagiri massif, 3200–3600 m a.s.l.); its distribution may also be related to glacial history. Similarly to *S. clavatosensillus*, the mesostigmatic species, *Paragamasus (Aclerogamasus) insertus*, may also be a glacial relict since it is known only from the Gory Stolowe Mtns., Poland [174,175], where it occurred in rock crevices in very extreme conditions with long-lasting snow cover, and recently collected in Svalbard [18].

The composition of the present acarofauna of Svalbard likely results mostly from postglacial immigration [86]. The mites, being wingless, have rather low dispersal abilities. However, phoresy with migrating birds [13–17] or insects [18,95], and transport on driftwood or even direct dispersal in or on seawater with ocean currents [11], are possible dispersal pathways from the mainland to, or between, remote islands such as the Svalbard archipelago. Humans may also play a role in the dispersal of mites; as with the import of fertile agricultural soils transported from southern Russia and contemporary Ukraine to the Russian settlements on Spitsbergen (Barentsburg and Pyramiden) to enrich the soils of the greenhouses and grass lawns [18,19]. Mite communities in such transformed microhabitats differ noticeably from those of the adjacent Arctic tundra. Moreover, manure from abandoned cow sheds provided specific ameliorated environmental conditions enabling the survival of terrestrial invertebrate species not yet recorded elsewhere in Svalbard [18–20]. Introduction of new species to islands can be deleterious since the island communities may be disrupted, often resulting in the extinction of their endemic species [176–178].

Reconstructing the colonization of Svalbard after the LGM may be attempted by observing primary succession before retreating glaciers. For example, at the Midtre Lovén glacier foreland, two oribatid species, *Camisia foveolata* and *Tectocepheus velatus*, were the first colonizers. Mesostigmata appeared later, *Proctolaelaps parvanalis* being the first species recorded at this glacier foreland [10]. All these species are widely distributed in different parts of the archipelago (see Table 1). *Tectocepheus velatus* was also the earliest colonizer on a glacier foreland in southern Norway [179], while *Camisia foveolata* was one of the first colonizers on geothermally active lava fields in Iceland [180]. Oribatida are mainly saprophagous but species that are fungivores, bacteriovores, algivores, or omnivores, such as *T. velatus*, can find the appropriate food on seemingly bare ground [181]. In turn, the development of the first animal communities provides prey to the predatory Mesostigmata.

As shown in the present review, acarological studies of Svalbard are heavily geographically biased since most sampling has been carried out on the largest island—Spitsbergen; 90% of papers refer to this island and 90% of species have been found there (see Table 1). Almost 20% of papers refer to Bjørnøya and 25% of the species total for the archipelago have been found there, while 7% of papers refer to Edgeøya and 17% of the species are known from there. Nine other islands and island groups have been studied to a much lesser extent, with single records from other locations. It needs to be emphasized that some mites have been found exclusively on one island, other than Spitsbergen: five species on Edgeøya (*Saproscans baloghi*, *Liochthonius clavatus*, *L. strenzkei*, *Neoliochthonius piluliferus*, and *Fuscozetes coulsoni*), four on Bjørnøya (*Zerconopsis muestairi*, *Sperchon brevirostris*, *Subbelba montana*, and *Ameronothrus nigrofemoratus*), and two on Hopen (*Thalassarachna coeca* and *T. princeps*), including three species new to science. It is unclear why these species have not been collected from Spitsbergen despite the more comprehensive sampling efforts on this island. The importance of further studies in different parts of the Svalbard archipelago is highlighted, in particular since the archipelago is extremely diverse geographically and climatically.

In natural conditions, the species composition and abundance of mites depends mainly on the vegetation [89], which, in turn, depends largely on the climate. For example, a relatively high mesostigmatid diversity is present along the western coast of Spitsbergen (about 30 species) [86], which experiences a comparatively mild climate for the latitude, in contrast to the polar deserts, where only five species were recorded [30]. Within the same climatic conditions, the densities of the mites also vary greatly according to the vegetation types [22,30,88,89,136].

Since mites are connected in different ways with other parts of the ecosystem, it would be particularly interesting to apply a multi-disciplinary approach to trace the effect of climate change in Svalbard. For example, one of the understudied mite groups is the family Eriophyidae. This

family includes phytophagous species of great economic importance and with high invasive potential, and could be very useful for ecological studies on the effects of a changing climate at Svalbard [67]. Another poorly known group is the hyperorder Astigmata, parasitizing birds [74]. It is well-documented that climatic changes are affecting the diversity of seabirds by changing their foraging and breeding ecology, as well as increasing the abundance of temperate species [6], which could in turn affect bird-associated Astigmata.

Attention should also be paid to the least known mite order in Svalbard, the Trombidiformes, which is extremely diverse with respect to their feeding preferences (this taxon includes algivores, bacterivores, fungivores, predators, and parasites), and occupies terrestrial, freshwater, and marine habitats [182], but has not been studied at all from an ecological perspective in Svalbard.

Acarological research in the Arctic has developed rapidly in the last 20 years, markedly contributing to the knowledge of the mite communities and increasing somewhat our understanding of the factors determining these communities. Nonetheless, little is understood about the physical and morphological adaptations of mites, their adaptations to the extreme Arctic environment, or the genetic biodiversity of these isolated populations. Moreover, research has focused on the larger and more easily accessible regions of the archipelago to the detriment of the more environmentally extreme eastern and northern regions. There has also been a focus on the Sarcoptiformes and Mesostigmata while the Trombidiformes have been neglected. These areas will remain the subject of our research in the forthcoming years.

Author Contributions: Conceptualization, A.S. and D.J.G.; Investigation: Mesostigmata—D.J.G.; Trombidiformes—M.D.S.; Sarcoptiformes—A.S., S.S.; Svalbard peculiarities—S.J.C.; Methodology, A.S., S.S., M.D.S., S.J.C. and D.J.G.; Visualization, A.S.; Writing—original draft, review & editing, A.S., S.S., M.D.S., S.J.C. and D.J.G. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Norwegian Taxonomy Initiative: Knr 6-20.

Acknowledgments: We are grateful to Jacek Dabert (Institute of Environmental Biology, Adam Mickiewicz University, Poznań, Poland) for information about distribution of species of Astigmata, and to Reinhard Gerecke (Institute for Evolution and Ecology, University of Tübingen, Tübingen, Germany) for information about Thor's collection. This study was partly supported by the grant from The Norwegian Taxonomy Initiative – Knr 6-20, Mites of Atlantic Raised Bogs in Norway (MARB).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gillespie, M.A.K.; Alfredsson, M.; Barrio, I.C.; Bowden, J.; Convey, P.; Coulson, S.J.; Culler, L.E.; Dahl, M.T.; Daly, K.M.; Koponen, S.; et al. Circumpolar terrestrial arthropod monitoring: A review of ongoing activities, opportunities and challenges, with a focus on spiders. *Ambio* **2020**, *49*, 704–717, doi:10.1007/s13280-019-01185-y.
2. Coulson, S.J. The terrestrial invertebrate fauna of the Svalbard archipelago in a changing world: History of research and challenges. *Can. Entomol.* **2013**, *145*, 131–146, doi:10.4039/ce.2012.110.
3. IPCC. Summary for Policymakers. In *Global Warming of 1.5 °C. An IPCC Special Report on the Impacts of Global Warming of 1.5°C above Pre-Industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*; Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P.R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., et al., Eds.; Intergovernmental Panel on Climate Change: Geneva, Switzerland, 2018; pp. 3–27.
4. Hanssen-Bauer, I.; Førland, E.J.; Hisdal, H.; Mayer, S.; Sandø, A.B.; Sorteberg, A. (Eds.) *Climate in Svalbard 2100—A Knowledge Base for Climate Adaptation*; The Norwegian Centre for Climate Services report 1/2019; The Norwegian Centre for Climate Services: Oslo, Norway, 2019; Available online: <https://www.miljodirektoratet.no/publikasjoner/2019> (accessed on 19 April 2020).
5. Norwegian Meteorological Institute. Precipitation in Svalbard, Hopen and Jan Mayen, annual total. Environmental Monitoring of Svalbard and Jan Mayen (MOSJ), 2020. Available online: <http://www.mosj.no/en/climate/atmosphere/temperature-precipitation.html> (accessed on 8 August 2020).

6. Descamps, S.; Aars, J.; Fuglei, E.; Kovacs, K.M.; Lydersen, C.; Pavlova, O.; Pedersen, Å.Ø.; Ravolainen, V.; Strøm, H. Climate change impacts on wildlife in a High Arctic. *Glob. Chang. Biol.* **2017**, *23*, 490–502, doi:10.1038/s41598-019-47951-6.
7. Summerhayes, V.S.; Elton, C.S. Further contribution to the ecology of Spitsbergen. *J. Ecol.* **1928**, *16*, 193–268.
8. Ávila-Jiménez, M.L.; Solhøy, T.; Gwiazdowicz, D.J.; Fjellberg, A.; Dozsa-Farkas, K.; Monson, F.; De Smet, W.H.; Stur, E.; Ekrem, T.; Coulson, S.J. The terrestrial invertebrate fauna of Edgeøya, Svalbard: Arctic landscape community composition reflects biogeography patterns. *Polar Biol.* **2019**, *42*, 837–850, doi:10.1007/s00300-019-02471-x.
9. Seniczak, A.; Seniczak, S. Morphological ontogeny of *Fuscozetes coulsoni* sp. nov. (Acari: Oribatida: Ceratozetidae) from Svalbard, Norway. *Syst. Appl. Acarol.* **2020**, *25*, 680–696, doi:10.11158/saa.25.4.8.
10. Gwiazdowicz, D.J.; Zawieja, B.; Olejniczak, I.; Skubala, P.; Gdula, A.K.; Coulson, S.J. Changing microarthropod communities in front of a receding glacier in the High Arctic. *Insects* **2020**, *11*, 226, doi:10.3390/insects11040226.
11. Coulson, S.J.; Hodkinson, I.D.; Block, W.; Webb, N.R.; Harrison, J.A. Survival of terrestrial soil-dwelling arthropods on and in seawater: Implications for trans-oceanic dispersal. *Funct. Ecol.* **2002**, *16*, 353–356, doi:10.1046/j.1365-2435.2002.00636.x.
12. Coulson, S.J.; Moe, B.; Monson, F.; Gabrielsen, G.W. The invertebrate fauna of High Arctic seabird nests: The microarthropod community inhabiting nests on Spitsbergen, Svalbard. *Polar Biol.* **2009**, *32*, 1041–1046, doi:10.1007/s00300-009-0603-8.
13. Lebedeva, N.V.; Krivolutsky, D.A. Birds spread soil microarthropods to Arctic Islands. *Dokl. Biol. Sci.* **2003**, *391*, 392–332.
14. Lebedeva, N.V.; Lebedev, V.D.; Melekhina, E.N. New data on the oribatid mite (Oribatei) fauna of Svalbard. *Dokl. Biol. Sci.* **2006**, *407*, 182–186.
15. Lebedeva, N.V.; Melekhina, E.N.; Gwiazdowicz, D.J. New data on soil mites in the nests of *Larus hyperboreus* in the Spitsbergen archipelago. *Vestn. Juzn. Naucn. Cent. Ran* **2012**, *8*, 70–75.
16. Lebedeva, N.V.; Lebedev, V.D. Transport of oribatid mites to the polar areas by birds. In *Integrative Acarology*; Bertrand, M., Kreiter, S., McCoy, K.D., Migeon, A., Navajas, M., Tixier, M.S., Vial, L., Eds.; European Association of Acarologists: Montpellier, France, 2008; pp. 359–367.
17. Coulson, S.J. Association of the soil mite *Diapterobates notatus* (Thorell, 1871) (Acari, Oribatidae) with *Cynomya mortuorum* (Linnaeus, 1761) (Calliphoridae, Calliphorinae): Implications for the dispersal of oribatid mites. *Int. J. Acarol.* **2009**, *35*, 175–177, doi:10.1080/01647950902934242.
18. Coulson, S.J.; Fjellberg, A.; Gwiazdowicz, D.J.; Lebedeva, N.V.; Melekhina, E.N.; Solhøy, T.; Erséus, C.; Maraldo, K.; Miko, L.; Schatz, H.; et al. Introduction of invertebrates into the High Arctic via imported soils: The case of Barentsburg in the Svalbard. *Biol. Invasions* **2013**, *15*, 1–5, doi:10.1007/s10530-012-0277-y.
19. Coulson, S.J.; Fjellberg, A.; Gwiazdowicz, D.J.; Lebedeva, N.V.; Melekhina, E.N.; Solhøy, T.; Erséus, C.; Maraldo, K.; Miko, L.; Schatz, H.; et al. The invertebrate fauna of anthropogenic soils in the High-Arctic settlement of Barentsburg, Svalbard. *Polar Res.* **2013**, *32*, 19273, doi:10.3402/polar.v32i0.19273.
20. Coulson, S.J.; Fjellberg, A.; Melekhina, E.N.; Taskaeva, A.A.; Lebedeva, N.V.; Belkina, O.; Seniczak, S.; Seniczak, A.; Gwiazdowicz, D.J. Microarthropod communities of industrially disturbed or imported soils in the High Arctic; the abandoned coal mining town of Pyramiden, Svalbard. *Biodivers. Conserv.* **2015**, *24*, 1671–1690, doi:10.1007/s10531-015-0885-9.
21. Hughes, K.A.; Convey, P. The protection of Antarctic terrestrial ecosystems from inter and intra-continental transfer of non-indigenous species by human activities: A review of current systems and practices. *Glob. Environ. Chang.* **2010**, *20*, 96–112, doi:10.1016/j.gloenvcha.2009.09.005.
22. Seniczak, S.; Seniczak, A.; Gwiazdowicz, D.J.; Coulson, S.J. Community structure of oribatid and gamasid mites (Acari) in moss-grass tundra in Svalbard (Spitsbergen, Norway). *Arct. Antarct. Alp. Res.* **2014**, *46*, 591–599, doi:10.1657/1938-4246-46.3.591.
23. Muñoz-Leal, S.; González-Acuña, D. The tick *Ixodes uriae* (Acari: Ixodidae): Hosts, geographical distribution, and vector roles. *Ticks Tick-borne Dis.* **2015**, *6*, 843–868, doi:10.1016/j.ttbdis.2015.07.014.
24. Krantz, G.W.; Walter, D.E. *A Manual of Acarology*, 3rd ed.; Texas Tech University Press: Lubbock, TX, USA, 2009; pp. 1–807.
25. Beaulieu, F.; Dowling, A.P.G.; Klompen, H.; de Moraes, G.J.; Walter, D.E. Superorder Parasitiformes Reuter, 1909. In *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic*

- Richness; Zhang, Z.-Q., Ed. Magnolia Press: Auckland, New Zealand, 2011; pp. 123–128, doi:10.11646/zootaxa.3148.1.23.
- 26. Fauna Europaea. Available online: <https://fauna-eu.org> (accessed on 19 April 2020).
 - 27. Gwiazdowicz, D.J.; Teodorowicz, E.; Coulson, S.J. Redescription of *Zercon solenites* Haarløv, 1942 (Acari, Zerconidae) with a key to the Svalbard species of the genus *Zercon*. *Int. J. Acarol.* **2011**, *37* (Suppl. 1), 135–148, doi:10.1080/01647954.2010.543699.
 - 28. Kolodochka, L.A.; Gwiazdowicz, D.J. A new species of predaceous mite of the genus *Neoseiulus* Hughes (Acari, Phytoseiidae), with redescriptions of *N. magnanalis* (Thor) and *N. ellesmerei* (Chant & Hansell), from Svalbard, High Arctic. *Zootaxa* **2014**, *3793*, 441–452.
 - 29. Gwiazdowicz, D.J.; Rakowski, R. Redescription of *Proctolaelaps parvanalis* (Thor, 1930) (Acari: Ascidae) from Spitsbergen. *Entomol. Fenn.* **2009**, *20*, 281–286, doi:10.33338/ef.84491.
 - 30. Ávila-Jiménez, M.L.; Gwiazdowicz, D.J.; Coulson, S.J. The mesostigmatid mite (Acari: Parasitiformes) fauna of Svalbard: A revised inventory of a high Arctic Archipelago. *Zootaxa* **2011**, *3091*, 33–41, doi:10.11646/zootaxa.3091.1.2.
 - 31. Zhang, Z.-Q.; Fan, Q.-H.; Pesic, V.; Smit, H.; Bochkov, A.V.; Khaustov, A.A.; Baker, A.; Wohltmann, A.; Wen, T.H.; Amrine, J.W.; et al. Order Trombidiformes Reuter, 1909. In *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness*; Zhang, Z.-Q., Ed.; Magnolia Press: Auckland, New Zealand, 2011; pp. 129–138, doi:10.11646/zootaxa.3148.1.24.
 - 32. Bolton, S.J.; Chetverikov, P.E.; Klompen, H. Morphological support for a clade comprising two vermiform mite lineages: Eriophyoidea (Acariformes) and Nematalycidae (Acariformes). *Syst. Appl. Acarol.* **2017**, *22*, 1096–1131, doi:10.11158/saa.22.8.2.
 - 33. Klimov, P.B.; OConnor, B.M.; Chetverikov, P.E.; Bolton, S.J.; Pepato, A.R.; Mortazavi, A.L.; Tolstikov, A.V.; Bauchan, G.R.; Ochoa, R. Comprehensive phylogeny of acariform mites (Acariformes) provides insights on the origin of the four-legged mites (Eriophyoidea), a long branch. *Mol. Phylogenet. Evol.* **2018**, *119*, 105–117, doi:10.1016/j.ympev.2017.10.017.
 - 34. Hernandes, F.A.; Skvarla, M.J.; Fisher, J.R.; Dowling, A.P.G.; Ochoa, R.; Ueckermann, E.A.; Bauchan, G.R. Catalogue of snout mites (Acariformes: Bdellidae) of the world. *Zootaxa* **2016**, *4152*, 1–83, doi:10.1111/j.1439-0418.1971.tb03191.x.
 - 35. Skvarla, M.J.; Fisher, J.R.; Dowling, A.P. A review of Cunaxidae (Acariformes, Trombidiformes): Histories and diagnoses of subfamilies and genera, keys to world species, and some new locality records. *ZooKeys* **2014**, *418*, 1–103, doi:10.3897/zookeys.418.7629.
 - 36. Khaustov, A.A. A new genus and species in the mite family Eupodidae (Acari, Eupodoidea) from Crimea. *ZooKeys* **2014**, *422*, 11–22, doi:10.3897/zookeys.422.7802.
 - 37. Szudarek-Trepto, N.; Kazmierski, A.; Dabert, M.; Dabert, J. Molecular phylogeny of Eupodidae reveals that the family Cocceupodidae (Actinotrichida; Eupodoidea) and its genus *Filieupodes* are valid taxa. *Exp. Appl. Acarol.* **2020**, *80*, 43–57, doi:10.1007/s10493-019-00455-4.
 - 38. Meyer, M.K.P.; Ryke, P.A.J. Mites of the superfamily Eupodoidea (Acarina: Prostigmata) associated with South African plants. *S. Afr. J. Agric. Sci.* **1960**, *3*, 481–496.
 - 39. Strandtmann, R.W.; Davies, L. Eupodiform mites from Possession Island, Crozet Islands, with a key to the species of *Eupodes* (Acarina: Prostigmata). *Pac. Insects* **1972**, *14*, 39–56.
 - 40. Umina, P.A.; Hoffman, A.A.; Weeks, A.R. Biology, ecology and control of the *Penthaleus* species complex (Acari: Penthaleidae). *Exp. Appl. Acarol.* **2004**, *34*, 211–237, doi:10.1007/s10493-004-1804-z.
 - 41. Jasionowska, K. A morphological study of the genus *Penthalodes* (Acari, Prostigmata, Eupodida, Pentalodidae) with description of a new species. *Zootaxa* **2010**, *2672*, 29–49.
 - 42. Zacharda, M. Soil mites of the family Rhagidiidae (Actinedida: Eupodoidea). Morphology, systematics, ecology. *Acta Univ. Carol. Biol.* **1980**, *516*, 489–785.
 - 43. Zacharda, M.; Gude, M.; Kraus, S.; Hauck, C.; Molenda, R.; Růžička, V. The relict mite *Rhagidia gelida* (Acari, Rhagidiidae) as a biological cryoindicator of periglacial microclimate in European Highland Scree. *Arct. Antarct. Alp. Res.* **2005**, *37*, 402–408, doi:10.1657/1523-0430(2005)037[0402:TRMRGA]2.0.CO;2.
 - 44. Qin, T.K. A checklist and key to species of Eupodoidea (Acari: Prostigmata) from Australia and New Zealand and their subantarctic islands. *J. R. Soc. N. Z.* **1998**, *28*, 295–307.
 - 45. Bartsch, I. Checklist of marine and freshwater halacarid mite genera and species (Halacaridae: Acari) with notes on synonyms, habitats, distribution. *Zootaxa* **2009**, *1998*, 1–170, doi:10.11646/zootaxa.1998.1.1.

46. André, H.M. Redefinition of the genus *Triophtydeus* Thor, 1932 (Acari: Actinedida). *Zool. Meded.* **1985**, *59*, 189–195.
47. Thor, S. *Acarina-Tydeidae, Ereynetidae*; Das Tierreich, Walter de Gruyter: Berlin, Germany, 1933; Volume 60, pp. 1–84.
48. Makarova, O.; Behan-Pelletier, V. 18.3.6. Oribatida (=Cryptostigmata, Beetle Mites). In *The Greenland Entomofauna*; Böcher, J., Kristensen, N.P., Pape, T., Vilhelmsen, L., Eds.; An Identification Manual of Insects, Spiders and Their Allies. Koninklijke Brill NV: Leiden, The Netherlands, 2015; pp. 802–845.
49. André, H.M. In search of the true *Tydeus* (Acari, Tydeidae). *J. Nat. Hist.* **2005**, *39*, 975–1001, doi:10.1080/00222930400002838.
50. Silva, G.L.; Metzelthin, M.H.; Silva, O.S.; Ferla, N.J. Catalogue of the mite family Tydeidae (Acari: Prostigmata) with the world key to the species. *Zootaxa* **2016**, *4135*, 1–68, doi:10.11646/zootaxa.4135.1.1.
51. Baker, E.W. A review of the genera of the family Tydeidae (Acarina). *Adv. Acarol.* **1965**, *2*, 95–133.
52. André, H.M. A generic revision of the family Tydeidae (Acari: Actinedida). IV. Generic descriptions, keys and conclusions. *Bull. Ann. Soc. r. Belg. Ent.* **1980**, *116*, 103–130.
53. Makarova, O.I. Acarocenoses (Acariformes, Parasitiformes) in polar deserts: 1. Mite assemblages of the Severnaya Zemlya Archipelago: Structure of fauna and abundance. *Entomol. Rev.* **2002**, *82*, 839–856.
54. Abe, H. Annotated checklist of Japanese water mites (Acari: Prostigmata: Hydracarina). *Acta Arachnol.* **2005**, *54*, 111–145, doi:10.2476/asjaa.54.111.
55. Esen, Y.; Pesic, V.; Erman, O. Water mites of the genus *Sperchon* Kramer (Acari: Hydrachnidia: Sperchontidae) from Turkey, with description of two new species. *Zootaxa* **2010**, *2514*, 35–46, doi:10.1080/01650424.2012.718080.
56. Małkol, J.; Wohltmann, A. An annotated checklist of terrestrial Parasitengona (Actinotrichida: Prostigmata) of the world, excluding Trombiculidae and Walchiidae. *Ann. Zool.* **2012**, *62*, 359–562, doi:10.3161/000345412X656671.
57. Skoracki, M.; Zawierucha, K. *Chenophila nanseni* sp. n. (Acari: Syringophilidae) parasitising the barnacle goose in Svalbard. *Pol. Polar Res.* **2016**, *37*, 121–130, doi:10.1515/popore-2016-0002.
58. Beron, P. Acarorum Catalogus VII. Trombidiformes: Prostigmata: Raphignathoidea. Fam. Barbutiidae, Caligonellidae, Camerobiidae, Cryptognathidae, Dasythyreidae, Dytiscacaridae, Eupalopsellidae, Homocaligidae, Mecognathidae, Raphignathidae, Stigmaeidae, Xenocaligonellididae. *Penssoft Natl. Mus. Nat. Hist. Sofia Bulg.* **2020**, doi:10.3897/ab.e55087.
59. Pritchard, A.E.; Baker, E.W. *A Revision of the Spider Mite Family Tetranychidae*; Memoirs Series; Volume 2; Pacific Coast Entomological Society: San Francisco, CA, USA, 1955.
60. Migeon, A.; Dorkeld, F. Spider Mites Web: A Comprehensive Database for the Tetranychidae, 2020. Available online: <http://www1.montpellier.inra.fr/CBGP/spmweb> (accessed on 2 July 2020).
61. Khaustov, A.A. A description of new genus, *Kerdabania* gen. n. with four new species (Acari: Heterostigmata: Neopygmephoridae). *Acarina* **2009**, *17*, 171–188.
62. Schatz, H.; Behan-Pelletier, V.M.; O'Connor, B.M.; Norton, R.A. Suborder Oribatida van der Hammen, 1968. In *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness*; Zhang, Z.-Q., Ed.; Magnolia Press: Auckland, New Zealand, 2011; pp. 141–148.
63. Beaulieu, F.; Knee, W.; Nowell, V.; Schwarzbeld, M.; Lindo, Z.; Behan-Pelletier, V.M.; Lumley, L.; Young, M.R.; Smith, I.; Proctor, H.C.; Mironov, S.V.; et al. Acari of Canada. In *The Biota of Canada—A Biodiversity Assessment. Part 1: The Terrestrial Arthropods*; Langor, D.W., Sheffield, C.S., Eds.; ZooKeys **2019**, *819*, 77–168, doi:10.3897/zookeys.819.28307.
64. Walter, D.E.; Bolton, S.; Uusitalo, M.; Zhang, Z.-Q. Suborder Endeostigmata Reuter, 1909. In *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness*; Zhang, Z.-Q., Ed.; Magnolia Press: Auckland, New Zealand, 2011; pp. 139–140.
65. Uusitalo, M. Terrestrial species of the genus *Nanorchestes* (Endeostigmata: Nanorchestidae) in Europe. In *Trends in Acarology*; Sabelis, M.W., Bruin, J., Eds.; Springer: Dordrecht, The Netherlands, 2010; pp. 161–166, doi:10.1007/978-90-481-9837-5_25.
66. Thor, S.; Willmann, C. Acarina. 7la Eupodidae, Penthalodidae, Penthaleidae, Rhagidiidae, Pachygnathidae, Cunaxidae. *Das Tierreich* **1941**, *71*, 1–186.
67. Kiedrowicz, A.; Rector, B.G.; Zawierucha, K.; Szydło, W.; Skoracka, A. Phytophagous mites (Acari: Eriophyoidea) recorded from Svalbard, including the description of a new species. *Polar Biol.* **2016**, *39*, 1359–1368, doi:10.1007/s10531-015-0885-9.

68. Subías, L.S. Listado Sistemático, Sinonímico y Biogeográfico de los Ácaros Oribátidos (Acariformes, Oribatida) del Mundo (excepto fósiles) (15^a actualización). *Graellsia* **2004**, *2020*, 60 (número extraordinario), 3–305 (2004), actualized pdf in 2020. Available online: http://bba.bioucm.es/cont/docs/RO_1.pdf (accessed on 14 April 2020).
69. Seniczak, S.; Seniczak, A. Differentiation of external morphology of Oribatulidae (Acari: Oribatida) in light of the ontogeny of three species. *Zootaxa* **2012**, *3184*, 1–34, doi:10.11646/zootaxa.3184.1.1.
70. Seniczak, A.; Seniczak, S. Morphological ontogeny of *Achipteria punctata* (Acari: Oribatida: Achipteriidae). *Zootaxa* **2018**, *4540*, 54–72, doi:10.11646/zootaxa.4540.1.7.
71. Bayartogtokh, B.; Schatz, H.; Ekrem, T. Distribution and diversity of the soil mites of Svalbard, with redescriptions of three known species (Acari: Oribatida). *Int. J. Acarol.* **2011**, *37*, 467–484, doi:10.1080/01647954.2010.525525.
72. Miko, L.; Monson, F.D. Two interesting damaeid mites (Acari, Oribatida, Damaeidae Berlese, 1896) from the British Isles and Svalbard (Spitsbergen, Norway), with a description of *Kunstidamaeus arcticus* n.sp. *Acarologia* **2013**, *53*, 89–100, doi:10.1051/acarologia/20132077.
73. Klimov, P.B. A review of acarid mites of the tribe Caloglyphini (Acaridae, Acariformes) with description of a new genus and species from Siberia and Russian Far East. *Vestn. Zool.* **2000**, *34*, 27–35.
74. Dabert, M.; Coulson, S.J.; Gwiazdowicz, D.J.; Moe, B.; Hanssen, S.A.; Biersma, E.; Pilskog, H.E.; Dabert, J. Differences in speciation progress in feather mites (Analgoidea) inhabiting the same host: The case of *Zachvatkinia* and *Alloptes* living on arctic and long-tailed skuas. *Exp. Appl. Acarol.* **2015**, *65*, 163–179, doi:10.1007/s10493-014-9856-1.
75. Dabert, J. Personal Communication; Adam Mickiewicz University: Poznan, Poland, 2020.
76. McCoy, K.D. Consequences of Dispersal in Host–Parasite Systems: Population Dynamics, Genetic Structure, and Localadaptation of the Seabird tickixodes Uriae. Ph.D. Thesis, Pierreand Marie Curie University, Paris, France, 2001.
77. Coulson, S.J.; Lorentzen, E.; Strøm, H.; Gabrielsen, G.W. The parasitic tick *Ixodes uriae* (Acari: Ixodidae) on seabirds from Spitsbergen, Svalbard. *Polar Res.* **2009**, *28*, 399–402, doi:10.3402/polar.v28i3.6137.
78. Descamps, S. Winter temperature affects the prevalence of ticks in an Arctic sea-bird. *PLoS ONE* **2013**, *8*, e65374, doi:10.1371/journal.pone.0065374.
79. Teodorowicz, E.; Gwiazdowicz, D.J.; Coulson, S.J. Redescription of *Antennoseius (Vitzthumia) oudemansi* (Acari, Mesostigmata) from Spitsbergen, Svalbard. *Entomol. Fenn.* **2014**, *25*, 27–42, doi:10.33338/ef.41467.
80. Gwiazdowicz, D.J.; Teodorowicz, E.; Coulson, S.J. Redescription of *Arctoseius haarlovi* Lindquist, 1963 (Acari: Ascidae) from Spitsbergen. *Entomol. Fenn.* **2011**, *22*, 140–148, doi:10.33338/ef.4692.
81. Trouessart, E. Note sur les Acariens recueillis au Spitzberg pendant le voyage de la Manche. *Nouv. Arch. Miss. Sci. Litt.* **1893**, *5*, 255–263.
82. Trouessart, E. Révision des Acariens des régions Arctiques et description d'espèces nouvelles. *M. Soc. Natl. Sci. Nat. Math. Cherb.* **1895**, *29*, 183–206.
83. Zmudczyńska-Skarbek, K.; Barcikowski, M.; Drobniak, S.M.; Gwiazdowicz, D.J.; Richard, P.; Skubała, P.; Stempniewicz, L. Transfer of ornithogenic influence through different trophic levels of the Arctic terrestrial ecosystem of Bjørnøya (Bear Island), Svalbard. *Soil Biol. Biochem.* **2017**, *115*, 475–489, doi:10.1016/j.soilbio.2017.09.008.
84. Makarova, O.L. Gamasid mites (Parasitiformes, Mesostigmata) of the European Arctic and their distribution patterns. *Entomol. Rev.* **2013**, *93*, 113–133, doi:10.1134/S0013873813010156.
85. Gwiazdowicz, D.J.; Gulvik, M.E. Mesostigmatid mites (Acari, Mesostigmata) in Svalbard. In *Materiały XXXII Międzynarodowego Sympozjum Polarnego*; Uniwersytet Wrocławski, Wydział Nauk o Ziemi i Kształtowania Środowiska: Wrocław, Poland, 2008; pp. 32–34.
86. Coulson, S.J.; Schatz, H.; Gwiazdowicz, D.J.; Solhøy, T. On the oribatid and mesostigmatid mites (Acari) of the High Arctic island of Hopen. *Pol. Polar Res.* **2014**, *35*, 133–139, doi:10.2478/popore-2014-0002.
87. Coulson, S.J.; Fjellberg, A.; Snazell, R.; Gwiazdowicz, D.J.; Ávila-Jiménez, M.L. On the Collembola, Araneae and Gamasida from the Kinnvika region of Nordaustlandet, Svalbard. *Geogr. Ann. A* **2011**, *93*, 253–257, doi:10.2307/41432731.
88. Byzova, J.B.; Uvarov, A.V.; Petrova, A.D. Seasonal changes in communities of soil invertebrates in tundra ecosystems of Hornsund, Spitsbergen. *Pol. Polar Res.* **1995**, *16*, 245–266.
89. Gwiazdowicz, D.J.; Coulson, S.J. High-Arctic gamasid mites (Acari, Mesostigmata): Community composition on Spitsbergen, Svalbard. *Polar Res.* **2011**, *30*, 8311, doi:10.3402/polar.v30i0.8311.

90. Pilskog, H.E.; Solhøy, T.; Gwiazdowicz, D.J.; Grytnes, J.A.; Coulson, S.J. Invertebrate communities inhabiting nests of migrating passerine, wild fowl and sea birds breeding in the High Arctic, Svalbard. *Polar Biol.* **2014**, *37*, 981–998.
91. Thor, S. Beiträge zur Kenntnis der Invertebratenfauna von Svalbard. *Skr. Svalbard Ishavet* **1930**, *27*, 1–156.
92. Gwiazdowicz, D.J.; Solhøy, T.; Coulson, S.J.; Lebedeva, N.V.; Melekhina, E.N. First record of *Vulgarogamasus immanis* (Acari, Mesostigmata) in Svalbard. *Pol. Polar Res.* **2012**, *33*, 35–39.
93. Gwiazdowicz, D.J.; Teodorowicz, E. Description of *Halolaelaps coulsoni* n. sp. (Acari, Halolaelapidae) from the High Arctic. *Acarologia* **2017**, *57*, 393–406, doi:10.1051/acarologia/20174164.
94. Hirschmann, W. Die Gattung *Halolaelaps* Berlese & Trouessart 1889 nov. comb. *Acarologia* **1966**, *9*, 21–24.
95. Gwiazdowicz, D.J.; Coulson, S.J. First record of *Thinoseius spinosus* (Acari, Eviphididae) from the High Arctic Island of Spitsbergen (Svalbard) including a key to deutonymphs of genus *Thinoseius*. *Int. J. Acarol.* **2010**, *36*, 233–236, doi:10.1080/0164795100359858.
96. Makarova, O.L. To the study of mites of the genus *Arctoseius* Thor (Parasitiformes, Ascidae) from the Far North: 3. Ranges and ecological preferences of species. *Entomol. Rev.* **2000**, *80*, 143–150.
97. Lindquist, L.L.; Makarova, O.L. Two new circumpolar mite species of the genus *Arctoseius* (Parasitiformes, Mesostigmata, Ascidae). *Entomol. Rev.* **2011**, *91*, 1054–1072.
98. Gwiazdowicz, D.J.; Coulson, S.J.; Grytnes, J.A.; Pilskog, H.E. The bird ectoparasite *Dermanyssus hirundinis* (Acari, Mesostigmata) in the High Arctic; a new parasitic mite to Spitsbergen, Svalbard. *Acta Parasitol.* **2012**, *57*, 378–384, doi:10.2478/s11686-012-0050-5.
99. Summerhayes, V.S.; Elton, C.S. Contributions to the ecology of Spitsbergen and Bear Island. *J. Ecol.* **1923**, *11*, 214–286.
100. Krumpál, M.; Cyprich, D.; Zejda, J.; Ambros, M. The occurrence of field vole (*Microtus arvalis* Pallas 1778) and its acarofauna on Spitsbergen (Svalbard). *Biologia* **1991**, *46*, 881–885.
101. Hull, M.A. On some land mites (Acari) from Spitzbergen and Bjørnøya. Results of the Oxford University Expedition to Spitsbergen, 1921. *Ann. Mag. Nat. Hist.* **1922**, *10*, 621–623.
102. Thorell, T. Om Arachnider från Spetsbergen och Beeren-Eiland. *Öfvers. Kongl. Vetensk.-Akad. Förh.* **1871**, *6*, 683–702.
103. Thor, S. Neue Beiträge zur Kenntnis der Invertebraten-Fauna von Svalbard. *Zool. Anz.* **1934**, *107*, 114–139.
104. Bertram, G.C.L.; Lack, D. Notes on the animal ecology of Bear Island. *J. Anim. Ecol.* **1938**, *7*, 27–52.
105. Trägårdh, I. Beiträge zur Fauna der Bären-Insel. *Bihang till K. Svenska Vet. Akad. Handlingar.* **1900**, *26*, 1–24.
106. Haarløv, N. A morphologic–systematic–ecological investigation of Acarina and other representatives of the microfauna of the soil around Mørkefjord, Northeast Greenland. *Medd. Grønl.* **1942**, *128*, 1–71.
107. Trouessart, E. Note préliminaire sur les acariens marins (Halacaridae) recueillis par S.A.S. le Prince de Monaco, dans les mers arctiques. *Bull. Soc. Zool. Fr.* **1902**, *27*, 66–70.
108. Barr, W. The Helgoland Expedition to Svalbard: Die Deutsche Expedition in das Nordliche Eismeer, 1898. *Arctic* **1988**, *41*, 203–214.
109. Chertoprud, E.S.; Makarova, O.L.; Novichkova, A.A. First data on aquatic mites (Acari) of inland water bodies of West Spitsbergen, Svalbard. *Acarina* **2017**, *25*, 181–189, doi:10.21684/0132-8077-2017-25-2-181-189.
110. Thor, S. Norwegische Tydeidae VIII–XV, bemerkungen über die Gattung *Tydeus* und über Augen, Tracheen usw. *Zool. Anz.* **1932**, *98*, 69–91.
111. Soar, C.D. A species of Hydracarina found at Bear Island, June 17th, 1921—The Oxford University Expedition to Spitsbergen, 1921. *Microscopy* **1922**, *14*, 301–304.
112. Oudemans, A.C. Acarologische Aanteekeningen. XCIX. *Entomol. Ber.* **1930**, *8*, 11–20.
113. Thor, S. Norwegische Alycidae I–VII. *Zool. Anz.* **1931**, *194*, 229–238.
114. Niedbała, W. Oribatei (Acari) of Spitsbergen. *B. Pol. Acad. Sci.-Biol.* **1971**, *19*, 737–742.
115. Karppinen, E. Notes on the arthropod fauna of Spitsbergen. *Acta Entomol. Fenn.* **1967**, *33*, 18–26.
116. Forsslund, K.H. Notizen über Oribatei (Acari). I. *Ark. Zool.* **1957**, *2*, 583–593.
117. Forsslund, K.H. *Liochthonius muscorum* n. sp. und *L. lapponicus* (Träg.) (Acari, Oribatei). *Entomol. Tidskr.* **1964**, *85*, 236–239.
118. Byzova, J.B.; Uvarov, A.V. Mites of the tundra biotopes in the vicinity of bird colonies on Spitsbergen. In *The 6th all-Union Meeting on the Problems of Theoretical and Applied Acarology. Aszchabad, Turkmenistan. Abstracts*; Balashov, Y.S., Ed.; Russian Academy of Science: Saint Petersburg, Russia, 1990; pp. 152–153.
119. Coulson, S.J. Terrestrial and freshwater invertebrate fauna of the high Arctic Archipelago of Svalbard. *Zootaxa* **2007**, *1448*, 41–58, doi:10.4039/tce.2012.110.

120. Coulson, S. *Personal Communication*; Swedish University of Agricultural Sciences: Uppsala, Sweden, 2020.
121. Coulson, S.J.; Hodkinson, I.D.; Strathdee, A.T.; Bale, J.S.; Block, W.; Worland, M.R.; Webb, N.R. Simulated climate change: The interaction between vegetation type and microhabitat temperatures at Ny Ålesund, Svalbard. *Polar Biol.* **1993**, *13*, 67–70, doi:10.1007/BF00236585.
122. Coulson, S.J.; Hodkinson, I.D.; Block, W.; Webb, N.R.; Worland, M.R. Low summer temperatures: A potential mortality factor for high arctic soil microarthropods? *J. Insect Physiol.* **1995**, *41*, 783–792, doi:10.1016/0022-1910(95)00027-R.
123. Coulson, S.J.; Hodkinson, I.D.; Webb, N.R.; Block, W.; Bale, J.S.; Strathdee, A.T.; Worland, M.R.; Wooley, C. Effects of experimental temperature elevation on high-arctic soil microarthropod populations. *Polar Biol.* **1996**, *16*, 147–153, doi:10.1007/BF02390435.
124. Coulson, S.J.; Leinaas, H.P.; Ims, R.A.; Søvik, G. Experimental manipulation of the winter surface ice layer—The effects on a High Arctic soil microarthropod community. *Ecography* **2000**, *23*, 299–306, doi:10.1111/j.1600-0587.2000.tb00285.x.
125. Coulson, S.J.; Hodkinson, I.D.; Webb, N.R. Microscale distribution patterns in high Arctic soil microarthropod communities: The influence of plant species within the vegetation mosaic. *Ecography* **2003**, *26*, 801–809, doi:10.1111/j.0906-7590.2003.03646.x.
126. Hodkinson, I.D.; Coulson, S.J.; Webb, N.R.; Block, W.; Strathdee, A.T.; Bale, J.S. Feeding studies on *Onychiurus arcticus* (Tullberg) (Collembola: Onychiuridae) on West Spitsbergen. *Polar Biol.* **1993**, *14*, 17–19.
127. Webb, N.R.; Coulson, S.J.; Hodkinson, I.D.; Block, W.; Bale, J.S.; Strathdee, A.T. The effects of experimental temperature elevation on populations of cryptostigmatic mites in high Arctic soils. *Pedobiologia* **1998**, *42*, 298–308.
128. Hodkinson, I.D.; Bird, J.M. Anoxia tolerance in high Arctic terrestrial microarthropods. *Ecol. Entomol.* **2004**, *29*, 506–509, doi:10.1111/j.0307-6946.2004.00619.x.
129. Dollery, R.; Hodkinson, I.D.; Jónsdóttir, I.S. Impact of warming and timing of snow melt on soil microarthropod assemblages associated with *Dryas*-dominated plant communities in Svalbard. *Ecography* **2006**, *29*, 111–119, doi:10.1111/j.2006.0906-7590.04366.x.
130. Kulczyński, V. Zoologische Ergebnisse der russischen Expeditionen nach Spitzbergen. *Annu. Mus. Zool. Acad. Imp. Sci. St. Petersb.* **1902**, *7*, 347–354.
131. Solhøy, I.W. A redescription of *Mycobates sarekensis* (Trägårdh) (Acari: Oribatei). *Acarologia* **1997**, *38*, 69–77.
132. Block, W. Some arctic Oribatei. *Acarologia* **1966**, *8*, 161–162.
133. Trägårdh, I. Monographie der arktischen Acariden. *Fauna Arct.* **1904**, *4*, 1–78.
134. Seniczak, S.; Seniczak, A.; Coulson, S.J. Morphological ontogeny, distribution and descriptive population parameters of *Hermannia reticulata* (Acari: Oribatida: Hermanniidae), with comments on Crotonioidea. *Int. J. Acarol.* **2017**, *43*, 52–72, doi:10.1080/01647954.2016.1229812.
135. Seniczak, S.; Seniczak, A.; Coulson, S.J. Morphological ontogeny and distribution of *Hermannia scabra* (Acari: Oribatida: Hermanniidae) in Svalbard and descriptive population parameters. *Acarologia* **2017**, *57*, 877–892, doi:10.24349/acarologia/20174214.
136. Seniczak, S.; Plichta, W. Structural dependence of moss mite populations on patchiness of vegetation in moss-lichen-tundra at the north coast of Hornsund, West Spitsbergen. *Pedobiologia* **1978**, *18*, 145–152.
137. Gordeeva, E.V. *Moritzoppiella* gen. n.—A new genus of oribatid mites of the family Oppiidae (Oribatei, Acariformes). *Vestn. Zool.* **2000**, *34*, 17–26.
138. Hayward, S.A.L.; Worland, M.R.; Bale, J.S.; Convey, P. Temperature and the hygro preference of the Arctic collembolan *Onychiurus arcticus* and mite *Lauroppia translamellata*. *Physiol. Entomol.* **2000**, *25*, 266–272, doi:10.1046/j.1365-3032.2000.00191.x.
139. Thor, S. Übersicht der norwegischen Cryptostigmata mit einzelnen Neben bemerkungen. *NYT Mag. Nat.* **1937**, *77*, 275–307.
140. Schubart, H. Morphologische Grundlagen für die Klärung der Verwandtschaftsbeziehungen innerhalb der Milbenfamilie Ameronothridae (Acari, Oribatei). *Zoologica* **1975**, *123*, 23–91.
141. Søvik, G.; Leinaas, H.P. Variation in extraction efficiency between juvenile and adult oribatid mites: *Ameronothrus lineatus* (Oribatida, Acari) in a Macfadyen high-gradient canister extractor. *Pedobiologia* **2002**, *46*, 34–41, doi:10.1078/0031-4056-00111.
142. Søvik, G.; Leinaas, H.P. Adult survival and reproduction in an arctic mite, *Ameronothrus lineatus* (Acari, Oribatida): Effects of temperature and winter cold. *Can. J. Zool.* **2003**, *81*, 1579–1588, doi:10.1139/z03-113.

143. Søvik, G.; Leinaas, H.P. Long life cycle and high adult survival in an arctic population of the mite *Ameronothrus lineatus* (Acari, Oribatida) from Svalbard. *Polar Biol.* **2003**, *26*, 500–508, doi:10.1007/s00300-003-0510-3.
144. Søvik, G. Observations on ovoviparity and mixed-parity mode in arctic populations of *Ameronothrus lineatus* (Acari, Oribatida). *Acarologia* **2003**, *43*, 393–398.
145. Søvik, G.; Leinaas, H.P.; Ims, R.A.; Solhøy, T. Population dynamics and life history of the oribatid mite *Ameronothrus lineatus* (Acari, Oribatida) on the high Arctic Archipelago of Svalbard. *Pedobiologia* **2003**, *47*, 257–271, doi:10.1078/0031-4056-00189.
146. Seniczak, S.; Seniczak, A.; Graczyk, R.; Tømmervik, H.; Coulson, S.J. Distribution and population characteristics of the soil mites *Diapterobates notatus* and *Svalbardia paludicola* (Acari: Oribatida: Ceratozetidae) in High Arctic Svalbard (Norway). *Polar Biol.* **2017**, *40*, 1545–1555, doi:10.1007/s00300-017-2076-5.
147. Seniczak, S.; Seniczak, A.; Coulson, S.J. Morphology, distribution and certain population parameters of the Arctic mite *Oribatella arctica* (Acari: Oribatida: Oribatellidae). *Int. J. Acarol.* **2015**, *41*, 395–414.
148. Behan-Pelletier, V.M. Ceratozetidae of the western North American Arctic. *Can. Entomol.* **1985**, *117*, 1287–1366, doi:10.4039/Ent1171287-11.
149. Seniczak, S.; Seniczak, A.; Coulson, S.J. Morphology, distribution and biology of *Mycobates sarekensis* (Acari: Oribatida: Punctoribatidae). *Int. J. Acarol.* **2015**, *41*, 663–675, doi:10.1080/01647954.2015.1102968.
150. Fain, A.; Camerik, A.M. Notes on the mites of the genus *Ereynetes* Berlese (Acari: Ereynetinae), with description of five new species from South Africa. *Bull. Inst. R. Sci. Nat. Belg. Entomol.* **1994**, *64*, 145–164.
151. Makarova, O.L. The fauna of free-living mites (Acari) of Greenland. *Entomol. Rev.* **2015**, *95*, 108–125, doi:10.1134/S0013873815010133.
152. Gjelstrup, P.; Solhøy, T. Oribatid mites (Acari). In *The Zoology of Iceland, Steenstrupia 3*; Zoological Museum University of Copenhagen: Copenhagen, Denmark, 1994; Volume 57e, pp. 1–78.
153. Alfredsson, M.; Olafsson, E.; Eydal, M.; Unnsteinsdottir, E.R.; Hansford, K.; Wint, W.; Alexander, N.; Medlock, J.M. Surveillance of *Ixodes ricinus* ticks (Acari: Ixodidae) in Iceland. *Parasit. Vectors* **2017**, *10*, 466, doi:10.1186/s13071-017-2375-2.
154. Mehl, R. Checklist of Norwegian ticks and mites (Acari). *Fauna Norv. Ser. B* **1979**, *26*, 31–45.
155. Gwiazdowicz, D.J.; Gulvik, M.E. Checklist of Norwegian mesostigmatid mites (Acari, Mesostigmata). *Nor. J. Entomol.* **2005**, *52*, 117–125.
156. Seniczak, A.; Bolger, T.; Roth, S.; Seniczak, S.; Djursvoll, P.; Jordal, B.H. Diverse mite communities (Acari: Oribatida, Mesostigmata) from a broadleaf forest in western Norway. *Ann. Zool. Fenn.* **2019**, *56*, 121–136, doi:10.5735/086.056.0111.
157. Seniczak, A.; Seniczak, S.; Iturronobeitia, J.C.; Solhøy, T.; Flatberg, K.I. Diverse *Sphagnum* mosses support rich moss mite communities (Acari, Oribatida) in mires of western Norway. *Wetlands* **2019**, doi:10.1007/s13157-019-01236-w.
158. The Editors of Encyclopaedia Britannica, Svalbard, Encyclopædia Britannica, inc. Available online: <https://www.britannica.com/place/Svalbard> (accessed on 19 April 2020).
159. Veron, S.; Haevermans, T.; Govaerts, R.; Mouchet, M.; Pellens, R. Distribution and relative age of endemism across islands worldwide. *Sci. Rep.* **2019**, *9*, 11693, doi:10.1038/s41598-019-47951-6.
160. Losos, J.B.; Ricklefs, R.E. Adaptation and diversification on islands. *Nature* **2009**, *457*, 830–836, doi:10.1038/nature07893.
161. Melekhina, E.N. Taxonomic diversity and areology of oribatid mites (Oribatei) of the European North of Russia. *Proc. Komi Sci. Cent. Ural BranchRas* **2011**, *2*, 30–37.
162. Makarova, O.L.; Böcher, J. Diversity and geographical ranges of Greenland mites (Acari: Oribatida and Mesostigmata). In *Species and Communities in Extreme Environments*; Pensoft Publisher and KMK Science Press: Sofia, Moscow, Russia, 2009; pp. 165–186.
163. Dabert, M.; Witaliński, W.; Kaźmierski, A.; Olszanowski, Z.; Dabert, J. Molecular phylogeny of acariform mites (Acari, Arachnida): Strong conflict between phylogenetic signal and long-branch attraction artifacts. *Mol. Phylogenet. Evol.* **2010**, *56*, 222–241, doi:10.1016/j.ympev.2009.12.020.
164. Gerecke, R. Personal Communication; University of Tübingen: Tübingen, Germany, 2019.
165. Mehl, R.; Traavik, T. The tick *Ixodes uriae* (Acari, Ixodidae) in seabird colonies in Norway. *Fauna Norv. Ser. B* **1983**, *30*, 94–107.

166. Gwiazdowicz, D.J.; Coulson, S.J.; Ávila-Jiménez, M.L. First records of *Zercon andrei* Sellnick, 1958 and *Zerconopsis muestairi* Schweizer, 1949 (Acarina, Mesostigmata) from Bjørnøya, Svalbard. *Nor. J. Entomol.* **2009**, *56*, 117–119.
167. Seniczak, A.; Seniczak, S. Diversity of oribatid mites (Sarcoptiformes) in the Svalbard Archipelago: A historical overview. *Zootaxa* **2020**, *4834*, 41–65.
168. Behan-Pelletier, V.M. Oribatid mites (Acari: Oribatida) of the Yukon. In *Insects of the Yukon. Biological survey of Canada (Terrestrial Arthropods)*; Danks, H.V., Downes, J.A., Eds.; Canadian Museum of Nature: Ottawa, Canada, 1997; pp. 115–149.
169. Behan-Pelletier, V.; Schatz, H. Patterns of diversity in the Ceratozetoidea (Acari: Oribatida): A North American assessment. In *Trends in Acarology*; Sabelis, M.W., Bruun, J., Eds.; Springer: Dordrecht, The Netherlands, 2010; pp. 97–104, doi:10.1007/978-90-481-9837-5_15.
170. Ryabinin, N.A. Oribatid mites (Acari, Oribatida) in soils of the Russian Far East. *Zootaxa* **2015**, *3914*, 201–244.
171. Bayartogtokh, B. Fauna and communities of oribatid mites of Mongolia (Acari: Oribatida). Ph.D. Thesis, Russian Academy of Sciences, Moscow, Russia, 2007.
172. Fischer, B.M.; Schatz, H.; Querner, P.; Pauli, H. *Ceratozetes spitsbergensis* Thor, 1934: An Arctic mite new to continental Europe (Acari: Oribatida). *Int. J. Acarol.* **2016**, *42*, 135–139, doi:10.1080/01647954.2015.1133702.
173. Landvik, J.Y.; Brook, E.J.; Gualtieri, L.; Raisbeck, G.; Salvigsen, O.; Yiou, F. Northwest Svalbard during the last glaciation: Ice-free areas existed. *Geology* **2003**, *31*, 905–908, doi:10.1130/G19703.1.
174. Micherdziński, W. *Die Familie Parasitidae Oudemans 1901 (Acarina, Mesostigmata)*; Państwowe Wydawnictwo Naukowe: Kraków, Poland, 1969; pp. 1–690.
175. Kamczyc, J.; Gwiazdowicz, D.J. Soil mites (Acari, Mesostigmata) from Szczeliniec Wielki in the Stołowe Mountains National Park (SW Poland). *Biological Lett.* **2009**, *46*, 21–27, doi:10.2478/v10120-009-0010-4.
176. Frenot, Y.; Chown, S.L.; Whinam, J.; Selkirk, P.; Convey, P.; Skotnicki, M.; Bergstrom, D. Biological invasions in the Antarctic: Extent, impacts and implications. *Biol. Rev.* **2005**, *80*, 45–72, doi:10.1017/S1464793104006542.
177. Bergstrom, D.M.; Lucieer, A.; Kiefer, K.; Wasley, J.; Belbin, L.; Pedersen, T.K.; Chown, S.L. Indirect effects of invasive species removal devastate World Heritage Island. *J. Appl. Ecol.* **2009**, *46*, 73–81, doi:10.1111/j.1365-2664.2008.01601.x.
178. Convey, P. Terrestrial biodiversity in Antarctica—Recent advances and future challenges. *Polar Sci.* **2010**, *4*, 135–147, doi:10.1016/j.polar.2010.03.003.
179. Hågvar, S.; Solhøy, T.; Mong, C.E. Primary succession of soil mites (Acari) in a Norwegian glacier foreland, with emphasis on oribatid species. *Arct. Antarct. Alp. Res.* **2009**, *41*, 219–227, doi:10.1657/1938-4246-41.2.219.
180. Buda, J.; Olszanowski, Z.; Wierzgoń, M.; Zawierucha, K. Tardigrades and oribatid mites in bryophytes from geothermally active lava fields (Krafla, Iceland) and the description of *Pilatobius islandicus* sp. nov. (Eutardigrada). *Pol. Polar Res.* **2018**, *39*, 425–453, doi:10.24425/118755.
181. Hågvar, S.; Pedersen, A. Food choice of invertebrates during early glacier foreland succession. *Arct. Antarct. Alp. Res.* **2015**, *47*, 561–572.
182. Walter, D.E.; Proctor, H.C. *Mites: Ecology, Evolution & Behaviour*; Springer: Dordrecht, The Netherlands; Heidelberg, Germany; New York, NY, USA; London, UK, 2013; doi:10.1007/978-94-007-7164-2.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).