

Nanoplasmonic Laser Fusion Response to Földes and Pokol - Letter to the Editor

László P. Csernai¹, Norbert Kroó^{2,3}, István Papp², Daniel D. Strottman⁴

¹ Dept. of Physics and Technology, Univ. of Bergen, Norway

² Wigner Research Centre for Physics, Budapest, Hungary

³ Hungarian Academy of Sciences, Budapest, Hungary

⁴ Los Alamos National Laboratory, Los Alamos, NM, USA

(Dated: August 25, 2020)

Földes and Pokol in their letter "Inertial fusion without compression does not work either with or without nanoplasmonics" criticized our works [1, 2]. Here we refute their argumentation. Our proposed improvement is the combination of two basic research discoveries: (i) The possibility of detonations on space-time hyper-surfaces with time-like normal (i.e. simultaneous detonation in a whole volume) and (ii) to increase the ignition volume to the whole target, by regulating the laser light absorption using nanoantennas. These principles can be realized in an in-line, one dimensional configuration, in the simplest way with two opposing laser beams as in particle colliders [3, 4].

First of all the title of the letter of Földes and Pokol is misleading. Already in the abstract of [2] it is stated that we need to achieve limited compression to avoid the Rayleigh-Taylor instabilities. Furthermore it is written that the aim is to achieve fusion "without too much pre-compression". It is false from Földes and Pokol to state in their conclusion that we "... misjudged the possibility of the application of the uncompressed inertial fusion scheme.". We have never had such a statement.

In [2] the LLNL NIF compression was discussed in some detail just as it is done by Földes and Pokol [5]. The difficulties of this NIF setup by an ablator are elaborated in both of these publications.

Földes and Pokol does not mention our work, where the improvements we propose are elaborated in greater detail [3], and the experimental work showing that with the simplified linear irradiation from both sides can achieve the same compression as reached at LLNL NIF [4]. These works provide a theoretical and experimental response to the comments of Földes and Pokol.

In recent years the Laser Wake Field Acceleration (LWFA) became a well known concept with useful applications. An intensive laser pulse impinging on a target creates a high density plasma of $4 \times 10^{19} / \text{cm}^3$, and a wake field wave follows the pulse. This, non-linear wave in dense plasma is formed of the EM-field, electrons and ions. A typical laser pulse of 20 mJ intensity, 7 fs length and λ wave length can create a Laser Wake Field (LWF) dense plasma wave of about 10λ wavelength. Electrons do not prevent the development of this Laser Wake Field (LWF) waves. In the configuration presented in [3, 4] such LWF waves develop and move opposite to each other like in accelerators in collider configuration. This is suggested in ref. [3] for laser driven fusion. Here two known effects were combined. First, the possibility of detonations on space-time hyper-surfaces with time-like normal, so called time-like detonations [1, 6] which were found theoretically and experimentally in high en-

ergy heavy ion collision in the couple of last decades. This simultaneous volume ignition eliminates the possibility of Rayleigh-Taylor instabilities, which is a serious obstacle in achieving laboratory scale nuclear fusion. The second effect we use is to achieve simultaneous detonation in the whole volume of the target, by regulating the laser light absorption with nano-shells or nano-rods. This second effect is already experimentally proven in validation experiments, at Wigner RCP, Budapest, the institute of Földes [7].

Földes and Pokol under points 1., 2., 3., describe the thermonuclear setup of ICF where a dense hot-spot is created in $\approx 10\text{-}20$ ns, in the middle of the target and then the flame front "slowly" propagates with α -heating to the outside radius, R , of the target. As we emphasize in all previous publications we do not follow this dynamics, instead we aim to reach the condition for ignition in $\approx 10\text{-}100$ fs, in most of the target volume, i.e. on a time-like hypersurface. Then the flame front does not have to propagate anywhere, only the local burning rate matters! In the short ignition time the ions cannot expand.

This was also emphasized (even in the title) in the above mentioned, published experiment using colliding laser beams [4] achieved high target density for nuclear fusion. On the other hand they did not utilize LWF waves and nanoplasmonics and did not attempt to have simultaneous transition in the target. Thus, our present proposal differs from this experiment in the two novel aspects.

In the publications commented on by Földes and Pokol, we focused on our two fundamental and novel ideas and did not discuss the aspects, which should not be modified. In Laser Wake Field Collider, the target has two sides, which are initially accelerated towards each other in a pre-acceleration and pre-compression process, on a nanosecond time scale, just like at LLNL NIF and in [4]. The LWFA may reach the GeV/nucleon energy range.

In ref. [8] it was shown that intense laser pulse irra-

diating a combination target can accelerate Carbon ions to the TeV level by the laser plasma wakefield.

If we consider LWFC with a double layer pre-compressed to ion density, n_{pc} , and pre-accelerated to several GeV/nucleon energy, (i.e. to a velocity near to the speed of light, $v_{LWFC} \approx c$), the two LWF waves can inter penetrate and lead to an ignition reaction rate of

$$2 c \gamma^2 n_{pc}^2 \sigma,$$

due to the Lorentz contraction of the two ion bunches to γn_{pc} , and where σ is the ion-ion cross section. This may well exceed the thermal rate of $n\langle v_{th}\sigma \rangle$. If we accelerate the ions to 5 GeV/nucleon, then $\gamma \approx 6$, and if we achieve a pre-compression of factor 8 (considerably less than at NIF, where the 3-Dim compression reaches 800 g/cm³), then our burning rate is $2 \cdot \gamma^2 \frac{8^2 n^2}{800 n} (c/v_{th}) \approx 270$ times bigger than at NIF. Here we assumed that the average thermal collision speed is $v_{th} \approx c/2$.

So, the non-thermal non-equilibrium Laser Wake Field Collider mechanism may well exceed the thermal ignition rate by the adiabatic compression and heating at NIF. Especially if this ignition at NIF takes place for a central hot-spot only and then the flame should propagate over the rest of the target. The extreme energy need mentioned at the end of point 3., arises from the assumption of the thermal, setup, that we do not follow.

According to our preliminary calculations (yet unpublished) we can see that at around 125 - 150 fs after the target and projectile touch, the two LWF waves constructively interact and the EM field strength is maximal. This moment of time would be adequate for a short, intensive ignition pulse.

As we increase the target density the EM field penetration into the target is reduced, as the energy and mo-

mentum are passed to the target and projectile ions. The longitudinal momentum is transported to the kinetic motion of the target particles, which then contribute to the pinch effect reducing the EM strength and target beam directed momentum.

Under points 4., and 5., the target opacity and absorptivity are discussed. Here we want to emphasize that our nanoantennas are embedded within the target, so all energy reaching these remain in the target, so it is also absorbed. We have no extreme losses that occur due to evaporating the external hohlraum. The conversion of the absorbed energy to higher frequencies happens within the nanoantennas or can lead to direct nuclear burning.

Under point 6. Földes and Pokol assumes that the electrons will remain at the surface of the target. This fully contradicts with the structure of the LWF waves where the electrons are well separated from the ions. The electrons within the target are concentrated in the plasmonic waves on the surface of the nanoantennas. There is no significant electron reflection observed in LWFA, and even if, these electrons would just again remain in the target fuel in the LWFC configuration.

The conclusion of Földes and Pokol that "Therefore the nanostructures inside the target are of no use at all" is already proven to be false, as shown in the lower energy validation experiments at Wigner RCP, with laser beams of up to 25 mJ pulse energy, and polymer targets, which exhibit phase or structural transition at given threshold energy. This is also discussed already in ref. [3] before the experiments were performed.

Summarizing the presented results both theoretically, and based on recent experiments, (both in fusion related and high energy heavy ion fields), we can state that the conclusions of Földes and Pokol are unsubstantiated.

-
- [1] L.P. Csernai, & D.D. Strottman, Volume Ignition via Time-Like Detonation in Pellet Fusion, *Laser and Particle Beams* 33, 279-282 (2015).
- [2] L.P. Csernai, N. Kroó, & I. Papp, Radiation-Dominated Implosion with Nano-Plasmonics, *Laser and Particle Beams* 36, 171-178 (2018).
- [3] L.P. Csernai, M. Csete, I.N. Mishustin, A. Motorenko, I. Papp, L.M. Satarov, H. Stöcker & N. Kroó, Radiation-Dominated Implosion with Flat Target, *Physics and Wave Phenomena*, accepted February 3, 2020, (arXiv:1903.10896v3)
- [4] G. Zhang, M. Huan, A. Bonasera, Y. G. Ma, B. F. Shen, H. W. Wang, J. C. Xu, G. T. Fan, H. J. Fu, H. Xue, H. Zheng, L. X. Liu, S. Zhang, W. J. Li, X. G. Cao, X. G. Deng, X. Y. Li, Y. C. Liu, Y. Yu, Y. Zhang, C. B. Fu, and X. P. Zhang, Nuclear probes of an out-of-equilibrium plasma at the highest compression, *Phys. Lett. A* 383 (19), 2285-2289 (2019).
- [5] I.B. Földes and G.I. Pokol, "Inertial fusion without compression does not work either with or without nanoplasmonics, *Laser and Particle Beams* 39, 1-3 (2020).
- [6] L.P. Csernai, Detonation on Timelike Front for Relativistic Systems, School of Physics, University of Minnesota, Minneapolis, Minnesota, USA, *Zh. Eksp. Teor. Fiz.* 92, 397-386 (1987), & *Sov. Phys. JETP* 65, 219 (1987).
- [7] N. Kroó & P. Rácz, Plasmonics - The Interaction of Light with Metal Surface Electrons, *Laser Physics* 26, 084011 (2016).
- [8] F.L. Zheng, S.Z. Wu, C.T. Zhou, H.Y. Wang, X.Q. Yan and X.T. He, An ultra-short and TeV quasi-monoenergetic ion beam generation by laser wakefield accelerator in the snowplow regime, *Europhys. Lett. (EPL)*, 95, 55005 (2011).