



Metric Dimension Parameterized By Treewidth

Édouard Bonnet¹ · Nidhi Purohit^{1,2}

Received: 6 February 2020 / Accepted: 21 January 2021 / Published online: 23 February 2021
© The Author(s) 2021

Abstract

A resolving set S of a graph G is a subset of its vertices such that no two vertices of G have the same distance vector to S . The METRIC DIMENSION problem asks for a resolving set of minimum size, and in its decision form, a resolving set of size at most some specified integer. This problem is NP-complete, and remains so in very restricted classes of graphs. It is also W[2]-complete with respect to the size of the solution. METRIC DIMENSION has proven elusive on graphs of bounded treewidth. On the algorithmic side, a polynomial time algorithm is known for trees, and even for outerplanar graphs, but the general case of treewidth at most two is open. On the complexity side, no parameterized hardness is known. This has led several papers on the topic to ask for the parameterized complexity of METRIC DIMENSION with respect to treewidth. We provide a first answer to the question. We show that METRIC DIMENSION parameterized by the treewidth of the input graph is W[1]-hard. More refinedly we prove that, unless the Exponential Time Hypothesis fails, there is no algorithm solving METRIC DIMENSION in time $f(\text{pw})n^{o(\text{pw})}$ on n -vertex graphs of constant degree, with pw the pathwidth of the input graph, and f any computable function. This is in stark contrast with an FPT algorithm of Belmonte et al. (SIAM J Discrete Math 31(2):1217–1243, 2017) with respect to the combined parameter $\text{tl} + \Delta$, where tl is the tree-length and Δ the maximum-degree of the input graph.

Keywords Metric dimension · Treewidth · Parameterized hardness

✉ Nidhi Purohit
nidhipurohit95@gmail.com

Édouard Bonnet
edouard.bonnet@ens-lyon.fr

¹ CNRS, ENS de Lyon, Univ Lyon, Université Claude Bernard Lyon 1, LIP UMR5668, Villeurbanne, France

² University of Bergen, Bergen, Norway

1 Introduction

The METRIC DIMENSION problem has been introduced in the 1970s independently by Slater [22] and by Harary and Melter [13]. Given a graph G and an integer k , METRIC DIMENSION asks for a subset S of vertices of G of size at most k such that every vertex of G is uniquely determined by its distances to the vertices of S . Such a set S is called a *resolving set*, and a resolving set of minimum-cardinality is called a *metric basis*. The metric dimension of graphs finds application in various areas including network verification [2], chemistry [4], and robot navigation [18].

METRIC DIMENSION is an entry of the celebrated book on intractability by Garey and Johnson [12] where the authors show that it is NP-complete. In fact METRIC DIMENSION remains NP-complete in many restricted classes of graphs such as planar graphs [6], split, bipartite, co-bipartite graphs, and line graphs of bipartite graphs [9], interval graphs of diameter two [11], permutation graphs of diameter two [11], and in a subclass of unit disk graphs [15]. Furthermore METRIC DIMENSION cannot be solved in subexponential-time unless 3-SAT can [1]. On the positive side, the problem is polynomial-time solvable on trees [13, 18, 22]. Diaz et al. [6] generalize this result to outerplanar graphs. Fernau et al. [10] give a polynomial-time algorithm on chain graphs. Epstein et al. [9] show that METRIC DIMENSION (and even its vertex-weighted variant) can be solved in polynomial time on cographs and forests augmented by a constant number of edges. Hoffmann et al. [16] obtain a linear algorithm on cactus block graphs.

Hartung and Nichterlein [14] prove that METRIC DIMENSION is W[2]-complete (parameterized by the size of the solution k) even on subcubic graphs. Therefore an FPT algorithm solving the problem is unlikely. However Foucaud et al. [11] give an FPT algorithm with respect to k on interval graphs. This result is later generalized by Belmonte et al. [3] who obtain an FPT algorithm with respect to $tl + \Delta$ (where tl is the tree-length and Δ is the maximum-degree of the input graph), implying one for parameter $tl + k$. Indeed interval graphs, and even chordal graphs, have constant tree-length. Hartung and Nichterlein [14] presents an FPT algorithm parameterized by the vertex cover number, Eppstein [8], by the max leaf number, and Belmonte et al. [3], by the modular-width (a larger parameter than clique-width).

The complexity of METRIC DIMENSION parameterized by treewidth is quite elusive. It is discussed [8] or raised as an open problem in several papers [3, 6]. On the one hand, it was not known, prior to our paper, if this problem is W[1]-hard. On the other hand, the complexity of METRIC DIMENSION in graphs of treewidth at most two is still an open question.

1.1 Our Contribution

We settle the parameterized complexity of METRIC DIMENSION with respect to treewidth. We show that this problem is W[1]-hard, and we rule out, under the Exponential Time Hypothesis (ETH), an algorithm running in $f(tw)|V(G)|^{o(tw)}$, where G is the input graph, tw its treewidth, and f any computable function. Our reduction

even shows that an algorithm in time $f(\text{pw})|V(G)|^{o(\text{pw})}$ is unlikely on constant-degree graphs, for the larger parameter pathwidth pw . This is in stark contrast with the FPT algorithm of Belmonte et al. [3] for the parameter $\text{tl} + \Delta$ where tl is the tree-length and Δ is the maximum-degree of the graph. We observe that this readily gives an FPT algorithm for $\text{ctw} + \Delta$ where ctw is the connected treewidth, since $\text{ctw} \geq \text{tl}$. This unravels an interesting behavior of METRIC DIMENSION, at least on bounded-degree graphs: usual tree-decompositions are not enough for efficient solving. Instead one needs tree-decompositions with an additional guarantee that the vertices of a same bag are at a bounded distance from each other.

As our construction is quite technical, we chose to introduce an intermediate problem dubbed k -MULTICOLORED RESOLVING SET in the reduction from k -MULTICOLORED INDEPENDENT SET to METRIC DIMENSION. The first half of the reduction, from k -MULTICOLORED INDEPENDENT SET to k -MULTICOLORED RESOLVING SET, follows a generic and standard recipe to design parameterized hardness with respect to treewidth. The main difficulty is to design an effective *propagation gadget* with a constant-size left-right cut. The second half brings some new local attachments to the produced graph, to bridge the gap between k -MULTICOLORED RESOLVING SET and METRIC DIMENSION. Along the way, we introduce a number of gadgets: edge, propagation, forced set, forced vertex. They are quite streamlined and effective. Therefore, we believe these building blocks may help in designing new reductions for METRIC DIMENSION.

1.2 Organization of the Paper

In Sect. 2 we introduce the definitions, notations, and terminology used throughout the paper. In Sect. 3 we present the high-level ideas to establish our result. We define the k -MULTICOLORED RESOLVING SET problem which serves as an intermediate step for our reduction. In Sect. 4 we design a parameterized reduction from the $W[1]$ -complete k -MULTICOLORED INDEPENDENT SET to k -MULTICOLORED RESOLVING SET parameterized by treewidth. In Sect. 5 we show how to transform the produced instances of k -MULTICOLORED RESOLVING SET to METRIC DIMENSION-instances (while maintaining bounded treewidth). In Sect. 6 we conclude with some open questions.

2 Preliminaries

We denote by $[i, j]$ the set of integers $\{i, i + 1, \dots, j - 1, j\}$, and by $[i]$ the set of integers $[1, i]$. If \mathcal{X} is a set of sets, we denote by $\cup \mathcal{X}$ the union of them.

2.1 Graph Notations

All our graphs are undirected and simple (no multiple edge nor self-loop). We denote by $V(G)$, respectively $E(G)$, the set of vertices, respectively of edges, of the graph G . For $S \subseteq V(G)$, we denote the *open neighborhood* (or simply *neighborhood*) of S by $N_G(S)$, i.e., the set of neighbors of S deprived of S , and the *closed neighborhood* of

S by $N_G[S]$, i.e., the set $N_G(S) \cup S$. For singletons, we simplify $N_G(\{v\})$ into $N_G(v)$, and $N_G[\{v\}]$ into $N_G[v]$. We denote by $G[S]$ the subgraph of G induced by S , and $G - S := G[V(G) \setminus S]$. For $S \subseteq V(G)$ we denote by \bar{S} the complement $V(G) \setminus S$. For $A, B \subseteq V(G)$, $E(A, B)$ denotes the set of edges in $E(G)$ with one endpoint in A and the other one in B .

The length of a path in an unweighted graph is simply the number of edges of the path. For two vertices $u, v \in V(G)$, we denote by $\text{dist}_G(u, v)$, the distance between u and v in G , that is the length of the shortest path between u and v . The diameter of a graph is the longest distance between a pair of its vertices. The diameter of a subset $S \subseteq V(G)$, denoted by $\text{diam}_G(S)$, is the longest distance between a pair of vertices in S . Note that the distance is taken in G , not in $G[S]$. In particular, when G is connected, $\text{diam}_G(S)$ is finite for every S . A *pendant* vertex is a vertex with degree one. A vertex u is *pendant to* v if v is the only neighbor of u . Two distinct vertices u, v such that $N(u) = N(v)$ are called *false twins*, and *true twins* if $N[u] = N[v]$. In particular, true twins are adjacent. In all the above notations with a subscript, we omit it whenever the graph is implicit from the context.

2.2 Treewidth, Pathwidth, Connected Treewidth, and Tree-Length

A *tree-decomposition* of a graph G , is a tree T whose nodes are labeled by subsets of $V(G)$, called *bags*, such that for each vertex $v \in V(G)$, the bags containing v induce a non-empty subtree of T , and for each edge $e \in E(G)$, there is at least one bag containing both endpoints of e . A *connected tree-decomposition* further requires that each bag induces a connected subgraph in G . The width of a (connected) tree-decomposition is the size of its largest bag minus one. The treewidth (resp. connected treewidth) of a graph G is the minimum width of a tree-decomposition (resp. a connected tree-decomposition) of G . The length of a bag is the maximum distance in G between any pair of vertices in the bag. The length of a tree-decomposition is the maximum length of any bag. The tree-length of a graph G is the minimum length of a tree-decomposition of G . We denote the treewidth, connected treewidth, and tree-length of a graph by tw , ctw , and tl respectively. Since a connected graph on n vertices has diameter at most $n - 1$, it holds that $\text{ctw} \geq \text{tl}$.

The pathwidth is the same as treewidth except the tree T is now required to be a path, and hence is called a path-decomposition. In particular pathwidth is always larger than treewidth. Later we will need to upper bound the pathwidth of our constructed graph. Since writing down a path-decomposition is a bit cumbersome, we will rely on the following characterization of pathwidth. Kirousis and Papadimitriou [19] show the equality between the interval thickness number, which is known to be pathwidth plus one, and the *node searching number*. Thus we will only need to show that the number of searchers required to win the following one-player game is bounded by a suitable function. We imagine the edges of a graph to be contaminated by a gas. The task is to move around a team of searchers, placed at the vertices, in order to clean all the edges. A move consists of removing a searcher from the graph, adding a searcher at an unoccupied vertex, or displacing a searcher from a vertex to any other vertex (not necessarily adjacent). An edge is cleaned when both

its endpoints are occupied by a searcher. However after each move, all the cleaned edges admitting a free-of-searchers path from one of its endpoints to the endpoint of a contaminated edge are recontaminated. The node searching number is the minimum number of searchers required to win the game.

2.3 Parameterized Problems and Algorithms

Parameterized complexity aims to solve hard problems in time $f(k)|\mathcal{I}|^{O(1)}$, where k is a parameter of the instance \mathcal{I} which is hopefully (much) smaller than the total size of \mathcal{I} . More formally, a *parameterized problem* is a pair (Π, κ) where $\Pi \subseteq L$ for some language $L \subseteq \Sigma^*$ over a finite alphabet Σ (e.g., the set of words, graphs, etc.), and κ is a mapping from L to \mathbb{N} . An element $\mathcal{I} \in L$ is called an *instance* (or *input*). The mapping κ associates each instance to an integer called *parameter*. An instance is said *positive* if $\mathcal{I} \in \Pi$, and a *negative* otherwise. We denote by $|\mathcal{I}|$ the size of \mathcal{I} , that can be thought of as the length of the *word* \mathcal{I} . An *FPT algorithm* is an algorithm which solves a parameterized problem (Π, κ) , i.e., decides whether or not an input $\mathcal{I} \in L$ is positive, in time $f(\kappa(\mathcal{I}))|\mathcal{I}|^{O(1)}$ for some computable function f . We refer the interested reader to recent textbooks in parameterized algorithms and complexity [5, 7].

2.4 Exponential Time Hypothesis, FPT Reductions, and W[1]-Hardness

The *Exponential Time Hypothesis* (ETH) is a conjecture by Impagliazzo et al. [17] asserting that there is no $2^{o(n)}$ -time algorithm for 3-SAT on instances with n variables. Lokshantov et al. [20] survey conditional lower bounds under the ETH.

An *FPT reduction* from a parameterized problem $(\Pi \subseteq L, \kappa)$ to a parameterized problem $(\Pi' \subseteq L', \kappa')$ is a mapping $\rho : L \mapsto L'$ such that for every $\mathcal{I} \in L$:

- (1) $\mathcal{I} \in \Pi \Leftrightarrow \rho(\mathcal{I}) \in \Pi'$,
- (2) $|\rho(\mathcal{I})| \leq f(\kappa(\mathcal{I}))|\mathcal{I}|^{O(1)}$ for some computable function f , and
- (3) $\kappa(\rho(\mathcal{I})) \leq g(\kappa(\mathcal{I}))$ for some computable function g .

We further require that for every \mathcal{I} , we can compute $\rho(\mathcal{I})$ in FPT time $h(\kappa(\mathcal{I}))|\mathcal{I}|^{O(1)}$ for some computable function h . Condition (1) makes ρ a valid reduction, condition (2) together with the further requirement on the time to compute $\rho(\mathcal{I})$ make the mapping ρ *FPT*, and condition (3) controls that the new parameter $\kappa(\rho(\mathcal{I}))$ is bounded by a function of the original parameter $\kappa(\mathcal{I})$. One can therefore observe that using ρ in combination with an FPT algorithm solving (Π', κ') yields an FPT procedure to solve the initial problem (Π, κ) .

A standard use of an FPT reduction is to derive conditional lower bounds: if a problem (Π, κ) is thought not to admit an FPT algorithm, then an FPT reduction from (Π, κ) to (Π', κ') indicates that (Π', κ') is also unlikely to admit an FPT algorithm. We refer the reader to the textbooks [5, 7] for a formal definition of W[1]-hardness. For the purpose of this paper, we will just state that W[1]-hard are parameterized problems that are unlikely to be FPT, and that the following problem is

W[1]-complete even when all the V_i have the same number of elements, say t (see for instance [21]).

k -MULTICOLORED INDEPENDENT SET (k -MIS) Parameter: k
Input: An undirected graph G , an integer k , and (V_1, \dots, V_k) a partition of $V(G)$.
Question: Is there a set $I \subseteq V(G)$ such that $|I \cap V_i| = 1$ for every $i \in [k]$, and $G[I]$ is edgeless?

Every parameterized problem that k -MULTICOLORED INDEPENDENT SET FPT-reduces to is W[1]-hard. Our paper is thus devoted to designing an FPT reduction from k -MULTICOLORED INDEPENDENT SET to METRIC DIMENSION parameterized by tw . Let us observe that the ETH implies that one (equivalently, every) W[1]-hard problem is not in the class of problems solvable in FPT time ($FPT \neq W[1]$). Thus if we admit that there is no subexponential algorithm solving 3-SAT, then k -MULTICOLORED INDEPENDENT SET is not solvable in time $f(k)|V(G)|^{O(1)}$. Actually under this stronger assumption, k -MULTICOLORED INDEPENDENT SET is not solvable in time $f(k)|V(G)|^{o(k)}$. A concise proof of that fact can be found in the survey on the consequences of ETH [20].

2.5 Metric Dimension, Resolved Pairs, Distinguished Vertices

A pair of vertices $\{u, v\} \subseteq V(G)$ is said to be *resolved* by a set S if there is a vertex $w \in S$ such that $\text{dist}(w, u) \neq \text{dist}(w, v)$. A vertex u is said to be *distinguished* by a set S if for any $w \in V(G) \setminus \{u\}$, there is a vertex $v \in S$ such that $\text{dist}(v, u) \neq \text{dist}(v, w)$. A *resolving set* of a graph G is a set $S \subseteq V(G)$ such that every two distinct vertices $u, v \in V(G)$ are resolved by S . Equivalently, a resolving set is a set S such that every vertex of G is distinguished by S . Then METRIC DIMENSION asks for a resolving set of size at most some threshold k . Note that a resolving set of minimum size is sometimes called a *metric basis* for G .

METRIC DIMENSION (MD) Parameter: $tw(G)$
Input: An undirected graph G and an integer k .
Question: Does G admit a resolving set of size at most k ?

Here we anticipate on the fact that we will mainly consider METRIC DIMENSION parameterized by treewidth. Henceforth we sometimes use the notation Π/tw to emphasize that Π is not parameterized by the natural parameter (size of the resolving set) but by the treewidth of the input graph.

3 Outline of the W[1]-Hardness Proof of METRIC DIMENSION/ tw

We will show the following.

Theorem 3.1 *Unless the ETH fails, there is no computable function f such that METRIC DIMENSION can be solved in time $f(\text{pw})n^{o(\text{pw})}$ on constant-degree n -vertex graphs.*

We first prove that the following variant of METRIC DIMENSION is $W[1]$ -hard.

<p>k-MULTICOLORED RESOLVING SET (k-MRS)</p> <p>Input: An undirected graph G, an integer k, a set \mathcal{X} of q disjoint subsets of $V(G)$: X_1, \dots, X_q, and a set \mathcal{P} of pairs of vertices of G: $\{x_1, y_1\}, \dots, \{x_h, y_h\}$.</p> <p>Question: Is there a set $S \subseteq V(G)$ of size q such that</p> <ul style="list-style-type: none"> – (i) for every $i \in [q]$, $S \cap X_i = 1$, and – (ii) for every $p \in [h]$, there is an $s \in S$ satisfying $\text{dist}_G(s, x_p) \neq \text{dist}_G(s, y_p)$? 	<p>Parameter: $\text{tw}(G)$</p>
--	--

In words, in this variant the resolving set is made by picking exactly one vertex in each set of \mathcal{X} , and not all the pairs should be resolved but only the ones in a prescribed set \mathcal{P} . We call *critical pair* a pair of \mathcal{P} . In the context of k -MULTICOLORED RESOLVING SET, we call *legal set* a set which satisfies the former condition, and *resolving set* a set which satisfies the latter. Thus a solution for k -MULTICOLORED RESOLVING SET is a legal resolving set.

The reduction from k -MULTICOLORED INDEPENDENT SET starts with a well-established trick to show parameterized hardness by treewidth. We create m “empty copies” of the k -MIS-instance $(G, k, (V_1, \dots, V_k))$, where $m := |E(G)|$ and $t := |V_i|$. We force exactly one vertex in each color class of each copy to be in the resolving set, using the set \mathcal{X} . In each copy, we introduce an edge gadget for a single (distinct) edge of G . Encoding an edge of k -MIS in the k -MRS-instance is fairly simple: we build a pair (of \mathcal{P}) which is resolved by every choice but the one *selecting both its endpoints* in the resolving set. We now need to force a *consistent choice of the vertex chosen in V_i* over all the copies. We thus design a propagation gadget. A crucial property of the propagation gadget, for the pathwidth of the constructed graph to be bounded, is that it admits a cut of size $O(k)$ disconnecting one copy from the other. Encoding a choice in V_i in the distances to four special vertices, called *gates*, we manage to build such a gadget with constant-size “left-right” separator per color class. This works by introducing t pairs (of \mathcal{P}) which are resolved by the south-west and north-east gates but not by the south-east and north-west ones. Then we link the vertices of a copy of V_i in a way that the higher their index, the more pairs they resolve in the propagation gadget to their left, and the fewer pairs they resolve in the propagation gadget to their right.

We then turn to the actual METRIC DIMENSION problem. We design a gadget which simulates requirement (i) by forcing a vertex of a specific set X in the resolving set. This works by introducing two pairs that are only resolved by vertices of X . We attach this new gadget, called *forcing set gadget*, to all the k color classes of the m copies. Finally we have to make sure that a candidate solution resolves all the pairs, and not only the ones prescribed by \mathcal{P} . For that we attach two adjacent “pendant” vertices to strategically chosen vertices. One of these two vertices have to be in the resolving set since they are true twins, hence not resolved by any other vertex. Then

everything is as if the unique common neighbor v of the true twins was added to the resolving set. Therefore we can perform this operation as long as v does not resolve any of the pairs of \mathcal{P} .

To facilitate the task of the reader, henceforth we stick to the following conventions:

- Index $i \in [k]$ ranges over the k rows of the k -MRS/MD-instance or color classes of k -MIS.
- Index $j \in [m]$ ranges over the m columns of the k -MRS/MD-instance or edges of k -MIS.
- Index $\gamma \in [t]$, ranges over the t vertices of a color class.

We invite the reader to look up Table 1 when in doubt about a notation/symbol relative to the construction.

4 Parameterized Hardness of k -MULTICOLORED RESOLVING SET/tw

In this section, we give an FPT reduction from the W[1]-complete k -MULTICOLORED INDEPENDENT SET to k -MULTICOLORED RESOLVING SET parameterized by treewidth. More precisely, given a k -MULTICOLORED INDEPENDENT SET-instance $(G, k, (V_1, \dots, V_k))$ we produce in polynomial-time an equivalent k -MULTICOLORED RESOLVING SET-instance $(G', k', \mathcal{X}, \mathcal{P})$ where G' has pathwidth (hence treewidth) $O(k)$.

4.1 Construction

Let $(G, k, (V_1, \dots, V_k))$ be an instance of k -MULTICOLORED INDEPENDENT SET where (V_1, \dots, V_k) is a partition of $V(G)$ and $V_i := \{v_{i,\gamma} \mid 1 \leq \gamma \leq t\}$. We arbitrarily number $e_1, \dots, e_j, \dots, e_m$ the m edges of G .

4.1.1 Overall Picture

We start with a high-level description of the k -MRS-instance $(G', k', \mathcal{X}, \mathcal{P})$. For each color class V_i , we introduce m copies $V_i^1, \dots, V_i^j, \dots, V_i^m$ of a selector gadget to G' . Each set V_i^j is added to \mathcal{X} , so a solution has to pick exactly one vertex within each selector gadget. One can imagine the vertex-sets V_i^1, \dots, V_i^m to be aligned on the i -th row, with V_i^j occupying the j -th column (see Fig. 1). Each V_i^j has t vertices denoted by $v_{i,1}^j, v_{i,2}^j, \dots, v_{i,t}^j$, where each $v_{i,\gamma}^j$ “corresponds” to $v_{i,\gamma} \in V_i$. We make $v_{i,1}^j, v_{i,2}^j \dots v_{i,t}^j$ a path with $t - 1$ edges.

For each edge $e_j \in E(G)$, we insert an edge gadget $\mathcal{G}(e_j)$ containing a pair of vertices $\{c_j, c'_j\}$ that we add to \mathcal{P} . Gadget $\mathcal{G}(e_j)$ is attached to V_i^j and $V_{i'}^j$, where $e_j \in E(V_i, V_{i'})$. The edge gadget is designed in a way that the only legal sets that do not resolve $\{c_j, c'_j\}$ are the ones that precisely pick $v_{i,\gamma}^j \in V_i^j$ and $v_{i',\gamma'}^j \in V_{i'}^j$ such that $e_j = v_{i,\gamma} v_{i',\gamma'}$. We add a propagation gadget $P_i^{j,j+1}$ between two consecutive copies V_i^j

Table 1 Glossary of the construction

Symbol/term	Definition/action
$\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$	Critical pair of the propagation gadget $P_i^{j,j+1}$
A_i^j	Set of vertices $\bigcup_{\gamma \in [t]} \{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$
bb_i^j	Bottom brown vertex, $v(v_{i,t}^j, r_i^j)$
bc_i^j	Bottom cyan vertex (smallest index γ)
bl_i^j	Neighbor of $v_{i,t}^j$ in $P_i^{j-1,j}$
Blue vertex	One of the four neighbors of V_i^j in the propagation gadgets
br_i^j	Neighbor of $v_{i,t}^j$ in $P_i^{j,j+1}$
Brown vertex	Vertices $v(v_{i,1}^j, p_i^j)$ and $v(v_{i,t}^j, r_i^j)$
$\{c_j, c'_j\}$	Critical pair of the edge gadget $\mathcal{G}(e_j)$
Cyan vertex	Neighbor of V_i^j in the paths to $\mathcal{G}(e_j)$
E_i^j	Vertices in the paths from V_i^j to $\mathcal{G}(e_j)$
$e_{i,\gamma}^j$	Alternative labeling of the cyan vertices, neighbor of $v_{i,\gamma}^j$
F	Set of all forced vertices, $\bigcup_{i \in [k], j \in [m]} F_i^j \cup \bigcup_{j \in [m]} F_j$
F_i^j	Set of forced vertices attached to neighbors of $\{\pi_i^j, \rho_i^j, nw_i^j, se_i^j\}$
F_j	Set of forced vertices attached to neighbors of g_j
$f(v)$	Forced vertex attached to a vertex v
$f'(v)$	True twin of $f(v)$
$\mathcal{G}(e_j)$	Edge gadget on $\{g_j, c_j, c'_j\}$ between V_i^j and $V_{i'}^j$, where $e_j \in E(V_i, V_{i'})$
mc_i^j	Middle cyan vertex (not top nor bottom)
ne_i^j	North-east gate of $P_i^{j,j+1}$
nw_i^j	North-west gate of $P_i^{j,j+1}$
ne_i^j, sw_i^j	Resolve the critical pairs of $P_i^{j,j+1}$
nw_i^j, se_i^j	Do not resolve the critical pairs of $P_i^{j,j+1}$
$v(u, v)$	Neighbor of u in the path $P(u, v)$
\mathcal{P}	List of critical pairs
$\{p_i^j, q_i^j\}$	Pair only resolved by vertices in $V_i^j \cup P(v_{i,1}^j, p_i^j) \cup \{q_i^j\}$
π_i^j	Gate of $\{p_i^j, q_i^j\}$, linked by paths to most neighbors of V_i^j
$P_i^{j,j+1}$	Propagation gadget between V_i^j and V_i^{j+1}
$P(u, v)$	Path added in the construction expressly between u and v
$\{r_i^j, s_i^j\}$	Pair only resolved by vertices in $V_i^j \cup P(v_{i,t}^j, r_i^j) \cup \{s_i^j\}$
ρ_i^j	Gate of $\{r_i^j, s_i^j\}$, linked by paths to most neighbors of V_i^j
se_i^j	South-east gate of $P_i^{j,j+1}$
sw_i^j	South-west gate of $P_i^{j,j+1}$
t	Size of each V_i
tb_i^j	Top brown vertex, $v(v_{i,1}^j, p_i^j)$
tc_i^j	Top cyan vertex (largest index γ)
tl_i^j	Neighbor of $v_{i,1}^j$ in $P_i^{j-1,j}$
tr_i^j	Neighbor of $v_{i,1}^j$ in $P_i^{j,j+1}$

Table 1 (continued)

Symbol/term	Definition/action
V_i	Partite set of G
V_i^j	“Copy of V_i ”, stringed by a path, in G' and G''
$v_{i,\gamma}^j$	Vertex of V_i^j representing $v_{i,\gamma} \in V(G)$
W_j	Endpoints in $V_i^j \cup V_{i'}^j$ of paths from $V_i^j \cup V_{i'}^j$ to $\mathcal{G}(e_j)$
\mathcal{X}	Set containing all the sets V_i^j for $i \in [k]$ and $j \in [m]$
X_j	Neighbors of W_j on the paths to $\mathcal{G}(e_j)$ (cyan vertices)

and V_i^{j+1} , where the indices in the superscript are taken modulo m . The role of the propagation gadget is to ensure that the choices in each V_i^j ($j \in [m]$) corresponds to the same vertex in V_i .

The intuitive idea of the reduction is the following. We say that a vertex of G' is *selected* if it is put in the resolving set of G' , a tentative solution. The propagation gadget $P_i^{j,j+1}$ ensures a consistent choice among the m copies V_i^1, \dots, V_i^m . The edge gadget ensures that the selected vertices of G' correspond to an independent set in the original graph G . If both the endpoints of an edge e_j are selected, then the pair $\{c_j, c'_j\}$ is not resolved.

We now detail the construction.

4.1.2 Selector Gadget

For each $i \in [k]$ and $j \in [m]$, we add to G' a path on $t - 1$ edges $v_{i,1}^j, v_{i,2}^j, \dots, v_{i,t}^j$, and denote this set of vertices by V_i^j . Each $v_{i,\gamma}^j$ corresponds to $v_{i,\gamma} \in V_i$. We call j -th column the set $\bigcup_{i \in [k]} V_i^j$, and i -th row, the set $\bigcup_{j \in [m]} V_i^j$. We set $\mathcal{X} := \{V_i^j\}_{i \in [k], j \in [m]}$. By definition of k -MULTICOLORED RESOLVING SET, a solution S has to satisfy that for every $i \in [k], j \in [m], |S \cap V_i^j| = 1$. We call *legal set* a set S of size $k' = km$ that satisfies this property. We call *consistent set* a legal set S which takes the “same” vertex in each row, that is, for every $i \in [k]$, for every pair $(v_{i,\gamma}^j, v_{i,\gamma'}^j) \in (S \cap V_i^j) \times (S \cap V_i^j)$, then $\gamma = \gamma'$.

4.1.3 Edge Gadget

For each edge $e_j = v_{i,\gamma} v_{i',\gamma'} \in E(G)$, we add an edge gadget $\mathcal{G}(e_j)$ in the j -th column of G' . $\mathcal{G}(e_j)$ consists of a path on three vertices: $c_j g_j c'_j$. The pair $\{c_j, c'_j\}$ is added to the list of critical pairs \mathcal{P} . We link both $v_{i,\gamma}^j$ and $v_{i',\gamma'}^j$ to g_j by a private path¹ of length $t + 2$. We link the at least two and at most four vertices $v_{i,\gamma-1}^j, v_{i,\gamma+1}^j, v_{i',\gamma'-1}^j, v_{i',\gamma'+1}^j$ (whenever they exist) to c_j by a private path of length $t + 2$. This defines at most six

¹ We use the expression *private path* to emphasize that the different sources get a pairwise internally vertex-disjoint path to the target.




Fig. 1 The overall picture with $k = 3$ color classes, $t = 5$ vertices per color class, $m = 6$ edges, $e_1 = v_{1,3}v_{2,4}$, $e_2 = v_{1,4}v_{2,1}$, $e_3 = v_{1,5}v_{3,1}$, etc. The dashed lines on the left and right symbolize that the construction is cylindrical (Color figure online)

paths from $V_i^j \cup V_{i'}^j$ to $\mathcal{G}(e_j)$. Let us denote by W_j the at most six endpoints of these paths in $V_i^j \cup V_{i'}^j$. For each $v \in W_j$, we denote by $P(v, j)$ the path from v to $\mathcal{G}(e_j)$. We set $E_i^j := \bigcup_{v \in W_j \cap V_i^j} P(v, j)$ and $E_{i'}^j := \bigcup_{v \in W_j \cap V_{i'}^j} P(v, j)$. We denote by X_j the set of the at most six neighbors of W_j on the paths to $\mathcal{G}(e_j)$. Henceforth we may refer to the vertices in some X_j as the *cyan vertices*. Individually we denote by $e_{i,\gamma}^j$ the cyan vertex neighbor of $v_{i,\gamma}^j$ in $P(v_{i,\gamma}^j, j)$. We observe that for fixed i and j , $e_{i,\gamma}^j$ exists for at most three values of γ . We add an edge between two cyan vertices if their respective neighbors in V_i^j are also linked by an edge (or equivalently, if they have consecutive “indices γ ”). These extra edges are useless in the k -MRS-instance, but will turn out useful in the MD-instance. See Fig. 2 for an illustration of the edge gadget.

The rest of the construction will preserve that for every $v \in (V_i^j \cup V_{i'}^j) \setminus \{v_{i,\gamma}^j, v_{i',\gamma'}^j\}$, $\text{dist}(v, c'_j) = \text{dist}(v, c_j) + 2$, and for each $v \in \{v_{i,\gamma}^j, v_{i',\gamma'}^j\}$, $\text{dist}(v, c_j) = \text{dist}(v, g_j) + 1 = \text{dist}(v, c'_j)$. In other words, the only two vertices of $V_i^j \cup V_{i'}^j$ not resolving the critical pair $\{c_j, c'_j\}$ are $v_{i,\gamma}^j$ and $v_{i',\gamma'}^j$, corresponding to the endpoints of e_j .

4.1.4 Propagation Gadget

Between each pair $(V_i^j, V_{i'}^{j+1})$, where $j + 1$ is taken modulo m , we insert an identical copy of the propagation gadget, and we denote it by $P_i^{j,j+1}$. It ensures that if the

vertex $v_{i,\gamma}^j$ is in a legal resolving set S , then the vertex of $S \cap V_i^{j+1}$ should be some $v_{i,\gamma'}^{j+1}$ with $\gamma \leq \gamma'$. The cylindricity of the construction and the fact that exactly one vertex of V_i^j is selected, will therefore impose that the set S is consistent.

$P_i^{j,j+1}$ comprises four vertices $sw_i^j, se_i^j, nw_i^j, ne_i^j$, called *gates*, and a set A_i^j of $2t$ vertices $a_{i,1}^j, \dots, a_{i,t}^j, \alpha_{i,1}^j, \dots, \alpha_{i,t}^j$. We make both $a_{i,1}^j, a_{i,2}^j \dots a_{i,t}^j$ and $\alpha_{i,1}^j, \alpha_{i,2}^j \dots \alpha_{i,t}^j$ a path with $t - 1$ edges. For each $\gamma \in [t]$, we add the pair $\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$ to the set of critical pairs \mathcal{P} . Removing the gates disconnects A_i^j from the rest of the graph.

We now describe how we link the gates to V_i^j, V_i^{j+1} , and A_i^j . We link $v_{i,1}^j$ (the “top” vertex of V_i^j) to sw_i^j and $v_{i,t}^j$ (the “bottom” vertex of V_i^j) to nw_i^j both by a path of length 2. We also link $v_{i,1}^{j+1}$ to se_i^j by a path of length 3, and $v_{i,t}^{j+1}$ to ne_i^j by a path of length 2. Then we make nw_i^j adjacent to $a_{i,1}^j$ and $\alpha_{i,1}^j$, while we make ne_i^j adjacent to $\alpha_{i,t}^j$ only. We make se_i^j adjacent to $a_{i,t}^j$ and $\alpha_{i,t}^j$, while we make sw_i^j adjacent to $a_{i,1}^j$ only. Finally, we add an edge between ne_i^j and nw_i^j , and between sw_i^j and se_i^j . See Fig. 3 for an illustration of the propagation gadget $P_i^{j,j+1}$ with $t = 5$.

Let us motivate the gadget $P_i^{j,j+1}$. One can observe that the gates ne_i^j and sw_i^j resolve the critical pairs of the propagation gadget, while the gates nw_i^j and se_i^j do not. Consider that the vertex added to the resolving set in V_i^j is $v_{i,\gamma}^j$. Its shortest paths to critical pairs *below* it (that is, with index $\gamma' > \gamma$) go through the gate sw_i^j , whereas its shortest paths to critical pairs at its level or above (that is, with index $\gamma' \leq \gamma$) go through the gate nw_i^j . Thus $v_{i,\gamma}^j$ only resolves the critical pairs $\{a_{i,\gamma'}^j, \alpha_{i,\gamma'}^j\}$ with $\gamma' > \gamma$. On the contrary, the vertex of the resolving set in V_i^{j+1} only resolves the critical pairs $\{a_{i,\gamma'}^j, \alpha_{i,\gamma'}^j\}$ at its level or above. This will force that its level is γ or below. Hence the vertices of the resolving in V_i^j and V_i^{j+1} should be such that $\gamma' \geq \gamma$. Since there is also a propagation gadget between V_i^m and V_i^1 , this circular chain of inequalities forces a global equality.

4.1.5 Wrapping Up

We put the pieces together as described in the previous subsections. At this point, it is convenient to give names to the neighbors of V_i^j in the propagation gadgets $P_i^{j-1,j}$ and $P_i^{j,j+1}$. We may refer to them as *blue vertices* (as they appear in Fig. 4). We denote by tl_i^j the neighbor of $v_{i,1}^j$ in $P_i^{j-1,j}$, tr_i^j the neighbor of $v_{i,1}^j$ in $P_i^{j,j+1}$, bl_i^j the neighbor of $v_{i,t}^j$ in $P_i^{j-1,j}$, and br_i^j the neighbor of $v_{i,t}^j$ in $P_i^{j,j+1}$. We add the following edges and paths.

For any pair i, j such that e_j has an endpoint in V_i , the vertices $tl_i^j, tr_i^j, bl_i^j, br_i^j$ are linked to g_j by a private path of length the distance of their unique neighbor in V_i^j to c_j . We add an edge between se_i^j and se_i^{j+1} , and between nw_i^j and nw_i^{j+1} (where $j + 1$ is modulo m). Finally, for every $e_j \in E(V_i, V_{i'})$, we add four paths between $se_{i'}^j, se_{i'}^j, nw_{i'}^j, nw_{i'}^j$ and $g_j \in \mathcal{G}(e_j)$. More precisely, for each $i'' \in \{i, i'\}$, we add a path from g_j to $se_{i''}^j$ of length $\text{dist}(g_j, sw_{i''}^j) - 4$, and a path from g_j to $nw_{i''}^j$ of length $\text{dist}(g_j, nw_{i''}^j) - 4$. These distances are taken in the graph before we introduced the new paths, and one can observe that the length of these paths is at least t . This finishes the construction.




Fig. 2 The edge gadget $\mathcal{G}(e_4)$ with $e_4 = v_{1,5}v_{3,3}$. Weighted edges are short-hands for subdivisions of the corresponding length. The edges between the cyan vertices will not be useful for the k -MRS-instance, but will later simplify the construction of the MD-instance (Color figure online)





Fig. 3 The propagation gadget $P_i^{j,j+1}$. The critical pairs $\{\alpha_{i,y}^j, \alpha_{i,y}^{j+1}\}$ are surrounded by thin dashed lines. The blue (resp. red) integer on a vertex of A_i^j is its distance to the blue (resp. red) vertex in V_i^j (resp. V_i^{j+1}). Note that the blue vertex distinguishes the critical pairs below it, while the red vertex distinguishes critical pairs at its level or above (Color figure online)

Fig. 4 The forget set gadget for V_i^j . The cyan vertices are the at most three neighbors of V_i^j in the potential paths to $\mathcal{G}(e_j)$. The lengths of the paths from the neighbors of V_i^j to π_i^j and ρ_i^j is defined in Sect. 5.1.1 (Color figure online)



We recall that, by a slight abuse of language, a *resolving set* in the context of k -MULTICOLORED RESOLVING SET is a set which resolves all the critical pairs of \mathcal{P} . In particular, it is not necessarily a resolving set in the sense of METRIC DIMENSION. With that terminology, a solution for k -MULTICOLORED RESOLVING SET is a legal resolving set.

4.2 Correctness of the Reduction

We now check that the reduction is correct. We start with the following technical lemma. If a set X contains a pair that no vertex of $N(X)$ (that is $N[X] \setminus X$) resolves, then no vertex outside X can distinguish the pair.

Lemma 4.1 *Let X be a subset of vertices, and $a, b \in X$ be two distinct vertices. If for every vertex $v \in N(X)$, $\text{dist}(v, a) = \text{dist}(v, b)$, then for every vertex $v \notin X$, $\text{dist}(v, a) = \text{dist}(v, b)$.*

Proof Let v be a vertex outside of X . We further assume that v is not in $N(X)$, otherwise we can already conclude that it does not distinguish $\{a, b\}$. A shortest path from v to a , has to go through $N(X)$. Let w_a be the first vertex of $N(X)$ met in this shortest path from v to a . Similarly, let w_b be the first vertex of $N(X)$ met in a shortest path from v to b . Since $w_a, w_b \in N(X)$, they satisfy $\text{dist}(w_a, a) = \text{dist}(w_a, b)$ and $\text{dist}(w_b, a) = \text{dist}(w_b, b)$. Then, $\text{dist}(v, a) \leq \text{dist}(v, w_b) + \text{dist}(w_b, a) = \text{dist}(v, w_b) + \text{dist}(w_b, b) = \text{dist}(v, b)$, and $\text{dist}(v, b) \leq \text{dist}(v, w_a) + \text{dist}(w_a, b) = \text{dist}(v, w_a) + \text{dist}(w_a, a) = \text{dist}(v, a)$. Thus $\text{dist}(v, a) = \text{dist}(v, b)$. □

We use the previous lemma to show that every vertex of a V_i^j only resolves critical pairs in gadgets it is attached to. This will be useful in the two subsequent lemmas.

Lemma 4.2 *For any $i \in [k]$, $j \in [m]$, and $v \in V_i^j$, v does not resolve any critical pair outside of P_i^{j-1} , P_i^{j+1} (where indices in the superscript are taken modulo m), and $\{c_j, c'_j\}$. Furthermore, if $e_j \in E(G)$ has no endpoint in $V_i \subseteq V(G)$, then v does not resolve $\{c_j, c'_j\}$.*

Proof We first show that $v \in V_i^j$ does not resolve any critical pair in propagation gadgets that are not P_i^{j-1} and P_i^{j+1} . Let $\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$ be a critical pair in a propagation gadget different from P_i^{j-1} and P_i^{j+1} . Let X be the connected component containing P_i^{j+1} of $G' - (\{nw_{i'}^{j-1}, se_{i'}^{j-1}, nw_{i'}^{j+1}, se_{i'}^{j+1}\} \cup C_e)$, where C_e comprises $\{c'_j, g'_j\}$ if e_j has an endpoint in $V_{i'}$ and $\{c_{j+1}, g_{j+1}\}$ if e_{j+1} has an endpoint in $V_{i'}$. Thus C_e has size 0, 2, or 4. One can observe that $N(X) = \{nw_{i'}^{j-1}, se_{i'}^{j-1}, nw_{i'}^{j+1}, se_{i'}^{j+1}\} \cup C_e$, that $V_i^j \cup V_i^{j+1} \subseteq X$, and that no “other V_i^j ” intersects X . In particular V_i^j is fully contained in $G - X$. We now check that no vertex of $N(X)$ resolves the pair $\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$ (which is inside X).

For each $u \in \{nw_{i'}^{j-1}, nw_{i'}^{j+1}\}$, it holds that $\text{dist}(u, a_{i,\gamma}^j) = \gamma + 1 = \text{dist}(u, \alpha_{i,\gamma}^j)$ (the shortest paths go through $nw_{i'}^j$), while for each $u \in \{se_{i'}^{j-1}, se_{i'}^{j+1}\}$, it holds that $\text{dist}(u, a_{i,\gamma}^j) = t - \gamma + 2 = \text{dist}(u, \alpha_{i,\gamma}^j)$ (the shortest paths go through $se_{i'}^j$). If they are part of C_e , g_j and c_j also do not resolve $\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$, the shortest paths going through the gates $nw_{i'}^j$ or $se_{i'}^j$, and respectively g_j and then the gates $nw_{i'}^j$ or $se_{i'}^j$. For the same reason, g_{j+1} and c_{j+1} do not resolve $\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$. Then we conclude by Lemma 4.1 that no vertex of V_i^j (in particular v) resolves $\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$, or any critical pair in P_i^j .

Let us now show that the pair $\{c_j, c'_j\}$ is not resolved by any vertex of $\cup \mathcal{A} \setminus (V_{i'}^j \cup V_{i''}^j)$ such that $e_j \in E(V_{i'}, V_{i''})$. Let $Y := \{tl_{i'}^j, tr_{i'}^j, bl_{i'}^j, br_{i'}^j, tl_{i''}^j, tr_{i''}^j, bl_{i''}^j, br_{i''}^j, nw_{i'}^j, se_{i'}^j, nw_{i''}^j, se_{i''}^j\}$, and X be the connected component containing g_j in $G' - Y$. Again one can observe that $N(X) = Y$, X contains $V_{i'}^j \cup V_{i''}^j$ but does not intersect any “other V_i^j ”. We therefore show that no vertex of Y resolves $\{c_j, c'_j\}$, and conclude with Lemma 4.1. All the vertices of Y have a private path to g_j whose length is such that they have a shortest path to c_j going through g_j . Therefore $\forall u \in Y, \text{dist}(u, c_j) = \text{dist}(u, g_j) + 1 = \text{dist}(u, c'_j)$. \square

The two following lemmas show the equivalences relative to the expected use of the edge and propagation gadgets. They will be useful in Sects. 4.2.1 and 4.2.2.

Lemma 4.3 *A legal set S resolves the critical pair $\{c_j, c'_j\}$ with $e_j = v_{i,\gamma} v_{i,\gamma'}$ if and only if the vertex $v_{i,\gamma}^j$ in $V_i^j \cap S$ and the vertex $v_{i,\gamma'}^j$ in $V_i^j \cap S$ satisfy $(\gamma, \gamma') \neq (\gamma_i, \gamma'_i)$.*

Proof By Lemma 4.2, no vertex of $S \setminus \{v_{i,\gamma}^j, v_{i,\gamma'}^j\}$ resolves $\{c_j, c'_j\}$. By construction of G' , $v_{i,\gamma}^j$ (resp. $v_{i,\gamma'}^j$) is the only vertex of V_i^j (resp. V_i^j) that does not resolve $\{c_j, c'_j\}$. Indeed the shortest paths of $v_{i,\gamma''}^j$, for $\gamma'' \geq \gamma + 1$, to $\{c_j, c'_j\}$ go through $v_{i,\gamma+1}^j$ which resolves the pair. Note that a shortest path between V_i^j and V_i^j has length at least

$2t + 4$, so a shortest path from $v_{i,\gamma''}^j$ to $\{c_j, c'_j\}$ cannot go through $V_{i,\gamma}^j$. Similarly the shortest paths of $v_{i,\gamma''}^j$, for $\gamma'' \leq \gamma - 1$, to $\{c_j, c'_j\}$ go through $V_{i,\gamma-1}^j$ which also resolves the pair. Thus only $v_{i,\gamma}^j$ (resp. $v_{i,\gamma'}^j$), whose shortest paths to $\{c_j, c'_j\}$ go via g_j , does not resolve this pair among V_i^j (resp. $V_{i'}^j$). Hence, the critical pair $\{c_j, c'_j\}$ is not resolved by S if and only if $v_{i,\gamma_1}^j = v_{i,\gamma}^j$ and $v_{i',\gamma'_1}^j = v_{i',\gamma'}^j$. \square

Lemma 4.4 *A legal set S resolves all the critical pairs of $P_i^{j,j+1}$ if and only if the vertex $v_{i,\gamma}^j$ in $V_i^j \cap S$ and the vertex $v_{i',\gamma'}^{j+1}$ in $V_{i'}^{j+1} \cap S$ satisfy $\gamma \leq \gamma'$.*

Proof By Lemma 4.2, no vertex of $S \setminus \{v_{i,\gamma}^j, v_{i,\gamma'}^{j+1}\}$ resolves a critical pair of $P_i^{j,j+1}$. Let us show that the critical pairs that $v_{i,\gamma}^j$ resolves in A_i^j are exactly the pairs $\{\alpha_{i,z}^j, \alpha'_{i,z}^j\}$ with $z > \gamma$. For any $z \in [t]$, it holds that $\text{dist}(v_{i,\gamma}^j, \alpha_{i,z}^j) = \min(t + 2 + z - \gamma, t + 2 + \gamma - z) = t + 2 + \min(z - \gamma, \gamma - z)$, and $\text{dist}(v_{i,\gamma}^j, \alpha'_{i,z}^j) = \min(t + 2 + z - \gamma, t + 3 + \gamma - z) = t + 2 + \min(z - \gamma, \gamma - z + 1)$. So if $z > \gamma$, $\text{dist}(v_{i,\gamma}^j, \alpha_{i,z}^j) = t + 2 + \gamma - z \neq t + 2 + \gamma - z + 1 = \text{dist}(v_{i,\gamma}^j, \alpha'_{i,z}^j)$. Whereas if $z \leq \gamma$, $\text{dist}(v_{i,\gamma}^j, \alpha_{i,z}^j) = t + 2 + z - \gamma = \text{dist}(v_{i,\gamma}^j, \alpha'_{i,z}^j)$.

Similarly, we show that the critical pairs that $v_{i,\gamma'}^{j+1}$ resolves in $A_{i'}^{j+1}$ are exactly the pairs $\{\alpha_{i,z}^{j+1}, \alpha'_{i,z}^{j+1}\}$ with $z \leq \gamma'$. For every $z \in [t]$, it holds that $\text{dist}(v_{i,\gamma'}^{j+1}, \alpha_{i,z}^{j+1}) = \min(t + 3 + z - \gamma', t + 3 + \gamma' - z) = t + 3 + \min(z - \gamma', \gamma' - z)$, and $\text{dist}(v_{i,\gamma'}^{j+1}, \alpha'_{i,z}^{j+1}) = \min(t + 2 + z - \gamma', t + 3 + \gamma' - z) = t + 2 + \min(z - \gamma', \gamma' - z + 1)$. So if $z \leq \gamma'$, $\text{dist}(v_{i,\gamma'}^{j+1}, \alpha_{i,z}^{j+1}) = t + 3 + z - \gamma' \neq t + 2 + z - \gamma' = \text{dist}(v_{i,\gamma'}^{j+1}, \alpha'_{i,z}^{j+1})$. Whereas if $z > \gamma'$, $\text{dist}(v_{i,\gamma'}^{j+1}, \alpha_{i,z}^{j+1}) = t + 3 + \gamma' - z = \text{dist}(v_{i,\gamma'}^{j+1}, \alpha'_{i,z}^{j+1})$. This implies that all the critical pairs of $A_{i'}^{j+1}$ are resolved by S if and only if $\gamma \leq \gamma'$. \square

We can now prove the correctness of the reduction. The construction can be computed in polynomial time in $|V(G)|$, and G' itself has size bounded by a polynomial in $|V(G)|$. We postpone checking that the pathwidth is bounded by $O(k)$ to the end of the second step, where we produce an instance of MD whose graph G'' admits G' as an induced subgraph.

4.2.1 k -Multicolored Independent Set in $G \Rightarrow$ Legal Resolving Set in G' .

Let $\{v_{1,\gamma_1}, \dots, v_{k,\gamma_k}\}$ be a k -multicolored independent set in G . We claim that $S := \bigcup_{j \in [m]} \{v_{1,\gamma_1}^j, \dots, v_{k,\gamma_k}^j\}$ is a legal resolving set in G' (of size km). The set S is legal by construction. Since for every $i \in [k]$, and $j \in [m]$, v_{i,γ_i}^j and v_{i,γ_i}^{j+1} are in S ($j + 1$ is modulo m), all the critical pairs in the propagation gadgets are resolved by S , by Lemma 4.4. Since $\{v_{1,\gamma_1}, \dots, v_{k,\gamma_k}\}$ is an independent set in G , there is no $e_j = v_{i,\gamma} v_{i',\gamma'} \in E(G)$, such that $(\gamma, \gamma') = (\gamma_i, \gamma_{i'})$. Thus every critical pair $\{c_j, c'_j\}$ is resolved by S , by Lemma 4.3.

4.2.2 Legal Resolving Set in $G' \Rightarrow k$ -Multicolored Independent Set in G .

Assume that there is a legal resolving set S in G' . For every $i \in [k]$, for every $j \in [m]$, the vertex $v_{i,\gamma(i,j)}^j$ in $V_i^j \cap S$ and the vertex $v_{i,\gamma(i,j+1)}^{j+1}$ in $V_i^{j+1} \cap S$ ($j + 1$ is modulo m) are such that $\gamma(i,j) \leq \gamma(i,j + 1)$, by Lemma 4.4. Thus $\gamma(i, 1) \leq \gamma(i, 2) \leq \dots \leq \gamma(i, m - 1) \leq \gamma(i, m) \leq \gamma(i, 1)$, and $\gamma_i := \gamma(i, 1) = \gamma(i, 2) = \dots = \gamma(i, m - 1) = \gamma(i, m)$. We claim that $\{v_{1,\gamma_1}, \dots, v_{k,\gamma_k}\}$ is a k -multicolored independent set in G . Indeed, there cannot be an edge $e_j = v_{i,\gamma_i} v_{i',\gamma_{i'}} \in E(G)$, since otherwise the critical pair $\{c_j, c_{j'}\}$ is not resolved, by Lemma 4.3.

5 Parameterized Hardness of METRIC DIMENSION/tw

In this section, we produce in polynomial time an instance (G'', k'') of METRIC DIMENSION equivalent to $(G', \mathcal{X}, km, \mathcal{P})$ of k -MULTICOLORED RESOLVING SET. The graph G'' has also pathwidth $O(k)$. Now, an instance is just a graph and an integer. There is no longer \mathcal{X} and \mathcal{P} to constrain and respectively loosen the “resolving set” at our convenience. This creates two issues: (1) the vertices outside the former set \mathcal{X} can now be put in the resolving set, potentially yielding undesired solutions² and (2) our candidate solution (when there is a k -multicolored independent set in G) may not distinguish all the vertices.

5.1 Construction

We settle both issues by attaching new gadgets to G' . Eventually the new graph G'' will contain G' as an induced subgraph. To settle the issue (1), we design a *forced set* gadget. A forced set gadget attached to V_i^j contains two pairs of vertices which are only resolved by vertices of V_i^j . Thus the gadget simulates the action of \mathcal{X} .

There are a few pairs which are not resolved by a solution of k -MULTICOLORED RESOLVING SET. To make sure that all pairs are resolved, we add vertices which need be selected in the resolving set. Technically we could use the previous gadget on a singleton set. But we can make it simpler: we just attach two pendant neighbors, that we then make adjacent, to some chosen vertices. A pair of pendant neighbors are true twins in the whole graph. So we know that at least one of these two vertices have to be in the resolving set. Hence we call that the *forced vertex* gadget, and one of the true twins, a *forced vertex*. It is important that these forced vertices do not resolve any pair of \mathcal{P} . So we can only add pendant twins to vertices themselves not resolving any pair of \mathcal{P} .

² Also, it is now possible to put two or more vertices of the same V_i^j in the resolving set S .

5.1.1 Forced Set Gadget

To deal with the issue (1), we introduce two new pairs of vertices for each V_i^j . The intention is that the only vertices resolving both these pairs simultaneously are precisely the vertices of V_i^j . For any $i \in [k]$ and $j \in [m]$, we add to G' two pairs of vertices $\{p_i^j, q_i^j\}$ and $\{r_i^j, s_i^j\}$, and two gates π_i^j and ρ_i^j . Vertex π_i^j is adjacent to p_i^j and q_i^j , and vertex ρ_i^j is adjacent to r_i^j and s_i^j .

We link $v_{i,1}^j$ to p_i^j and $v_{i,t}^j$ to r_i^j , each by a path of length t . It introduces two new neighbors of $v_{i,1}^j$ and $v_{i,t}^j$ (the brown vertices in Fig. 4). We denote them by tb_i^j and bb_i^j , respectively. The blue and brown vertices are linked to π_i^j and ρ_i^j in the following way. We link tl_i^j and tr_i^j to π_i^j by a private path of length t , and to ρ_i^j by a private path of length $2t - 1$. We link bl_i^j and br_i^j to π_i^j by a private path of length $2t - 1$, and to ρ_i^j by a private path of length t . (Let us clarify that the names of the blue vertices bl_i^j and br_i^j are for “bottom-left” and “bottom-right”, and *not* for “blue” and “brown”.) We link tb_i^j (neighbor of $v_{i,1}^j$) to ρ_i^j by a private path of length $2t - 1$. We link bb_i^j (neighbor of $v_{i,t}^j$) to π_i^j by a private path of length $2t - 1$. Note that the general rule to set the path length is to match the distance between the neighbor in V_i^j and p_i^j (resp. r_i^j). With that in mind we link, if it exists, the *top cyan vertex* $tc_{i,\gamma}^j$ (the one with smallest index γ) neighboring V_i^j to π_i^j with a path of length $\text{dist}(v_{i,\gamma}^j, p_i^j) = t + \gamma - 1$ where $v_{i,\gamma}^j$ is the unique vertex in $N(tc_{i,\gamma}^j) \cap V_i^j$. Observe that with the notations of the previous section $tc_{i,\gamma}^j = e_{i,\gamma}^j$. We also link, if it exists, the *bottom cyan vertex* $bc_{i,\gamma}^j$ (the one with largest index γ) to ρ_i^j with a path of length $\text{dist}(v, r_i^j)$ where v is again the unique neighbor of $bc_{i,\gamma}^j$ in V_i^j .

It can be observed that we only have two paths (and not all six) from the at most three cyan vertices to the gates π_i^j and ρ_i^j . This is where the edges between the cyan vertices will become relevant. See Fig. 4 for an illustration of the forced vertex gadget.

5.1.2 Forced Vertex Gadget

We now deal with the issue (2). By *we add* (or *attach*) a *forced vertex* to an already present vertex v , we mean that we add two adjacent neighbors to v , and that these two vertices remain of degree 2 in the whole graph G'' . Hence one of the two neighbors will have to be selected in the resolving set since they are true twins. We call *forced vertex* one of these two vertices (picking arbitrarily).

For every $i \in [k]$ and $j \in [m]$, we add a forced vertex to the gates nw_i^j and se_i^j of P_i^{j+1} . We also add a forced vertex to each vertex in $N(\{\pi_i^j, \rho_i^j\}) \setminus \{p_i^j, q_i^j, r_i^j, s_i^j\}$. This represents a total of 12 vertices (6 neighbors of π_i^j and 6 neighbors of ρ_i^j). For every $j \in [m]$, we attach a forced vertex to each vertex in $N(g_j) \setminus \{c_j, c'_j\}$. This constitutes 14 neighbors (hence 14 new forced vertices). Therefore we set $k'' := km + 12km + 2km + 14m = 15km + 14m$.

5.1.3 Finishing Strokes and Useful Notations

We use the convention that $P(u, v)$ denotes the path from u to v which was specifically built from u to v . In other words, for $P(u, v)$ to make sense, there should be a point in the construction where we say that we add a (private) path between u and v . For the sake of legibility, $P(u, v)$ may denote either the set of vertices or the induced subgraph. We also denote by $v(u, v)$ the neighbor of u in the path $P(u, v)$. Observe that $P(u, v)$ is a symmetric notation but not $v(u, v)$.

We add a path of length $\text{dist}(v(\pi_i^j, \text{tr}_i^j), \text{sw}_i^j) = t$ between $v(\pi_i^j, \text{tr}_i^j)$ and se_i^j , and a path of length $\text{dist}(v(\pi_i^j, \text{bl}_i^j), \text{ne}_i^{j-1}) = 2t - 1$ between $v(\pi_i^j, \text{bl}_i^j)$ and nw_i^{j-1} . Similarly, we add a path of length $\text{dist}(v(\rho_i^j, \text{tr}_i^j), \text{sw}_i^j) = 2t - 1$ between $v(\rho_i^j, \text{tr}_i^j)$ and se_i^j , and a path of length $\text{dist}(v(\rho_i^j, \text{bl}_i^j), \text{ne}_i^{j-1}) = t$ between $v(\rho_i^j, \text{bl}_i^j)$ and nw_i^{j-1} . We added these four paths so that no forced vertex resolves any critical pair in the propagation gadgets $P_i^{j-1,j}$ and $P_i^{j,j+1}$.

Finally we add an edge between $v(g_j, \text{nw}_i^j)$ and $v(c_j, \text{bc}_i^j)$ whenever V_i^j have exactly three cyan vertices. We do that to resolve the pair $\{v(c_j, \text{tc}_i^j), v(c_j, \text{bc}_i^j)\}$, and more generally every pair $\{x, y\} \in P(c_j, \text{tc}_i^j) \times P(c_j, \text{bc}_i^j)$ such that $\text{dist}(c_j, x) = \text{dist}(c_j, y)$. This finishes the construction of the instance $(G'', k'' := 15 \text{ km} + 14 \text{ m})$ of METRIC DIMENSION (Fig. 5).

5.2 Correctness of the Reduction

The two next lemmas will be crucial in Sect. 5.2.1. The first lemma shows how the forcing set gadget simulates the action of former set \mathcal{X} .

Lemma 5.1 *For every $i \in [k]$ and $j \in [m]$,*

- $\forall v \in V_i^j, v$ resolves both pairs $\{p_i^j, q_i^j\}$ and $\{r_i^j, s_i^j\}$,
- $\forall v \notin V_i^j, v$ resolves at most one pair of $\{p_i^j, q_i^j\}$ and $\{r_i^j, s_i^j\}$,
- $\forall v \notin V_i^j \cup P(v_{i,1}^j, p_i^j) \cup P(v_{i,r}^j, r_i^j) \cup \{q_i^j, s_i^j\}, v$ does not resolve $\{p_i^j, q_i^j\}$ nor $\{r_i^j, s_i^j\}$.

Proof Let $Y := \{\text{tl}_i^j, \text{tr}_i^j, \text{bl}_i^j, \text{br}_i^j\} \cup (X_j \cap N(V_i^j)) \cup (N(\{\pi_i^j, \rho_i^j\}) \setminus \{p_i^j, q_i^j, r_i^j, s_i^j\})$, and recall that $X_j \cap N(V_i^j)$ is the set of cyan vertices neighbors of V_i^j (if they exist). Let us assume that these cyan vertices exist (otherwise the proof is just simpler). In particular, there are at least two cyan neighbors $\text{tc}_i^j, \text{bc}_i^j \in X_j \cap N(V_i^j)$. Let X be the connected component of $G - Y$ containing $\{\pi_i^j, \rho_i^j\}$. For every vertex $u \in \{\text{tl}_i^j, \text{tr}_i^j, \text{bl}_i^j, \text{br}_i^j, \text{tc}_i^j, \text{bc}_i^j\}$, by the way we chose the length of $P(u, \pi_i^j)$ (resp. $P(u, \rho_i^j)$), there is a shortest path from u to p_i^j (resp. r_i^j) that goes through π_i^j (resp. ρ_i^j). Thus $\text{dist}(u, p_i^j) = \text{dist}(u, \pi_i^j) + 1 = \text{dist}(u, q_i^j)$ and $\text{dist}(u, r_i^j) = \text{dist}(u, \rho_i^j) + 1 = \text{dist}(u, s_i^j)$.

Let mc_i^j be the middle cyan vertex if it exists (the one which is not the top nor the bottom one). There is shortest a path from mc_i^j to p_i^j (resp. r_i^j) going via tc_i^j (resp. bc_i^j) and then π_i^j (resp. ρ_i^j). This is where the edges $\text{mc}_i^j \text{tc}_i^j$ and $\text{mc}_i^j \text{bc}_i^j$ are useful. Hence mc_i^j does not resolve $\{p_i^j, q_i^j\}$ nor $\{r_i^j, s_i^j\}$, either. It is direct that no vertex of $N(\{\pi_i^j, \rho_i^j\}) \setminus \{p_i^j, q_i^j, r_i^j, s_i^j\}$ resolves $\{p_i^j, q_i^j\}$ nor $\{r_i^j, s_i^j\}$. Thus no vertex of Y resolves




Fig. 5 The different gadgets attached to V_i^j (recall that $G(e_j)$ is only optionally linked to V_i^j). Gray edges are the edges in the propagation gadgets already depicted in Fig. 3. Black vertices are forced vertices. For the sake of clarity, we did not represent the paths already in the k -MRS-instance from the blue vertices to g_j , and the full forced set gadget of Fig. 4 (it is only symbolized with the dash-dotted edges). A forced vertex is added to each neighbor of the red vertices, except $p_i^j, q_i^j, r_i^j, s_i^j, c_j, c'_j$. Finally we will add four more paths and potentially two edges (see the finishing touches of Sect. 5.1.3) (Color figure online)

any of $\{p_i^j, q_i^j\}$ and $\{r_i^j, s_i^j\}$. Therefore by Lemma 4.1, no vertex outside X resolves any of $\{p_i^j, q_i^j\}$ and $\{r_i^j, s_i^j\}$.

We observe that $X = V_i^j \cup P(v_{i,1}^j, p_i^j) \cup P(v_{i,t}^j, r_i^j) \cup \{\pi_i^j, q_i^j, \rho_i^j, s_i^j\}$. Because of the path from the top brown vertex to p_i^j , vertices of $P(v_{i,1}^j, p_i^j) \setminus \{v_{i,1}^j\} \cup \{q_i^j\}$, which do resolve $\{p_i^j, q_i^j\}$, do not resolve $\{r_i^j, s_i^j\}$. Similarly because of the path from the bottom brown vertex to π_i^j , vertices of $P(v_{i,t}^j, r_i^j) \setminus \{v_{i,t}^j\} \cup \{s_i^j\}$, which do resolve $\{r_i^j, s_i^j\}$, do not resolve $\{p_i^j, q_i^j\}$. Finally for every $u \in V_i^j$, $\text{dist}(u, q_i^j) = \text{dist}(u, p_i^j) + 2$ and $\text{dist}(u, r_i^j) = \text{dist}(u, s_i^j) + 2$. Therefore vertices of V_i^j are the only ones resolving both $\{p_i^j, q_i^j\}$ and $\{r_i^j, s_i^j\}$, while no vertex of $G - X$ resolves any of these pairs. \square

We denote by $f(v)$ the forced vertex attached to a vertex v . For Sect. 5.2.1, we also need the following lemma, which states that the forced vertices do not resolve critical pairs.

Lemma 5.2 *No forced vertex resolves a pair of \mathcal{P} .*

Proof We first show that no critical pair in some P_i^{j+1} is resolved by a forced vertex. We use a similar plan as for the proof of Lemma 4.2. Let $Y := \{nw_i^{j-1}, se_i^{j-1}, nw_i^{j+1}, se_i^{j+1}\} \cup C_e$, where C_e comprises $\{c_j, g_j\}$ if e_j has an endpoint in V_i and $\{c_{j+1}, g_{j+1}\}$ if e_{j+1} has an endpoint in V_i . Let X be the connected

component of $G'' - Y$ containing $P_i^{j,j+1}$. Note that the distances between the vertices of Y and the critical pairs in $P_i^{j,j+1}$ are the same between G' and G'' . Hence as we showed in Lemma 4.2, no vertex of Y resolves a critical pair in $P_i^{j,j+1}$. Thus by Lemma 4.1 no vertex outside X resolves a critical pair in $P_i^{j,j+1}$.

We now check that no forced vertex in X resolves a critical pair in $P_i^{j,j+1}$. We show that every forced vertex in X has a shortest path to $\{nw_i^j, ne_i^j\}$ ending in nw_i^j , and a shortest path to $\{sw_i^j, se_i^j\}$ ending in se_i^j . It is clear for $f(nw_i^j)$ and for $f(se_i^j)$, as well as for all the forced vertices attached to neighbors of g_j (in case e_j has an endpoint in V_i). Indeed recall that the length of $P(g_j, nw_i^j)$ (resp. $P(g_j, se_i^j)$) is four less than the distance to nw_i^j (resp. sw_i^j) ignoring the path $P(g_j, nw_i^j)$ (resp. $P(g_j, se_i^j)$). So the shortest paths from the latter forced vertices go to g_j and then to nw_i^j (resp. se_i^j). Similarly in case e_{j+1} has an endpoint in V_i , the shortest paths from the forced vertices attached to the neighbors of c_{j+1} to $\{nw_i^j, ne_i^j\}$ (resp. $\{sw_i^j, se_i^j\}$) go to g_{j+1} , then to nw_i^{j+1} and nw_i^j (resp. then to se_i^{j+1} and se_i^j).

Note that all the forced vertices attached to neighbors of π_i^j and ρ_i^j (resp. π_i^{j+1} and ρ_i^{j+1}) have a shortest path to $\{nw_i^j, ne_i^j\}$ ending in nw_i^j (resp. to $\{sw_i^j, se_i^j\}$ ending in se_i^j). Finally due to the paths $P(v(\pi_i^j, tr_i^j), se_i^j)$ and $P(v(\rho_i^j, tr_i^j), se_i^j)$, all the forced vertices attached to neighbors of π_i^j and ρ_i^j have a shortest path to $\{sw_i^j, se_i^j\}$ ending in se_i^j . And due to the paths $P(v(\pi_i^{j+1}, bl_i^{j+1}), nw_i^j)$ and $P(v(\rho_i^{j+1}, bl_i^{j+1}), nw_i^j)$, all the forced vertices attached to neighbors of π_i^{j+1} and ρ_i^{j+1} have a shortest path to $\{nw_i^j, ne_i^j\}$ ending in nw_i^j .

We now show that no critical pair $\{c_j, c'_j\}$ is resolved by a forced vertex. We set $Y' := \{tl_i^j, tr_i^j, bl_i^j, br_i^j, tl_{i'}^j, tr_{i'}^j, bl_{i'}^j, br_{i'}^j, nw_i^j, se_i^j, nw_{i'}^j, se_{i'}^j, \pi_i^j, \rho_i^j, \pi_{i'}^j, \rho_{i'}^j\}$, with $e_j \in E(V_i, V_{i'})$, and X' be the connected component of $G'' - Y'$ containing g_j . We showed in Lemma 4.2, and it remains true in G'' , that no vertex of $Y' \setminus \{\pi_i^j, \rho_i^j, \pi_{i'}^j, \rho_{i'}^j\}$ resolves $\{c_j, c'_j\}$. We observe that π_i^j and ρ_i^j have shortest paths to c_j going through g_j (via a vertex of $\{tl_i^j, tr_i^j, bl_i^j, br_i^j\}$). Similarly $\pi_{i'}^j$ and $\rho_{i'}^j$ have shortest paths to c_j going through g_j . Therefore no vertex of $\{\pi_i^j, \rho_i^j, \pi_{i'}^j, \rho_{i'}^j\}$ resolves the pair $\{c_j, c'_j\}$. Hence by Lemma 4.1, no vertex outside X' resolves $\{c_j, c'_j\}$. The only forced vertices in X' are attached to neighbors of g_j , thus they do not resolve $\{c_j, c'_j\}$. □

5.2.1 MD-Instance has a Solution \Rightarrow k -MRS-Instance has a Solution

Let S be a resolving set for the METRIC DIMENSION-instance. We show that $S' := S \cap \bigcup_{i \in [k], j \in [m]} V_i^j$ is a solution for k -MULTICOLORED RESOLVING SET. The set $S \setminus S'$ is made of $14 \text{ km} + 14 \text{ m}$ forced vertices, none of which is in some $V_i^j \cup P(v_{i,1}^j, p_i^j) \cup \{q_i^j\} \cup P(v_{i,r}^j, r_i^j) \cup \{s_i^j\}$. Thus by Lemma 5.1, $S \setminus S'$ does not resolve any pair $\{p_i^j, q_i^j\}$ or $\{r_i^j, s_i^j\}$. Now S' is a set of $k'' - (14 \text{ km} + 14 \text{ m}) = \text{km}$ vertices resolving all the 2 km pairs $\{p_i^j, q_i^j\}$ and $\{r_i^j, s_i^j\}$. Again by Lemma 5.1, this is only possible if $|S' \cap V_i^j| = 1$. Thus S' is a legal set of size $k' = \text{km}$. Let us now check that S' resolves every pair of \mathcal{P} in the graph G' .

By Lemma 5.2, $S \setminus S'$ does not resolve any pair of \mathcal{P} in the graph G'' . Thus S' resolves all the pairs of \mathcal{P} in G'' . Since the distances between V_i^j and the critical pairs

in the edge and propagation gadgets V_i^j is attached to are the same in G' and in G'' , S' also resolves every pair of \mathcal{P} in G' . Thus S' is a solution for the k -MRS-instance.

5.2.2 k -MRS-Instance has a Solution \Rightarrow MD-Instance has a Solution

For every $i \in [k]$, $j \in [m]$, let

$$F_i^j := \bigcup_{u \in \{nw_i^j, se_i^j\} \cup N(\{\pi_i^j, \rho_i^j\}) \setminus \{p_i^j, q_i^j, r_i^j, s_i^j\}} \{f(u)\}, \text{ and}$$

$$F_j := \bigcup_{u \in N(g_j) \setminus \{c_j, c'_j\}} \{f(u)\}.$$

Let S be a solution for k -MULTICOLORED RESOLVING SET. Thus $|S| = km$. Let $F := \bigcup_{i \in [k], j \in [m]} F_i^j \cup \bigcup_{j \in [m]} F_j$. We show that $S' := S \cup F$ is a solution of METRIC DIMENSION. First we observe that $|S'| = km + 14km + 14m = k''$. Since the distances between the sets V_i^j and the critical pairs (of \mathcal{P}) are the same in G' and in G'' , the pairs of \mathcal{P} are resolved by S . In what follows, we show that F resolves all the other pairs. For every $i \in [k]$, $j \in [m]$, we define the subset of vertices:

$$\Pi_i^j := \bigcup_{u \in \{tr_i^j, tl_i^j, br_i^j, bl_i^j, bb_i^j, tc_i^j\}} P(\pi_i^j, u) \cup P(v_{i,1}^j, p_i^j) \cup \{q_i^j\},$$

$$R_i^j := \bigcup_{u \in \{tr_i^j, tl_i^j, br_i^j, bl_i^j, tb_i^j, bc_i^j\}} P(\rho_i^j, u) \cup P(v_{i,t}^j, r_i^j) \cup \{s_i^j\}, \text{ and}$$

$$G_j := \bigcup_{u \in \{tr_i^j, tl_i^j, br_i^j, bl_i^j, tr'_j, bl'_j, br'_j, nw_i^j, se_i^j, nw'_j, se'_j\}} P(g_j, u) \cup E_i^j \cup \overline{E'_j} \cup \{c'_j\}.$$

Informally Π_i^j (R_i^j , G_j , respectively) consists of the vertices on the paths incident to π_i^j (ρ_i^j , g_j , respectively). Our objective is the following result.

Lemma 5.3 *Every vertex in G'' is distinguished by S' .*

We start with the forced vertices and their true twin. We denote by $f'(v)$ the true twin of the forced vertex $f(v)$.

Lemma 5.4 *All the vertices $f(v)$ and $f'(v)$ are distinguished by F .*

Proof Any vertex $f(v)$ is distinguished by being the only vertex at distance 0 of itself $f(v) \in F$. Since $f(v)$ has only two neighbors $f'(v)$ and v , it also resolves every pair $\{f'(v), w\}$ where w is not v . The pair $\{f'(v), v\}$ is resolved by any vertex $f \in F \setminus \{f(v)\}$. Indeed $\text{dist}(f, f'(v)) = \text{dist}(f, v) + 1$. Thus $f'(v)$ is distinguished. \square

In general, to show that all the vertices in a set X are distinguished, we proceed in two steps. First we show that every internal pair of X is resolved. Then, we prove that every pair of $X \times \overline{X}$ is also resolved. Let us recall that \overline{X} is the

complement of x , here $V(G'') \setminus X$. For instance, the two following lemmas show that every vertex of Π_i^j is distinguished by S' .

Lemma 5.5 *Every pair of distinct vertices $x, y \in \Pi_i^j$ is resolved by S' .*

Proof Let U_i^j be the set $\{tl_i^j, tr_i^j, bl_i^j, br_i^j, tc_i^j, bb_i^j\}$. We first consider two vertices $x \neq y \in P(\pi_i^j, u)$, for some $u \in U_i^j$. As $\text{dist}_{G''}(\pi_i^j, u)$ is equal to the length of $P(\pi_i^j, u)$, it holds that $\text{dist}_{G''}(\pi_i^j, x) = \text{dist}_{P(\pi_i^j, u)}(\pi_i^j, x) \neq \text{dist}_{P(\pi_i^j, u)}(\pi_i^j, y) = \text{dist}_{G''}(\pi_i^j, y)$. Without loss of generality, we assume that $\text{dist}(\pi_i^j, x) < \text{dist}(\pi_i^j, y)$. If $x \neq \pi_i^j$, then x and y have distinct distances to $v(\pi_i^j, u)$. Hence $\text{dist}(f(v(\pi_i^j, u)), x) \neq \text{dist}(f(v(\pi_i^j, u)), y)$ and S' resolves $\{x, y\}$. Now if $x = \pi_i^j$, then $f(v(\pi_i^j, u'))$ resolves $\{x, y\}$ for any $u' \in U_i^j \setminus \{u\}$.

Secondly we consider $x \in P(\pi_i^j, u)$ and $y \in P(\pi_i^j, u')$, for some $u \neq u' \in U_i^j$. If $\text{dist}(\pi_i^j, x) \neq 2 + \text{dist}(\pi_i^j, y)$, then $f(v(\pi_i^j, x))$ resolves $\{x, y\}$. Indeed $\text{dist}(f(v(\pi_i^j, x)), x) = \text{dist}(\pi_i^j, x) \neq 2 + \text{dist}(\pi_i^j, y) = \text{dist}(f(v(\pi_i^j, x)), y)$. Else if $\text{dist}(\pi_i^j, x) = 2 + \text{dist}(\pi_i^j, y)$, then $f(v(\pi_i^j, y))$ resolves $\{x, y\}$ (since $\text{dist}(\pi_i^j, y) \neq 2 + \text{dist}(\pi_i^j, x)$).

Two distinct vertices on $P(v_{i,1}^j, p_i^j)$ are resolved by, say, $f(v(\pi_i^j, br_i^j)) \in F$. A vertex of $P(v_{i,1}^j, p_i^j)$ and a vertex of $P(\pi_i^j, u)$, for some $u \in U_i^j$, are resolved by either $f(v(\pi_i^j, u))$ or $f(v(\pi_i^j, u'))$ for a $u' \in U_i^j \setminus \{u\}$. Finally q_i^j and a vertex in $P(v_{i,1}^j, p_i^j) \setminus \{p_i^j\}$ are resolved by, say, $f(v(\pi_i^j, br_i^j))$, whereas q_i^j and a vertex in $P(p_i^j, u)$ is resolved by either $f(v(\pi_i^j, u))$ or $f(v(\pi_i^j, u'))$ for a $u' \in U_i^j \setminus \{u\}$. Therefore every pair of distinct vertices in Π_i^j is resolved by F , except $\{p_i^j, q_i^j\}$ which is resolved by S . \square

Lemma 5.6 *Every pair $\{x, y\} \in \Pi_i^j \times \overline{\Pi_i^j}$ is resolved by F .*

Proof Again let U_i^j be the set $\{tl_i^j, tr_i^j, bl_i^j, br_i^j, tc_i^j, bb_i^j\}$. We first assume x is in $P(\pi_i^j, u)$ for some $u \in U_i^j \setminus \{tr_i^j, bl_i^j\}$. Let y be a vertex of Π_i^j such that $\text{dist}(f(v(\pi_i^j, u)), x) = \text{dist}(f(v(\pi_i^j, u)), y)$, otherwise $f(v(\pi_i^j, u))$ already resolves $\{x, y\}$. Every shortest path from $f(v(\pi_i^j, u))$ to y go through π_i^j . One can observe that there is a $u' \in U_i^j \setminus \{u\}$ such that $f(v(\pi_i^j, u'))$ has a shortest path also going through π_i^j . Hence $f(v(\pi_i^j, u'))$ has the same distance to y (as $f(v(\pi_i^j, u))$) but a larger distance to x . Hence $f(v(\pi_i^j, u'))$ resolves $\{x, y\}$.

We now consider an $x \in P(\pi_i^j, u)$ for some $u \in \{tr_i^j, bl_i^j\}$. Again let y be a vertex of Π_i^j such that $\text{dist}(f(v(\pi_i^j, u)), x) = \text{dist}(f(v(\pi_i^j, u)), y)$. If all the shortest paths of $f(v(\pi_i^j, u))$ to y goes through π_i^j , we conclude as in the previous paragraph. So they go through $P(v(\pi_i^j, u), se_i^j)$ (if $u = tr_i^j$) or $P(v(\pi_i^j, u), nw_i^{j-1})$ (if $u = bl_i^j$). Since $\text{dist}(f(v(\pi_i^j, u)), x) \leq 2t - 1$, it also holds that $\text{dist}(f(v(\pi_i^j, u)), y) \leq 2t - 1$. The path $P(v(\pi_i^j, tr_i^j), se_i^j)$ has length t and the path $P(v(\pi_i^j, bl_i^j), nw_i^{j-1})$ has length $2t - 1$. Therefore one of $f(se_i^j), f(se_i^{j-1}), f(nw_i^j), f(nw_i^{j-1})$ resolves $\{x, y\}$.

We now assume x is in $P(v_{i,1}^j, p_i^j) \cup \{q_i^j\}$ and $y \in \Pi_i^j$. Then $f(v(\pi_i^j, br_i^j))$ resolves $\{x, y\}$ if y is not in the path $P(v(\pi_i^j, tr_i^j), se_i^j)$ or $P(v(\pi_i^j, u), nw_i^{j-1})$. Otherwise at least one of $f(v(\pi_i^j, br_i^j)), f(v(\pi_i^j, tr_i^j)), f(v(\pi_i^j, bl_i^j))$ resolves $\{x, y\}$. In conclusion, every pair of vertices $\{x, y\} \in \Pi_i^j \times \overline{\Pi_i^j}$ is resolved by F . \square

Lemmas 5.5 and 5.6 prove that every vertex in Π_i^j is distinguished by S' . Using the same arguments, we get symmetrically that every vertex of R_i^j is distinguished by S' .

Lemma 5.7 *All the vertices in the paths $P(v(\pi_i^j, tr_i^j), se_i^j)$, $P(v(\rho_i^j, tr_i^j), se_i^j)$, $P(v(\pi_i^j, bl_i^j), nw_i^{j-1})$, $P(v(\rho_i^j, bl_i^j), nw_i^{j-1})$ are distinguished by F .*

Proof Any vertex $x \in P(v(\pi_i^j, tr_i^j), se_i^j)$ is uniquely determined by its distances to $f(se_i^j)$, $f(se_i^{j-1})$, and $v(\pi_i^j, tr_i^j)$. Any vertex $x \in P(v(\pi_i^j, bl_i^j), nw_i^{j-1})$ is uniquely determined by its distances to $f(nw_i^j)$, $f(nw_i^{j-1})$, and $v(\pi_i^j, bl_i^j)$. The two other cases are symmetric. □

So far we showed that the vertices added in the forced set and forced vertex gadgets are all distinguished. We now focus on the vertices in propagation gadgets. Let $\Delta_i := A_i^j \cup \{nw_i^j, ne_i^j, sw_i^j, se_i^j\}$.

Lemma 5.8 *Every pair of distinct vertices $x, y \in \Delta_i^j$ is resolved by S' .*

Proof Since the distances between vertices of V_i^j and vertices of Δ_i^j are the same between G' and G'' , S resolves all the critical pairs $\{a_{i,\gamma}^j, \alpha_{i,\gamma}^j\}$. Thus we turn our attention to the pairs which are not critical pairs. Since $\text{dist}(nw_i^j, a_{i,\gamma}^j) = \gamma$ and $\text{dist}(nw_i^j, \alpha_{i,\gamma}^j) = \gamma$, every pair $\{a_{i,\gamma}^j, a_{i,\gamma'}^j\}$, $\{a_{i,\gamma}^j, \alpha_{i,\gamma'}^j\}$, or $\{\alpha_{i,\gamma}^j, \alpha_{i,\gamma'}^j\}$, with $\gamma \neq \gamma'$ is resolved by $f(nw_i^j)$.

Gate nw_i^j (resp. se_i^j) and any other vertex in Δ_i^j is resolved by $f(nw_i^j)$ (resp. $f(se_i^j)$). Gate ne_i^j (resp. sw_i^j) is resolved from any vertex of $\Delta_i^j \setminus \{a_{i,1}^j, \alpha_{i,1}^j\}$ (resp. $\Delta_i^j \setminus \{a_{i,t}^j, \alpha_{i,t}^j\}$) by $f(nw_i^j)$ (resp. $f(se_i^j)$). Finally, ne_i^j (resp. sw_i^j) and a vertex of $\{a_{i,1}^j, \alpha_{i,1}^j\}$ (resp. $\{a_{i,t}^j, \alpha_{i,t}^j\}$) is resolved by $f(se_i^j)$ (resp. $f(nw_i^j)$). □

Now when we check that a pair made of a vertex in Δ_i^j and a vertex outside Δ_i^j is resolved, we can further assume that the second vertex is not in some $\Pi_i^j \cup R_i^j$ since we already showed that these vertices were distinguished.

Lemma 5.9 *Every pair $\{x, y\} \in \Delta_i^j \times \overline{\Delta_i^j}$ is resolved by S' .*

Proof We may assume that y is not a vertex that was previously shown distinguished. Thus y is not in some $\Pi_i^j \cup R_i^j$ nor in a path of Lemma 5.7. Then we claim that the pair $\{x, y\}$ is resolved by at least one of $f(se_i^j)$, $f(se_i^{j-1})$, $f(se_i^{j+1})$, $f(nw_i^j)$. Indeed assume that $f(se_i^j)$ does not resolve $\{x, y\}$, and consider a shortest path from $f(se_i^j)$ to y . Either this shortest path goes through se_i^{j-1} (resp. se_i^{j+1}), and in that case $f(se_i^{j-1})$ (resp. $f(se_i^{j+1})$) resolves $\{x, y\}$. Either it takes the path to g_j (if e_j has an endpoint in V_i) or to tl_i^{j+1} , and then $f(nw_i^j)$ resolves $\{x, y\}$. Or it takes a path to V_i^j , and then $f(se_i^{j-1})$ resolves $\{x, y\}$. □

Lemmas 5.8 and 5.9 show that that every vertex in Δ_i^j is distinguished by S' . The common neighbor of se_i^{j-1} and tl_i^j is distinguished by $\{f(se_i^{j-1}), f(v(\pi_i^j, tl_i^j))\}$. We are

now left with showing that the vertices in the edge gadgets, in the sets V_i^j , and in the paths incident to the edge gadgets, are distinguished.

Lemma 5.10 *Every pair of distinct vertices $x, y \in G_j$ is resolved by S' .*

Proof Let $v_{i,\gamma}$ and $v_{i',\gamma'}$ be the two endpoints of e_j , and $U_i^j := \{tl_i^j, tr_i^j, bl_i^j, br_i^j, tl_{i'}^j, tr_{i'}^j, bl_{i'}^j, br_{i'}^j, se_i^j, nw_i^j, se_{i'}^j, nw_{i'}^j, v_{i,\gamma}^j, v_{i',\gamma'}^j\}$. Every pair in $\bigcup_{u \in U_i^j} P(g_j, u)$ is resolved. Indeed, similarly to Lemma 5.5, two distinct vertices x, y on a path $P(g_j, u)$ ($u \in U_i^j$) are resolved by $f(v(g_j, u))$, while two vertices on distinct paths $P(g_j, u)$ and $P(g_j, u')$ ($u \neq u' \in U_i^j$) are resolved by at least one of $f(v(g_j, u))$ and $f(v(g_j, u'))$.

We now show that any pair in $\Gamma_i^j := E_i^j \cup E_{i'}^j \setminus \{P(g_j, v_{i,\gamma}^j), P(g_j, v_{i',\gamma'}^j)\}$ is resolved. Two distinct vertices $x, y \in \Gamma_i^j$ are resolved by, say, $f(v(g_j, se_i^j))$ if they are on the same path, or more generally if they have different distances to c_j . Thus let us assume that x and y are at the same distance from c_j . If $x \in E_i^j$ and $y \in E_{i'}^j$ (or vice versa) then the pair $\{x, y\}$ is resolved by the vertex in $S \cap V_i^j$ or the vertex in $S \cap V_{i'}^j$. If $x \neq y \in E_i^j$ (resp. $\in E_{i'}^j$), then $\{x, y\}$ is resolved by $f(v(g_j, nw_i^j))$ (resp. $f(v(g_j, nw_{i'}^j))$). This is the reason why we added an edge between $v(g_j, nw_i^j)$ and $v(c_j, bc_i^j)$ (recall Sect. 5.1.3).

We now consider pairs $\{x, y\}$ of $\bigcup_{u \in U_i^j} P(g_j, u) \times \Gamma_i^j$. Any of these pairs are resolved by at least one of $f(v(g_j, u))$, $f(v(g_j, u'))$, $f(v(g_j, nw_i^j))$, $f(v(g_j, nw_{i'}^j))$, where x is on the path $P(c_j, u)$ and u' is any vertex in $U_i^j \setminus \{u, nw_i^j, nw_{i'}^j\}$. Finally c_j' is distinguished from all the other vertices in G'' but c_j by the forced vertices attached to the neighbors of g_j .

Thus every pair $\{x, y\}$ in G_j is resolved by F , except $\{c_j, c_j'\}$ which is resolved by S . □

Lemma 5.11 *Every pair $\{x, y\} \in G_j \times \overline{G_j}$ is resolved by F .*

Proof Consider an arbitrary pair $\{x, y\} \in G_j \times \overline{G_j}$. We can assume that x is not c_j' , and that y is in one different $G_{j'}$ or in one $V_{j''}^j$ (since we already showed that the other vertices are distinguished). Again let $v_{i,\gamma}$ and $v_{i',\gamma'}$ be the two endpoints of e_j , and $U_i^j := \{tl_i^j, tr_i^j, bl_i^j, br_i^j, tl_{i'}^j, tr_{i'}^j, bl_{i'}^j, br_{i'}^j, se_i^j, nw_i^j, se_{i'}^j, nw_{i'}^j, v_{i,\gamma}^j, v_{i',\gamma'}^j\}$. If x is on a path $P(g_j, u)$, then at least one of $f(v(g_j, u))$ and $f(v(g_j, u'))$, with u' being any vertex in $U_i^j \setminus \{u\}$, resolves $\{x, y\}$. If instead x is on a path $P(c_j, u)$ with $u \in \{v_{i,\gamma-1}^j, v_{i,\gamma+1}^j, v_{i',\gamma'-1}^j, v_{i',\gamma'+1}^j\}$, then at least one of $f(v(g_j, nw_i^j))$, $f(v(g_j, nw_{i'}^j))$, $f(v(g_j, u'))$, with u' being any vertex in U_i^j , resolves $\{x, y\}$. □

Lemmas 5.10 and 5.11 show that every vertex in G_j is distinguished by S' . We finally show that the vertices in V_i^j are distinguished. A pair of distinct vertices $x, y \in V_i^j$ is resolved by $f(nw_{i'}^j)$. We thus consider a pair $\{x, y\} \in V_i^j \times V_{i'}^j$. We can further assume that y is in some $V_{i''}^j$, since all the other vertices have already been shown distinguished. Then $\{x, y\}$ is resolved by at least one of $f(nw_i^j)$, $f(nw_{i'}^j)$, the vertex in $S \cap V_i^j$, and the vertex in $S \cap V_{i''}^j$. This finishes the proof of Theorem 5.3. Thus S' is a solution of the METRIC DIMENSION-INSTANCE.

The reduction is correct and it takes polynomial-time in $|V(G)|$ to compute G'' . The maximum degree of G'' is 16. It is the degree of the vertices g_j (nw_i^j and se_i^j have degree at most 11, π_i^j and ρ_i^j have degree 8, and the other vertices have degree at most 5). The last element to establish Theorem 3.1 is to show that $\text{pw}(G'')$ is in $O(k)$. Then solving METRIC DIMENSION on constant-degree graphs in time $f(\text{pw})n^{o(\text{pw})}$ could be used to solve k -MULTICOLORED INDEPENDENT SET in time $f(k)n^{o(k)}$, disproving the ETH.

5.3 G'' has Pathwidth $O(k)$

We use the pathwidth characterization of Kirousis and Papadimitriou [19] mentioned in the preliminaries, and give a strategy with $O(k)$ searchers cleaning all the edges of G'' . A basic and useful fact is that the searching number of a path is two.

Lemma 5.12 *Two searchers are enough to clean a path $u_1u_2 \dots u_n$.*

Proof We place two searchers at u_1 and u_2 . This cleans the edge u_1u_2 . Then we move the searcher in u_1 to u_3 . This cleans u_2u_3 (while u_1u_2 remains clean). Then we move the searcher in u_2 to u_4 , and so on. □

Lemma 5.13 $\text{pw}(G'') \leq 90k + 83$.

Proof For every $j \in [m]$, let $S_j := N[g_j] \cup X_j \cup \bigcup_{i \in [k]} N[\{v_{i,1}^j, v_{i,2}^j, \pi_i^j, \rho_i^j\}] \cup \{nw_i^j, ne_i^j, sw_i^j, se_i^j\}$. We notice that $|S_j| \leq 17 + 6 + 30k + 4 = 30k + 27$. Another important observation is that $S_1 \cup S_j$ disconnects the first j columns of G'' from the rest of G'' . Finally the connected components $G'' - (S_j \cup S_{j+1})$ that are not the main component (i.e., containing more than half of the graph if $m \geq 4$) are all paths.

We now suggest the following cleaning strategy with at most $90k + 83$ searchers. We place one searcher at each vertex of $S_1 \cup S_2 \cup S_3$. This requires $90k + 81$ searchers. By Lemma 5.12, with two additional searchers we clean all the connected components of $G'' - (S_1 \cup S_2 \cup S_3)$ that are paths. We then move all the searchers from S_2 to S_4 , and clean all the connected components of $G'' - (S_1 \cup S_3 \cup S_4)$ that are paths. Since $S_1 \cup S_3$ is a separator, the edges that were cleaned during the first phase are not recontaminated when we move from S_2 to S_4 . We then move the searchers of S_3 to S_5 , and so on. Eventually the searchers reach $S_1 \cup S_{m-1} \cup S_m$, and the last contaminated edges are cleaned. □

6 Perspectives

The main remaining open question is whether or not METRIC DIMENSION can be solved in polynomial time on graphs with constant treewidth. In the parameterized complexity language, now we know that MD/tw is W[1]-hard, is it in XP or

paraNP-hard? We believe that the tools and ideas developed in this paper could help answering this question negatively. The FPT algorithm of Belmonte et al. [3] also implies that METRIC DIMENSION is FPT with respect to $tl + k$ where k is the size of the resolving set, due to the bound $\Delta \leq 2^k + k - 1$ [18]. What about the parameterized complexity of METRIC DIMENSION with respect to $tw + k$? We conjecture that this problem is W[1]-hard as well, and once again, treewidth will contrast with tree-length.

It appears that bounded connected treewidth or tree-length is significantly more helpful than the mere bounded treewidth when it comes to solving MD. We wish to ask for the parameterized complexity of METRIC DIMENSION with respect to ctw only (on graphs with arbitrarily large degree). Finally, it would be interesting to determine if planarity can sometimes help to compute a metric basis. Therefore we also ask all the above questions in planar graphs.

Funding Open Access funding provided by University of Bergen (incl Haukeland University Hospital).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Barbero, F., Isenmann, L., Thiebaut, J.: On the distance identifying set meta-problem and applications to the complexity of identifying problems on graphs. In: 13th International Symposium on Parameterized and Exact Computation, IPEC 2018, August 20–24, 2018, Helsinki, Finland, pp. 10:1–10:14 (2018). <https://doi.org/10.4230/LIPIcs.IPEC.2018.10>
2. Beerliova, Z., Eberhard, F., Erlebach, T., Hall, A., Hoffmann, M., Mihalák, M., Ram, L.S.: Network discovery and verification. *IEEE J. Sel. Areas Commun.* **24**(12), 2168–2181 (2006). <https://doi.org/10.1109/JSAC.2006.884015>
3. Belmonte, R., Fomin, F.V., Golovach, P.A., Ramanujan, M.S.: Metric dimension of bounded tree-length graphs. *SIAM J. Discrete Math.* **31**(2), 1217–1243 (2017). <https://doi.org/10.1137/16M1057383>
4. Chartrand, G., Eroh, L., Johnson, M.A., Oellermann, O.: Resolvability in graphs and the metric dimension of a graph. *Discrete Appl. Math.* **105**(1–3), 99–113 (2000). [https://doi.org/10.1016/S0166-218X\(00\)00198-0](https://doi.org/10.1016/S0166-218X(00)00198-0)
5. Cygan, M., Fomin, F.V., Kowalik, L., Lokshtanov, D., Marx, D., Pilipczuk, M., Pilipczuk, M., Saurabh, S.: *Parameterized Algorithms*. Springer, Berlin (2015). <https://doi.org/10.1007/978-3-319-21275-3>
6. Díaz, J., Pottonen, O., Serna, M.J., van Leeuwen, E.J.: Complexity of metric dimension on planar graphs. *J. Comput. Syst. Sci.* **83**(1), 132–158 (2017). <https://doi.org/10.1016/j.jcss.2016.06.006>
7. Downey, R.G., Fellows, M.R.: *Fundamentals of Parameterized Complexity*. Texts in Computer Science. Springer, Berlin (2013). <https://doi.org/10.1007/978-1-4471-5559-1>
8. Eppstein, D.: Metric dimension parameterized by max leaf number. *J. Graph. Algorithms Appl.* **19**(1), 313–323 (2015). <https://doi.org/10.7155/jgaa.00360>
9. Epstein, L., Levin, A., Woeginger, G.J.: The (weighted) metric dimension of graphs: hard and easy cases. *Algorithmica* **72**(4), 1130–1171 (2015). <https://doi.org/10.1007/s00453-014-9896-2>
10. Fernau, H., Heggenes, P., van't Hof, P., Meister, D., Saei, R.: Computing the metric dimension for chain graphs. *Inf. Process. Lett.* **115**(9), 671–676 (2015). <https://doi.org/10.1016/j.ipl.2015.04.006>

11. Foucaud, F., Mertzios, G.B., Naserasr, R., Parreau, A., Valicov, P.: Identification, location-domination and metric dimension on interval and permutation graphs. II. Algorithms and complexity. *Algorithmica* **78**(3), 914–944 (2017). <https://doi.org/10.1007/s00453-016-0184-1>
12. Garey, M.R., Johnson, D.S.: *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman, New York (1979)
13. Harary, F., Melter, R.A.: On the metric dimension of a graph. *Ars Combin.* **2**(191–195), 1 (1976)
14. Hartung, S., Nichterlein, A.: On the parameterized and approximation hardness of metric dimension. In: *Proceedings of the 28th Conference on Computational Complexity, CCC 2013*, K. Io Alto, California, USA, 5–7 June, 2013, pp. 266–276 (2013). <https://doi.org/10.1109/CCC.2013.36>
15. Hoffmann, S., Wanke, E.: Metric dimension for Gabriel unit disk graphs is NP-complete. In: *Algorithms for Sensor Systems, 8th International Symposium on Algorithms for Sensor Systems, Wireless Ad Hoc Networks and Autonomous Mobile Entities, ALGOSENSORS 2012*, Ljubljana, Slovenia, September 13–14, 2012. Revised Selected Papers, pp. 90–92 (2012). https://doi.org/10.1007/978-3-642-36092-3_10
16. Hoffmann, S., Elterman, A., Wanke, E.: A linear time algorithm for metric dimension of cactus block graphs. *Theor. Comput. Sci.* **630**, 43–62 (2016). <https://doi.org/10.1016/j.tcs.2016.03.024>
17. Impagliazzo, R., Paturi, R., Zane, F.: Which problems have strongly exponential complexity? *J. Comput. Syst. Sci.* **63**(4), 512–530 (2001)
18. Khuller, S., Raghavachari, B., Rosenfeld, A.: Landmarks in graphs. *Discrete Appl. Math.* **70**(3), 217–229 (1996). [https://doi.org/10.1016/0166-218X\(95\)00106-2](https://doi.org/10.1016/0166-218X(95)00106-2)
19. Kirousis, L.M., Papadimitriou, C.H.: Interval graphs and searching. *Discrete Math.* **55**(2), 181–184 (1985). [https://doi.org/10.1016/0012-365X\(85\)90046-9](https://doi.org/10.1016/0012-365X(85)90046-9)
20. Lokshtanov, D., Marx, D., Saurabh, S.: Lower bounds based on the exponential time hypothesis. *Bull. EATCS* **105**, 41–72 (2011)
21. Pietrzak, K.: On the parameterized complexity of the fixed alphabet shortest common supersequence and longest common subsequence problems. *J. Comput. Syst. Sci.* **67**(4), 757–771 (2003). [https://doi.org/10.1016/S0022-0000\(03\)00078-3](https://doi.org/10.1016/S0022-0000(03)00078-3)
22. Slater, P.J.: Leaves of trees. *Congr. Numer.* **14**(549–559), 37 (1975)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.