

Additional file 1: Most common medicines used against nausea on 1st and 2nd ATC level according to timing of use in pregnancy among women with nausea in pregnancy (n=6701)*

Anatomical Therapeutic Chemical (ATC) classification index 1st and 2nd levels		Anytime during pregnancy n (%)	1st trimester n (%)	2nd trimester n (%)	3rd trimester n (%)
A	Alimentary tract and metabolism	564 (8.4)	448 (6.7)	249 (3.7)	141 (2.1)
A02	Drugs for acid related disorders	176 (2.6)	105 (1.6)	83 (1.2)	78 (1.2)
A03	<i>Drugs for functional gastrointestinal disorders</i>	316 (4.7)	277 (4.1)	137 (2.0)	52 (0.8)
A04	Antiemetics and antinauseants	90 (1.3)	81 (1.2)	47 (0.7)	22 (0.3)
A05	Bile and liver therapy	12 (0.2)	9 (0.1)	6 (0.1)	1 (0.0)
A07	Antidiarrheals, intestinal antiinflammatory/antiinfective agents	17 (0.3)	13 (0.2)	5 (0.1)	2 (0.0)
N	Nervous system	115 (1.7)	81 (1.2)	74 (1.1)	34 (0.5)
N02	<i>Analgesics</i>	83 (1.2)	54 (0.8)	52 (0.8)	24 (0.4)
N05	Psycholeptics	32 (0.5)	27 (0.4)	22 (0.3)	10 (0.2)
R	Respiratory system	620 (9.3)	536 (8.0)	319 (4.8)	111 (1.7)
R06	<i>Antihistamines for systemic use</i>	613 (9.2)	533 (8.0)	316 (4.7)	108 (1.6)
-	Unspecified	82 (1.2)	57 (0.9)	46 (0.7)	17 (0.3)
Total medicine use (any ATC)		1201 (17.9)^a	965 (14.4)	577 (8.6)	254 (3.8)
Total vitamin and/or mineral use (ATC-class: A11-13, B03 and Unspecified supplement)		106 (1.6)	90 (1.3)	42 (0.6)	9 (0.1)

*Only medicine groups used by more than 10 women are presented. Exposure timing is defined as follows: 1st trimester (gestational weeks 0-12), 2nd trimester (gestational week 13-24), 3rd trimester (gestational week 25 and up to childbirth).

^aNumbers do not add up to total due to some women may have used more than one medicine