

Additional data file 4. Clinicopathological features of patients (“GRONINGEN”) stratified according to *HOTAIR* expression.

Characteristics	<i>HOTAIR</i> RNA expression		P Value *
	negative (n = 58)	positive (n = 117)	
Age			0.262
≤ 62.7 yrs. (median age)	87	25	62
> 62.7 yrs. (median age)	88	33	55
FIGO			0.665
III	147	50	97
IV	28	8	20
Tumor grade			0.483
I/II	59	17	42
III	97	34	63
Unknown	19	7	12
Histology			0.988
serous cancer	118	39	79
mucinous cancer	15	5	10
endometrioid cancer	14	5	9
clear cell cancer	8	2	6
Others	20	7	13
Residual disease after surgery			0.161
no residual disease	29	8	21
residual disease ≤ 2cm	38	18	20
residual disease > 2cm	97	31	66
Unknown	11	1	10
Chemotherapy**			0.999
Cisplatin based	18	6	12
Carboplatin based	157	52	105
Survival status (5 years)			0.066
Alive	61	26	35
Dead	114	32	82

* P values were calculated with the use of the Chi square test.

** 98 and 59 patients received Carboplatin in combination with Cyclophosphamide and Paclitaxel, respectively; 10 and 8 patients received Cisplatin in combination with Cyclophosphamide and Paclitaxel, respectively.