



MEETING ABSTRACT

Open Access

# The MHC-II transactivator CIITA, a viral restriction factor inhibiting the replication of Human T-Cell Lymphotropic Virus Type 1

Giovanna Tosi<sup>1\*</sup>, Greta Forlani<sup>1</sup>, Vibeke Andresen<sup>2</sup>, Marco Turci<sup>3</sup>, Umberto Bertazzoni<sup>3</sup>, Genoveffa Franchini<sup>4</sup>, Roberto S Accolla<sup>1</sup>

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses  
Leuven and Gembloux, Belgium. 5-8 June 2011

Human T-cell Lymphotropic Virus type-1 (HTLV-1) is the causative agent of an aggressive malignancy of CD4 + T lymphocytes. It is believed that the viral transactivator Tax-1 is a major player in T-cell transformation. Thus, targeting Tax-1 protein is regarded as a possible strategy to arrest viral replication and ultimately to counteract neoplastic transformation.

Here, we demonstrated that CIITA, the master regulator of MHC class II gene transcription, inhibits HTLV-1 replication by blocking the transactivating function of Tax-1. Co-immunoprecipitation experiments have shown that CIITA and Tax-1 physically interact in vivo and that the first 108 amino acids of Tax-1 were necessary for this binding. Two adjacent regions (1-252 and 253-410) of CIITA bound independently to Tax-1, but only region 1-252 mediated Tax-1 inhibition, in agreement with the fact that CIITA residues from positions 64-124 were required to block Tax-1 transactivation. CIITA inhibitory action on Tax-1 function correlated with the nuclear localization of CIITA and was independent of the transcription factor NF-YB, previously involved in CIITA-mediated inhibition of Tax-2 of HTLV-2, a virus with still elusive pathogenic action. Furthermore, CIITA severely impaired the physical and functional interaction of Tax-1 with the cellular co-activator PCAF, which is required for the optimal activation of HTLV-1 promoter. Accordingly, the over-expression of PCAF restored Tax-1-dependent transactivation of the viral LTR promoter inhibited by CIITA.

These findings strongly support our original observation that CIITA, beside increasing the antigen-presenting function for pathogen antigens, acts as an endogenous restriction factor against human retroviruses by blocking virus replication and spreading.

#### Author details

<sup>1</sup>Department of Experimental Medicine, University of Insubria, Varese, 21100, Italy. <sup>2</sup>Institute of Medicine, Hematology section, University of Bergen, Bergen, Norway. <sup>3</sup>Department of Life and Reproduction Sciences, Section of Biology and Genetics, University of Verona, Verona, Italy. <sup>4</sup>Animal Models and Retroviral Vaccines Section, National Cancer Institute, Bethesda, MD, USA.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A171

**Cite this article as:** Tosi *et al.*: The MHC-II transactivator CIITA, a viral restriction factor inhibiting the replication of Human T-Cell Lymphotropic Virus Type 1. *Retrovirology* 2011 **8**(Suppl 1):A171.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [giovanna.tosi@yahoo.com](mailto:giovanna.tosi@yahoo.com)

<sup>1</sup>Department of Experimental Medicine, University of Insubria, Varese, 21100, Italy

Full list of author information is available at the end of the article